

**NEOVA DNA DAMAGE CONTROL - SILC SHEER BROAD SPECTRUM SPF 40-  
octinoxate, titanium dioxide, zinc oxide emulsion  
Pharma Cosmetics**

-----

**Active Ingredients**

Octinoxate 3.0%, Titanium Dioxide 3.0%, Zinc Oxide 9.0%

**Purpose**

Sunscreen

**Uses**

- Helps prevent sunburn
- If used as directed and with other sun protection measures (**see Directions**), decreases the risk of skin cancer and early skin aging caused by the sun.

**Warnings**

For external use only

**Do not use** on damaged or broken skin.

**When using this product** keep out of eyes. Rinse with water to remove.

**Stop use and ask a doctor if** rash occurs.

**Keep out of reach of children.** If product is swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

- Apply liberally 15 minutes before sun exposure.
- Reapply:

°After 80 minutes of swimming or sweating.

°Immediately after towel drying.

°At least every two hours.

- Children under 6 months of age: ask a doctor.
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

°Limit time in the sun, especially from 10 a.m. - 2 p.m.

°Wear long-sleeve shirts, pants, hats, and sunglasses.

## **Inactive Ingredients**

Alumina, Butylene Glycol, Citric Acid, Cyclopentasiloxane, Dimethicone, Dimethicone /PEG-10/15 Crosspolymer, Ergothioneine, HDI/Trimethylol Hexyllactone Crosspolymer, Iodopropynyl Butylcarbamate,

Iron Oxides, Lauryl PEG-9 Polymethylsiloxyethyl Dimethicone, Lecithin, Hydrogen Dimethicone, Micrococcus

Lysate, Octyldodecyl Neopentanoate, Phenoxyethanol, Plankton Extract, Purified Water, Silica, Sodium

Chloride, Sodium Hydroxide, Triethoxycaprylylsilane.

## **Other Information**

Protect this product from excessive heat and direct sun.

## **Questions or Comments?**

Call toll free 1-888-966-1010.

## **Product label**

SILC SHEER  
2.0  
BROAD SPECTRUM  
SPF 40

**Drug Facts**

Active Ingredients	Purpose
Octinoxate 3.0%	Sunscreen
Titanium Dioxide 3.0%	Sunscreen
Zinc Oxide 8.0%	Sunscreen

**Uses** • Helps prevent sunburns. • If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun.

**Warnings** For external use only

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs.

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

• Apply liberally 15 minutes before sun exposure.

• Reapply:

• After 80 minutes of sweating or sweating.

• Immediately after towel drying.

• At least every 2 hours.

• Children under 6 months of age: ask a doctor.

• Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging.

To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

• Limit time in the sun, especially from 10 a.m. - 2 p.m.

• Wear long-sleeved shirts, pants, hats, and sunglasses.

**Inactive Ingredients**

Alumina, Isotylene Glycol, Citric Acid, Cyclopentasiloxane, Dimethicone, Dimethicone / PEG-10/15 Copolymer, Ergosterol, HDI/Hexamethylene Diisocyanate Crosspolymer, Isododecyl Isitylcarbamate, Iron Oxides, Lauryl PEG-9 Polymethylsilyloxyethyl Dimethicone, Lecithin, Hydrogen Dimethicone, Microcrystalline Lysine, Octyldecyl Neopentanoate, Phenoxyethanol, Plantain Extract, Purified Water, Silica, Sodium Chloride, Sodium Hydroxide, Tetrahydroxyethylsiloxane.

**Other Information**

Protect this product from excessive heat and direct sun.

**Questions or Comments?**

Call toll free 1-888-888-1110.

Dermatologist tested + Hypoallergenic.

The future is SILC SHEER 2.0. Now smarter, brighter and full of great ideas.

- + 2 physical UV filters for complete, broad spectrum photo protection
- + 2 levels of advanced skin reformers (DNA Repair enzymes + L-ergothioneine)
- + 2 light-diffusing factors to illuminate and reduce visible imperfections. On contact.

Pharmia Cosmetics, Inc.  
Oradell, NJ 07648  
95 bis rue de Rocher  
75008 Paris, France  
1.888.888.1110 • www.neoava.com

U.S. Patent No. 8,331,159  
Made in the USA  
N03001 | Allsize

Follow Us:    

NEOVA<sup>®</sup> SmartScience. SmartSunscreens.

DNA DAMAGE CONTROL

SILC SHEER 2.0  
COLOR CORRECTING PHOTO FINISH TINT

 SmartSunscreens Technology

BROAD SPECTRUM SPF 40

WATER RESISTANT (80 MINUTES)

74 mL e 2.5 fl. oz.


Award-Winning DNA Repair Technology  
The same DNA repair science awarded the 2015 Nobel Prize in Chemistry is used in NEOVA SmartSunscreens.

NEOVA DNA Damage Control (SILC SHEER 2.0), with a skin perfecting, natural tint for the face, provides a new level of protection like no other: SmartSunscreens Technology delivers advanced defense against the aging and burning effects of UVA/UVB rays and helps improve the visible signs of DNA skin damage from the sun. (Oil-free formula)

**Complete Photoprotection In One Advanced Formula**

- + Transparent Zinc and Micronized Titanium Dioxide, highly effective physical UV filters, defend against broad-spectrum UVA/UVB rays
- + Liposome-encapsulated DNA Repair enzymes restrict and reduce the signs of sun-inflicted damage
- + L-ergothioneine, a super antioxidant, suppresses free-radical damage





## NEOVA DNA DAMAGE CONTROL - SILC SHEER BROAD SPECTRUM SPF 40

octinoxate, titanium dioxide, zinc oxide emulsion

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:72251-001
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>OCTINOXATE</b> (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	3 g in 100 mL
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	3 g in 100 mL
<b>ZINC OXIDE</b> (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	9 g in 100 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>ALUMINUM OXIDE</b> (UNII: LMI26O6933)	
<b>BUTYLENE GLYCOL</b> (UNII: 3XUS85K0RA)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>CYCLOMETHICONE 5</b> (UNII: 0THT5PCI0R)	
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)	
<b>ERGOTHIONEINE</b> (UNII: BDZ3DQM98W)	
<b>HEXAMETHYLENE DIISOCYANATE/TRIMETHYLOL HEXYLLACTONE CROSSPOLYMER</b> (UNII: WB5K9Y35Y9)	
<b>IODOPROPYNYL BUTYLCARBAMATE</b> (UNII: 603P14DHEB)	
<b>FERROSFERRIC OXIDE</b> (UNII: XM0M87F357)	
<b>LAURYL PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE</b> (UNII: 25G622K2RA)	
<b>EGG PHOSPHOLIPIDS</b> (UNII: 1Z74184RGV)	
<b>METHICONE (20 CST)</b> (UNII: 6777U11MKT)	
<b>OCTYLDODECYL NEOPENTANOATE</b> (UNII: X8725R883T)	
<b>PHENOXYETHANOL</b> (UNII: HIE492ZZ3T)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	
<b>TRIETHOXYCAPRYLYLSILANE</b> (UNII: LDC331P08E)	
<b>MICROCOCCUS LUTEUS</b> (UNII: LV6L29Z6AX)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72251-001-74	74 mL in 1 TUBE; Type 0: Not a Combination Product	04/17/2018	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	10/08/2012	

**Labeler** - Pharma Cosmetics (080622701)

Revised: 10/2023

Pharma Cosmetics