

METHOCARBAMOL- methocarbamol tablet
Asclemed USA, Inc.

Methocarbamol Tablets 500mg

Methocarbamol Tablets, USP 500 mg


Rx only

DESCRIPTION

Methocarbamol Tablets, USP, 500 mg, a carbamate derivative of guaifenesin, is a central nervous system (CNS) depressant with sedative and musculoskeletal relaxant properties.

The chemical name of methocarbamol is 3-(2-methoxyphenoxy)-1, 2-propanediol 1-carbamate and has the empirical formula C₁₁H₁₅NO₅. Its molecular weight is 241.24.

The structural formula is shown below.



Methocarbamol is a white powder, sparingly soluble in water and chloroform, soluble in alcohol (only with heating) and propylene glycol, and insoluble in benzene and *n*-hexane.

Each tablet, for oral administration, contains either 500 mg of methocarbamol, USP. The inactive ingredients present are colloidal silicon dioxide, magnesium stearate, povidone, pregelatinized corn starch, purified water, sodium starch glycolate, and stearic acid.

CLINICAL PHARMACOLOGY

The mechanism of action of methocarbamol in humans has not been established, but may be due to general central nervous system (CNS) depression. It has no direct action on the contractile mechanism of striated muscle, the motor end plate or the nerve fiber.

Pharmacokinetics

In healthy volunteers, the plasma clearance of methocarbamol ranges between 0.20 and 0.80 L/h/kg, the mean plasma elimination half-life ranges between 1 and 2 hours, and the plasma protein binding ranges between 46% and 50%.

Methocarbamol is metabolized via dealkylation and hydroxylation. Conjugation of methocarbamol also is likely. Essentially all methocarbamol metabolites are eliminated in the urine. Small amounts of unchanged methocarbamol also are excreted in the urine.

Special populations

Elderly

The mean (\pm SD) elimination half-life of methocarbamol in elderly healthy volunteers (mean [\pm SD] age, 69 [\pm 4] years) was slightly prolonged compared to a younger (mean [\pm SD] age, 53.3 [\pm 8.8] years), healthy population (1.5 [\pm 0.4] hours versus 1.1 [\pm 0.27] hours, respectively). The fraction of bound methocarbamol was slightly decreased in the elderly versus younger volunteers (41 to 43% versus 46 to 50%, respectively).

Renally impaired

The clearance of methocarbamol in 8 renally-impaired patients on maintenance hemodialysis was reduced about 40% compared to 17 normal subjects, although the mean (\pm SD) elimination half-life in these two groups was similar: 1.2 (\pm 0.6) versus 1.1 (\pm 0.3) hours, respectively.

Hepatically impaired

In 8 patients with cirrhosis secondary to alcohol abuse, the mean total clearance of methocarbamol was reduced approximately 70% compared to that obtained in 8 age- and weight-matched normal subjects. The mean (\pm SD) elimination half-life in the cirrhotic patients and the normal subjects was 3.38 (\pm 1.62) hours and 1.11 (\pm 0.27) hours, respectively. The percent of methocarbamol bound to plasma proteins was decreased to approximately 40 to 45% compared to 46 to 50% in the normal subjects.

INDICATIONS AND USAGE

Methocarbamol is indicated as an adjunct to rest, physical therapy, and other measures for the relief of discomfort associated with acute, painful musculoskeletal conditions. The mode of action of methocarbamol has not been clearly identified, but may be related to its sedative properties.

Methocarbamol does not directly relax tense skeletal muscles in man.

CONTRAINDICATIONS

Methocarbamol is contraindicated in patients hypersensitive to methocarbamol or to any of the tablet components.

WARNINGS

Since methocarbamol may possess a general CNS depressant effect, patients receiving methocarbamol tablets should be cautioned about combined effects with alcohol and other CNS depressants.

Safe use of methocarbamol has not been established with regard to possible adverse effects upon fetal development. There have been reports of fetal and congenital abnormalities following in utero exposure to methocarbamol. Therefore, methocarbamol tablets should not be used in women who are or may become pregnant and particularly during early pregnancy unless in the judgment of the physician the potential benefits outweigh the possible hazards (see **PRECAUTIONS, Pregnancy**).

Use in Activities Requiring Mental Alertness

Methocarbamol may impair mental and/or physical abilities required for performance of hazardous tasks, such as operating machinery or driving a motor vehicle. Patients should be cautioned about operating machinery, including automobiles, until they are reasonably certain that methocarbamol therapy does not adversely affect their ability to engage in such activities.

PRECAUTIONS

Information for Patients

Patients should be cautioned that methocarbamol may cause drowsiness or dizziness, which may impair their ability to operate motor vehicles or machinery.

Because methocarbamol may possess a general CNS-depressant effect, patients should be cautioned about combined effects with alcohol and other CNS depressants.

Drug Interactions

See **WARNINGS** and **PRECAUTIONS** for interaction with CNS drugs and alcohol.

Methocarbamol may inhibit the effect of pyridostigmine bromide. Therefore, methocarbamol should be used with caution in patients with myasthenia gravis receiving anticholinesterase agents.

Drug/Laboratory Test Interactions

Methocarbamol may cause a color interference in certain screening tests for 5-hydroxyindoleacetic acid (5-HIAA) using nitrosonaphthol reagent and in screening tests for urinary vanillylmandelic acid (VMA) using the Gitlow method.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term studies to evaluate the carcinogenic potential of methocarbamol have not been performed. No studies have been conducted to assess the effect of methocarbamol on mutagenesis or its potential to impair fertility.

Pregnancy

Teratogenic Effects—Pregnancy Category C

Animal reproduction studies have not been conducted with methocarbamol. It is also not known whether methocarbamol can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Methocarbamol should be given to a pregnant woman only if clearly needed.

Safe use of methocarbamol has not been established with regard to possible adverse effects upon fetal development. There have been reports of fetal and congenital abnormalities following in utero exposure to methocarbamol. Therefore, methocarbamol should not be used in women who are or may become pregnant and particularly during early pregnancy unless in the judgment of the physician the potential benefits outweigh the possible hazards (see **WARNINGS**).

Nursing Mothers

Methocarbamol and/or its metabolites are excreted in the milk of dogs; however, it is not known whether methocarbamol or its metabolites are excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when methocarbamol is administered to a nursing woman.

Pediatric Use

Safety and effectiveness of methocarbamol in pediatric patients below the age of 16 have not been established.

ADVERSE REACTIONS

Adverse reactions reported coincident with the administration of methocarbamol include:

*Body as a whole:*Anaphylactic reaction, angioneurotic edema, fever, headache

*Cardiovascular system:*Bradycardia, flushing, hypotension, syncope, thrombophlebitis

*Digestive system:*Dyspepsia, jaundice (including cholestatic jaundice), nausea and vomiting

*Hemic and lymphatic system:*Leukopenia

*Immune system:*Hypersensitivity reactions

*Nervous system:*Amnesia, confusion, diplopia, dizziness or lightheadedness, drowsiness, insomnia, mild muscular incoordination, nystagmus, sedation, seizures (including grand mal), vertigo

*Skin and special senses:*Blurred vision, conjunctivitis, nasal congestion, metallic taste, pruritus, rash, urticaria

OVERDOSAGE

Limited information is available on the acute toxicity of methocarbamol. Overdose of methocarbamol is frequently in conjunction with alcohol or other CNS depressants and includes the following symptoms: nausea, drowsiness, blurred vision, hypotension, seizures, and coma.

In post-marketing experience, deaths have been reported with an overdose of methocarbamol alone or in the presence of other CNS depressants, alcohol or psychotropic drugs.

Treatment

Management of overdose includes symptomatic and supportive treatment. Supportive measures include maintenance of an adequate airway, monitoring urinary output and vital signs, and administration of intravenous fluids if necessary. The usefulness of hemodialysis in managing overdose is unknown.

DOSAGE AND ADMINISTRATION

Methocarbamol, 500 mg — Adults: Initial dosage: 3 tablets q.i.d.
Maintenance dosage: 2 tablets q.i.d.

Six grams a day are recommended for the first 48 to 72 hours of treatment. (For severe conditions 8 grams a day may be administered). Thereafter, the dosage can usually be reduced to approximately 4 grams a day.

HOW SUPPLIED

Methocarbamol Tablets, USP 500 mg — white, round, convex face, debossed “611” over bisect and “O” below bisect on one side and plain on the reverse side. Available in:

Bottles of 30, NDC 76420-667-30 (repackaged from NDC 70868-910-10)

Bottles of 60, NDC 76420-667-60 (repackaged from NDC 70868-910-10)

Bottles of 90, NDC 76420-667-90 (repackaged from NDC 70868-910-10)

Bottles of 100, NDC 76420-667-01 (re-labeled from NDC 70868-910-10)

Store at 20°– 25°C (68°– 77°F) [See USP Controlled Room Temperature].

Dispense in tight container.

Relabeled and Repackaged by:

Enovachem PHARMACEUTICALS

Torrance, CA 90501

PACKAGE LABEL. PRINCIPAL DISPLAY PANEL

Packaged By:



Methocarbamol Tablets, USP 500mg

NDC: 76420-667-30

Qty: 30

Manufactured For: KEY Therapeutics

Source NDC: 70868-910-10

Description: white/round/convex face; debossed 611; O

Lot #: 00000000

Exp:

Batch #: 00000000

Drug Status: RX

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].



(01) 0 0376420 67730 7

(17)

(10) 00000000

(21)

Methocarbamol Tablets, USP 500mg

NDC: 76420-667-30

S/N:

Qty: 30

Methocarbamol Tablets, USP 500mg

NDC: 76420-667-30

S/N:

Qty: 30

Methocarbamol Tablets, USP 500mg

NDC: 76420-667-30

S/N:

Qty: 30

Packaged By:



Methocarbamol Tablets, USP 500mg

NDC: 76420-667-60

Qty: 60

Manufactured For: KEY Therapeutics



(01) 0 0376420 67760 4

Methocarbamol Tablets, USP 500mg

NDC: 76420-667-60

S/N:

Qty: 60

Methocarbamol Tablets, USP 500mg

NDC: 76420-667-60

S/N:

Qty: 60



Manufactured For: KEY Therapeutics
 Source NDC: 70868-910-10
 Description: white/round/convex face;debossed 611;O
 Lot #: 00000000 Exp:
 Batch #: 00000000
 Drug Status: RX

(17)
 (10) 00000000
 (21)

Qty: 60

Methocarbamol Tablets, USP 500mg
 NDC: 76420-667-60
 S/N:
 Qty: 60

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
 KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].

Packaged By:



Methocarbamol Tablets, USP 500mg

NDC: 76420-667-90

Qty: 90

Manufactured For: KEY Therapeutics
 Source NDC: 70868-910-10
 Description: white/round/convex face;debossed 611;O
 Lot #: 00000000 Exp:
 Batch #: 00000000
 Drug Status: RX



(01) 0 0376420 67790 1
 (17)
 (10) 00000000
 (21)

Methocarbamol Tablets, USP 500mg
 NDC: 76420-667-90
 S/N:
 Qty: 90

Methocarbamol Tablets, USP 500mg
 NDC: 76420-667-90
 S/N:
 Qty: 90

Methocarbamol Tablets, USP 500mg
 NDC: 76420-667-90
 S/N:
 Qty: 90

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
 KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].

Relabeled By:



Methocarbamol Tablets, USP 500mg

NDC: 76420-667-01

Qty: 100

Manufactured For: KEY Therapeutics
 Source NDC: 70868-910-10
 Description: white/round/convex face;debossed 611;O
 Lot #: 00000000 Exp:
 Batch #: 00000000
 Drug Status: RX



(01) 0 0376420 67701 7
 (17)
 (10) 00000000
 (21)

Methocarbamol Tablets, USP 500mg
 NDC: 76420-667-01
 S/N:
 Qty: 100

Methocarbamol Tablets, USP 500mg
 NDC: 76420-667-01
 S/N:
 Qty: 100

Methocarbamol Tablets, USP 500mg
 NDC: 76420-667-01
 S/N:
 Qty: 100

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT.
 KEEP OUT OF REACH OF CHILDREN. STORE AT 20-25C (68-77F) [SEE USP CONTROLLED ROOM TEMP].

METHOCARBAMOL

methocarbamol tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:76420-667(NDC:70868-910)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
METHOCARBAMOL (UNII: 125OD7737X) (METHOCARBAMOL - UNII:125OD7737X)	METHOCARBAMOL	500 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE K90 (UNII: RDH86HJV5Z)	
STARCH, CORN (UNII: O8232NY3SJ)	
WATER (UNII: 059QF0KO0R)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	19mm
Flavor		Imprint Code	611;O
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76420-667-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	02/08/2024	
2	NDC:76420-667-60	60 in 1 BOTTLE; Type 0: Not a Combination Product	02/08/2024	
3	NDC:76420-667-90	90 in 1 BOTTLE; Type 0: Not a Combination Product	02/08/2024	
4	NDC:76420-667-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/08/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040489	01/01/2021	

Labeler - Asclemed USA, Inc. (059888437)

Establishment

Name	Address	ID/FEI	Business Operations
ASCLEMED USA INC. DBA ENOVACHEM		059888437	relabel(76420-667) , repack(76420-667)