

**SUNSCREEN LOTION FOR FACE- sunscreen lotion for face cellular sheet  
Shenzhen XiaoMai Manufacturing Co., Ltd.**

-----

**ACTIVE INGREDIENT**

Homosalate: 10%  
Titanium Dioxide: 5%  
Octocrylene: 2%  
Zinc Oxide: 9%  
Niacinamide: 4%  
Ceramide 3: 0.5%  
Ceramide 6 II: 0.5%  
Ceramide 1: 0.5%  
Sodium Hyaluronate: 1.5%  
Collagen: 1.5%  
Centella Asiatica Extract: 2.5%

**PURPOSE**

helps prevent sunburn.

**Uses**

decreases the risk of skin cancer and early skin aging caused by the sun.

**WARNINGS**

For external use only.

**Do not use**

on damaged or broken skin.

**When using this product**

keep out of eyes.  
Rinse with water to remove.

**Stop use and ask a doctor**

if rash occurs.

## **Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

## **Directions for use**

apply liberally 15 minutes before sun exposure.  
reapply at least every 2 hours.  
use a water resistant sunscreen if swimming or sweating.

## **INACTIVE INGREDIENT**

Cetearyl Alcohol: 4.5%  
Ethylcelluloses: 1%  
Carbomer: 0.2%  
Disodium EDTA: 0.6%  
Glycerin: 5.5%  
Dimethicone: 3.5%  
Behentrimonium Methosulfate: 1.5%  
Aluminum Starch Octenylsuccinate: 2%  
Propylparaben: 0.6%  
Sodium Lauroyl Aspartate: 0.3%  
Phytosphingosine: 0.2%  
Lactic Acid: 0.2%  
Xanthan Gum: 0.2%  
Water: 42.7%

## **Other Information**

Protect the product in this container from excessive heat and direct sun.

**Drug Facts**

| Active ingredients  | Purpose   |
|---------------------|-----------|
| Homosalate 10%      | Sunscreen |
| Titanium Dioxide 5% | Sunscreen |
| Octocrylene 2%      | Sunscreen |
| Zinc oxide 9%       | Sunscreen |

**Uses**

- helps prevent sunburn.
- if used as directed with other sun protection measures, decreases the risk of skin cancer and early skin aging caused by the sun.

**Warnings**

For external use only.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

For sunscreen use:

- apply liberally 15 minutes before sun exposure.
- reapply at least every 2 hours.
- use a water resistant sunscreen if swimming or sweating.
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m.-2 p.m.
  - wear long-sleeved shirts, pants, hats, and sunglasses.
  - children under 6 months of age: Ask a doctor.

## CLEAR SKIN

FOR ACNE PRONE + OILY SKIN

SUNSCREEN LOTION FOR FACE

# 50

BROAD SPECTRUM SPF 50

WON'T CAUSE BREAKOUTS  
HYPOALLERGENIC

1.7 FL OZ / 50 ml e

**Other information**

- protect the product in this container from excessive heat and direct sun.

**Ingredients**

Homosalate, Titanium Dioxide, Octocrylene, Zinc Oxide, Aqua, Niacinamide, Cetearyl Alcohol, Glycerin, Dimethicone, Behentrimonium Methosulfate, Ceramide 3, Ceramide 6-II, Ceramide 1, Sodium Hyaluronate, Collagen, Aluminum Starch Octenylsuccinate, Centella Asiatica Leaf Extract, Ethylcellulose, Disodium EDTA, Propylparaben, Sodium Lauroyl Aspartate, Phytosphingosine Lactate, Carbomer, Xanthan Gum,

**Country of Origin** China

**Manufacturer**  
Shenzhen XiaoMai Manufacturing Co., Ltd.

**Address**  
61 Guantian Road, Niuhu, Shenzhen, China

**MFG** 2024/3/20  
**EXP** 2026/3/20

**Batch Number** 20240202

**EXPIRY DATE**  
Before 24 Months (Sealed/Unopened)  
Best Before 12 Months (After Opening)

**MCM Product LTD**  
Unit 36 A&B Arcade brownlow  
Hill, Liverpool, L3 5TA,  
TEL: 0044 202547864  
E-Mail: mjcm19002@gmail.com

**MCM Limited**  
Floor A&L, Uffley Building,  
Dublin 1, Ireland  
TEL: +353 084683065  
E-Mail: mjcm19002@gmail.com

## SUNSCREEN LOTION FOR FACE

sunscreen lotion for face cellular sheet

### Product Information

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:83872-108 |
| <b>Route of Administration</b> | TOPICAL        |                           |               |

### Active Ingredient/Active Moiety

| Ingredient Name   | Basis of Strength | Strength      |
|---|-------------------|---------------|
| <b>CERAMIDE 1</b> (UNII: 5THT33P7X7) (CERAMIDE 1 - UNII:5THT33P7X7)             | CERAMIDE 1        | 5 mg in 1 g   |
| <b>OCTOCRYLENE</b> (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)           | OCTOCRYLENE       | 20 mg in 1 g  |
| <b>NIACINAMIDE</b> (UNII: 25X5118RD4) (NIACINAMIDE - UNII:25X5118RD4)           | NIACINAMIDE       | 40 mg in 1 g  |
| <b>CERAMIDE 6 II</b> (UNII: F1X8L2B00J) (CERAMIDE 6 II - UNII:F1X8L2B00J)       | CERAMIDE 6 II     | 5 mg in 1 g   |
| <b>HOMOSALATE</b> (UNII: V06SV4M95S) (HOMOSALATE - UNII:V06SV4M95S)             | HOMOSALATE        | 100 mg in 1 g |
| <b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP) | TITANIUM DIOXIDE  | 50 mg in 1 g  |
| <b>ZINC OXIDE</b> (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)             | ZINC OXIDE        | 90 mg in 1 g  |
| <b>CERAMIDE 3</b> (UNII: 4370DF050B) (CERAMIDE 3 - UNII:4370DF050B)             | CERAMIDE 3        | 5 mg in 1 g   |

### Inactive Ingredients

| Ingredient Name                                       | Strength     |
|---|--------------|
| <b>SODIUM LAUROYL ASPARTATE</b> (UNII: 328G13N8ZR)    | 3 mg in 1 g  |
| <b>PHYTOSPHINGOSINE</b> (UNII: GIN46U9Q2Q)            | 2 mg in 1 g  |
| <b>XANTHAN GUM</b> (UNII: TTV12P4NEE)                 | 2 mg in 1 g  |
| <b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)                 | 35 mg in 1 g |
| <b>BEHENTRIMONIUM METHOSULFATE</b> (UNII: 5SHP745C61) | 15 mg in 1 g |

|  |               |
|--|---------------|
| <b>GLYCERIN</b> (UNII: PDC6A3C00X)                         | 55 mg in 1 g  |
| <b>ALUMINUM STARCH OCTENYLSUCCINATE</b> (UNII: I9PJ006294) | 20 mg in 1 g  |
| <b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)                    | 6 mg in 1 g   |
| <b>LACTIC ACID</b> (UNII: 33X04XA5AT)                      | 2 mg in 1 g   |
| <b>WATER</b> (UNII: 059QF0KO0R)                            | 427 mg in 1 g |

### Product Characteristics

|                 |       |                     |  |
|-----------------|-------|---------------------|--|
| <b>Color</b>    | white | <b>Score</b>        |  |
| <b>Shape</b>    |       | <b>Size</b>         |  |
| <b>Flavor</b>   |       | <b>Imprint Code</b> |  |
| <b>Contains</b> |       |                     |  |

### Packaging

| # | Item Code        | Package Description                                 | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:83872-108-01 | 50 g in 1 BOTTLE; Type 0: Not a Combination Product | 04/17/2024           |                    |

### Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M020                                     | 04/17/2024           |                    |

**Labeler** - Shenzhen XiaoMai Manufacturing Co., Ltd. (712999147)

### Establishment

| Name                                     | Address | ID/FEI    | Business Operations    |
|--|---------|-----------|------------------------|
| Shenzhen XiaoMai Manufacturing Co., Ltd. |         | 712999147 | manufacture(83872-108) |

Revised: 4/2024

Shenzhen XiaoMai Manufacturing Co., Ltd.