

PYRID 2X CANINE ANTHELMINTIC- pyrantel pamoate suspension

Neogen Corporation-Mercer

Pyrid 2x
(pyrantel pamoate oral suspension)
Canine Anthelmintic Suspension


DESCRIPTION & INDICATIONS AND USAGE SECTIONS

Pyrid 2x
(pyrantel pamoate oral suspension)
Canine Anthelmintic Suspension

NOT FOR USE IN HUMANS

DESCRIPTION: Pyrid 2x liquid wormer is a suspension of pyrantel pamoate in a palatable butterscotch flavored vehicle. Each mL contains 4.54 mg of pyrantel base as pyrantel pamoate.

Pyrantel pamoate is a compound belonging to a family classified chemically as tetrahydropyrimidines. It is a yellow, water-insoluble crystalline salt of the tetrahydropyrimidine base and pamoic acid containing 34.7% base activity. The chemical structure and name are given below:



INDICATIONS AND USAGE: Pyrid 2x suspension is a highly palatable formulation intended as a single treatment for the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies. The presence of these parasites should be confirmed by laboratory fecal exam. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

Pyrid 2x suspension may also be used to prevent reinfestation of *T.canis* in puppies and adult dogs and in lactating bitches after whelping.

PRECAUTIONS SECTION

PRECAUTIONS: THIS PRODUCT IS A SUSPENSION AND AS SUCH WILL SEPARATE. TO ENSURE UNIFORM RESUSPENSION AND TO ACHIEVE PROPER DOSAGE, **IT IS EXTREMELY IMPORTANT THAT THE PRODUCT BE SHAKEN THOROUGHLY BEFORE EVERY USE. KEEP OUT OF REACH OF CHILDREN**

DOSAGE AND ADMINISTRATION Section

DOSAGE AND ADMINISTRATION: Administer 1 teaspoon (5 mL) for each 10 lb of body weight. It is not necessary to withhold food prior to or after treatment. Dogs usually find this dewormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2-4 weeks after treatment.

For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T. canis* reinfestation.

SAFETY AND EFFICACY STUDIES SECTION

SAFETY AND EFFICACY STUDIES: Critical (worm count) studies in dogs demonstrated that pyrantel pamoate at the recommended dosage is highly efficacious against *T. leonina* (99%), *T. canis* (85%), *A. caninum* (97%), and *U. stenocephala* (94%).

One of the most outstanding and significant features of pyrantel pamoate is its wide margin of therapeutic safety in dogs. The acute oral LD(50) of pyrantel pamoate administered in gelatin capsules to female and male dogs is greater than 314 mg base per lb of body weight, which indicates a therapeutic index in excess of 138 times the recommended dosage. In subacute and chronic studies, no significant morphological abnormalities could be attributed to pyrantel pamoate when administered to dogs at daily dose rates up to 94 mg base per pound of body weight (40x) for periods of 19, 30, and 90 days. Clinical studies conducted in a

wide variety of geographic locations using more than 40 different breeds of dogs showed no drug-induced toxic effects. Included in these studies were nursing pups, weaned pups, adults, pregnant bitches, and males at stud.

Additional data has demonstrated the safe use of pyrantel pamoate in dogs having heartworm infestations and/or receiving medication for heartworms, dogs exposed to organophosphate flea collars or flea/tick-dip treatments, and dogs undergoing concurrent treatment or medication at the time of worming such as immunization and antibacterial treatment.

STORAGE AND HANDLING SECTION

RECOMMENDED STORAGE: STORE BELOW 86° F (30° C). Keep Pyrid 2x in a secure location out of reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.
HOW SUPPLIED: Pyrid 2x is supplied in 2 fl. oz. (60 mL), 16 fl. oz. (474 mL), and 32 fl. oz. (946 mL) bottles.

CONTACT INFORMATION SECTION

Approved by FDA under ANADA # 200-007

Contact Information: To report suspected adverse events or for medical assistance contact 1-800-498-5743. For technical assistance or to obtain a copy of the Safety Data Sheet(SDS), contact Neogen Corporation at 1-859-254-1221. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>

Manufactured for: Neogen Corporation 620 Lesher Place Lansing, MI 48912
USA provet.net

For customer questions call 1-859-254-1221
For medical emergencies call 1-800-498-5743



Pyrid™ 2x

(pyrantel pamoate oral suspension)

Canine Anthelmintic Suspension

IMPORTANT: SHAKE WELL BEFORE USING

Active Ingredient: 4.54 mg of Pyrantel Base as Pyrantel Pamoate per mL

NOT FOR USE IN HUMANS

Approved by FDA under ANADA #200-007

NET CONTENTS - 16 fl. oz. (474 mL)

PROVECTA®

IMPORTANT: SHAKE WELL BEFORE USING

DOSAGE AND ADMINISTRATION: Administer 1 teaspoon (5 mL) for each 10 lb of body weight. It is not necessary to withhold food prior to or after treatment. Dogs usually find this dewormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2-4 weeks after treatment. For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T. canis* reinfestation.

RECOMMENDED STORAGE: STORE BELOW 86°F (30°C).

KEEP OUT OF REACH OF CHILDREN. Keep Pyrid 2x in a secure location out of reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.

PRODUCT LOT & EXPIRATION DATE PRINTED ON BOTTOM OF CONTAINER ▼

NDC: 59051-6009-6



Pyrid™ 2x

(pyrantel pamoate oral suspension)

Canine Anthelmintic Suspension

Active Ingredient: 4.54 mg of Pyrantel Base as Pyrantel Pamoate per mL

NOT FOR USE IN HUMANS

Approved by FDA under ANADA # 200-007

NET CONTENTS — 16 fl. oz. (474 mL)

PROVECTA®

INDICATIONS AND USAGE: For the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies. The presence of these parasites should be confirmed by laboratory fecal exam. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. Pyrid 2x suspension may also be used to prevent reinfestation of *T. canis* in puppies and adult dogs and in lactating bitches after whelping.

Read accompanying professional information under this label.

Manufactured for:
Neogen Corporation
620 Leshner Place
Lansing, MI 48912 USA
provetapet.com

For customer questions call 1-859-254-1221
For medical emergencies call 1-800-498-5743

Item Number 6009008000

PH00163_0125

Take Time



Observe Label Directions



READ PACKAGE EXTENDED CONTENT LABEL FOR FULL PRESCRIBING INFORMATION ►



Pyrid™ 2x

(pyrantel pamoate oral suspension)

Canine Anthelmintic Suspension

IMPORTANT: SHAKE WELL BEFORE USING

Active Ingredient: 4.54 mg of Pyrantel Base as Pyrantel Pamoate per mL

NOT FOR USE IN HUMANS

Approved by FDA under ANADA #200-007

NET CONTENTS - 32 fl. oz. (946 mL)

PROVECTA®

NDC: 59051-6009-7



Pyrid™ 2x
(pyrantel pamoate oral suspension)

Canine Anthelmintic Suspension

IMPORTANT: SHAKE WELL BEFORE USING

Active Ingredient: 4.54 mg of Pyrantel Base as Pyrantel Pamoate per mL

NOT FOR USE IN HUMANS

Approved by FDA under ANADA # 200-007

NET CONTENTS — 32 fl. oz. (946 mL)

PROVECTA®

Pyrid 2x


(pyrantel pamoate oral suspension)

Canine Anthelmintic Suspension

NOT FOR USE IN HUMANS

DESCRIPTION: Pyrid 2x liquid wormer is a suspension of pyrantel pamoate in a palatable butterscotch flavored vehicle. Each mL contains 4.54 mg of pyrantel base as pyrantel pamoate.

Pyrantel pamoate is a compound belonging to a family classified chemically as tetrahydropyrimidines. It is a yellow, water-insoluble crystalline salt of the tetrahydropyrimidine base and pamoic acid containing 34.7% base activity. The chemical structure and name are given below:



INDICATIONS AND USAGE: Pyrid 2x suspension is a highly palatable formulation intended as a single treatment for the removal of large roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*) in dogs and puppies. The presence of these parasites should be confirmed by laboratory fecal exam. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

Pyrid 2x suspension may also be used to prevent reinfestation of *T.canis* in puppies and adult dogs and in lactating bitches after whelping.

PRECAUTIONS: THIS PRODUCT IS A SUSPENSION AND AS SUCH WILL SEPARATE. TO ENSURE UNIFORM RESUSPENSION AND TO ACHIEVE PROPER DOSAGE, IT IS EXTREMELY IMPORTANT THAT THE PRODUCT BE SHAKEN THOROUGHLY BEFORE EVERY USE.

KEEP OUT OF REACH OF CHILDREN

DOSAGE AND ADMINISTRATION: Administer 1 teaspoon (5 mL) for each 10 lb of body weight. It is not necessary to withhold food prior to or after treatment. Dogs usually find this dewormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. It is recommended that dogs maintained under conditions of constant exposure to worm infestation should have a follow-up fecal exam within 2-4 weeks after treatment.

For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2-3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated at monthly intervals to prevent *T. canis* reinfestation.

SAFETY AND EFFICACY STUDIES: Critical (worm count) studies in dogs demonstrated that pyrantel pamoate at the recommended dosage is highly efficacious against *T. leonina* (99%), *T. canis* (85%), *A. caninum* (97%), and *U. stenocephala* (94%).

One of the most outstanding and significant features of pyrantel pamoate is its wide margin of therapeutic safety in dogs. The acute oral LD₅₀ of pyrantel pamoate administered in gelatin capsules to female and male dogs is greater than 314 mg base per lb of body weight, which indicates a therapeutic index in excess of 138 times the recommended dosage. In subacute and chronic studies, no significant morphological abnormalities could be attributed to pyrantel pamoate when administered to dogs at daily dose rates up to 94 mg base per pound of body weight (40x) for periods of 19, 30, and 90 days. Clinical studies conducted in a wide variety of geographic locations using more than 40 different breeds of dogs showed no drug-induced toxic effects. Included in these studies were nursing pups, weaned pups, adults, pregnant bitches, and males at stud.

Additional data has demonstrated the safe use of pyrantel pamoate in dogs having heartworm infestations and/or receiving medication for heartworms, dogs exposed to organophosphate flea collars or flea/tick-dip treatments, and dogs undergoing concurrent treatment or medication at the time of worming such as immunization and antibacterial treatment.

RECOMMENDED STORAGE: STORE BELOW 86° F (30° C). Keep Pyrid 2x in a secure location out of reach of dogs, cats, and other animals to prevent accidental ingestion or overdose.

HOW SUPPLIED: Pyrid 2x is supplied in 2 fl. oz. (60 mL), 16 fl. oz. (474 mL), and 32 fl. oz. (946 mL) bottles.

Approved by FDA under ANADA # 200-007

Contact Information: To report suspected adverse events or for medical assistance contact 1-800-498-5743. For technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact Neogen Corporation at 1-859-254-1221. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>

Manufactured for:
Neogen Corporation
620 Lesher Place
Lansing, MI 48912 USA
provetapet.com

For customer questions call 1-859-254-1221
For medical emergencies call 1-800-498-5743
Item Number 6009009000
PH00164_0125



PRODUCT LOT & EXPIRATION DATE PRINTED ON BOTTOM OF CONTAINER ▼

PYRID 2X CANINE ANTHELMINTIC

pyrantel pamoate suspension

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:59051-6009
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PYRANTEL PAMOATE (UNII: 81BK194Z5M) (PYRANTEL - UNII:4QIH0N49E7)	PYRANTEL	4.54 mg in 1 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59051-6009-6	473.2 mL in 1 BOTTLE, PLASTIC		
2	NDC:59051-6009-7	946.4 mL in 1 BOTTLE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200007	08/18/2025	

Labeler - Neogen Corporation-Mercer (042125879)

Establishment

Name	Address	ID/FEI	Business Operations
ABS Corporation		048022727	manufacture

Establishment			
Name	Address	ID/FEI	Business Operations
COSMA Spa		428655732	api manufacture

Revised: 8/2025

Neogen Corporation-Mercer