

SENNOSIDES- sennosides capsule, liquid filled
Softgel Healthcare Pvt Ltd

Sennosides 8.6 mg

Active ingredient (in each softgel)

Sennosides USP 8.6 mg

Purpose

Laxative

Uses

- relieves occasional constipation (irregularity)
- generally causes a bowel movement in 6-12 hours

Warnings

Do not use

- laxative products for longer than 1 week unless directed by a doctor

Ask a doctor before use if you have

- stomach pain
- nausea
- vomiting
- notice a sudden change in bowel habits that continue over a period of 2 weeks

Stop use and ask a doctor if

- you have rectal bleeding or fail to have a bowel movement after use of a laxative. These may indicate a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms

Directions

- do not take more than directed
- take with full glass of water
- take preferably at bedtime or as directed by a doctor

Age	Starting Dosage	Maximum Dosage
adults and children 12 years of age and over	2 softgels once a day	4 softgels twice a day
children 6 to under 12 years	1 softgel once a day	2 softgels twice a day
children under 6 years of age	ask a doctor	ask a doctor

Other information

- store at room temperature between 20°C - 25°C (68°F - 77°F)


Inactive ingredients

ferrosoferric oxide, fumaric acid, gelatin, glycerin, lecithin soybean, methyl paraben, purified water, ferric oxide red, sorbitol 70% solution, soy bean oil, yellow bees wax.

NDC 35916-1936-2

Sennosides 8.6 mg Capsules

- For constipation relief
- Promotes bowel movements




500 SOFTGELS

Drug Facts	
Active Ingredient (in each softgel) Sennosides USP 8.6 mg	Purpose Laxative
Uses ■ relieves occasional constipation (irregularity) ■ generally causes a bowel movement in 6-12 hours	
Warnings Do not use ■ laxative products for longer than 1 week unless directed by a doctor	
Ask a doctor before use if you have ■ stomach pain ■ nausea ■ vomiting ■ notice a sudden change in bowel habits that continue over a period of 2 weeks	
Stop use and ask a doctor if ■ you have rectal bleeding or fail to have a bowel movement after use of a laxative. These may indicate a serious condition.	
If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms	
Directions ■ do not take more than directed ■ take with full glass of water ■ take preferably at bedtime or as directed by a doctor	
Age	Starting Dosage Maximum Dosage
adults and children 12 years of age and over	2 softgels once a day 4 softgels twice a day
children 6 to under 12 years	1 softgel once a day 2 softgels twice a day
children under 6 years of age	ask a doctor ask a doctor
Other information ■ store at room temperature between 20°C - 25°C (68°F - 77°F)	
Inactive ingredients ferrosoferric oxide, fumaric acid, gelatin, glycerin, lecithin soy bean, methyl paraben, purified water, ferric oxide red, sorbitol 70% solution, soy bean oil, yellow bees wax.	

xxxxxxx

Manufactured by:
Softgel Healthcare Pvt. Ltd.
Survey No. 201,
Vandalur-Kelambakkam Road,
Pudupakkam Village,
Kannuram Project,
Tamilnadu - 603 103, India.
Mfg. Lic. No.: XXXXXXXXXXXXX



Unvarnished Area
LH - 25x16 mm

SENNOSIDES

sennosides capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:35916-1936
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SENNOSIDES (UNII: 3FYP5M0IJX) (SENNOSIDES - UNII:3FYP5M0IJX)	SENNOSIDES	8.6 mg

Inactive Ingredients

Ingredient Name	Strength
SOYBEAN OIL (UNII: 241ATL177A)	
GELATIN (UNII: 2G86QN327L)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
SORBITOL SOLUTION 70% (UNII: 8KW3E207O2)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
YELLOW WAX (UNII: 2ZA36H0S2V)	
FUMARIC ACID (UNII: 88XHZ13131)	

Product Characteristics

Color	brown (Opaque)	Score	no score
Shape	OVAL	Size	11mm
Flavor		Imprint Code	901
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:35916-1936-2	500 in 1 BOTTLE; Type 0: Not a Combination Product	06/28/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	06/28/2024	

Labeler - Softgel Healthcare Pvt Ltd (675584180)

Revised: 6/2024

Softgel Healthcare Pvt Ltd