


CEFADROXIL- cefadroxil powder, for suspension
Hikma Pharmaceuticals USA Inc.

CEFADROXIL for Oral suspension, USP
250 mg/5 mL and 500 mg/5 mL
Rev. 03/2019
Rx only

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Cefadroxil for oral suspension and other antibacterial drugs, Cefadroxil for oral suspension should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.

DESCRIPTION

Cefadroxil monohydrate is a semisynthetic cephalosporin antibiotic intended for oral administration. It is a white to yellowish-white crystalline powder. It is soluble in water and it is acid-stable. It is chemically designated as 5-Thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylic acid, 7-[[amino(4-hydroxyphenyl)acetyl]amino]-3-methyl-8-oxo-, monohydrate, [6R-[6 α (,7 β ((R*))]]-. It has the formula C₁₆H₁₇N₃O₅•H₂O and the molecular weight of 381.40. It has the following structural formula:



Cefadroxil for oral suspension USP contains cefadroxil monohydrate USP. After reconstitution, each 5 mL contains Cefadroxil monohydrate USP equivalent to 250 mg or 500 mg of cefadroxil. In addition, cefadroxil for oral suspension USP contains the following inactive ingredients: FD&C Yellow No. 6, orange powder flavor, pineapple powder flavor, polysorbate 80, sodium benzoate, sugar (sucrose), and xanthan gum.

Cefadroxil for oral suspension, USP is off-white yellow powder, forming an orange colored suspension on constitution.

CLINICAL PHARMACOLOGY

Cefadroxil monohydrate is rapidly absorbed after oral administration. Following single doses of 500 mg and 1000 mg, average peak serum concentrations were approximately 16 and 28 mcg/mL, respectively. Measurable levels were present 12 hours after administration. Over 90% of the drug is excreted unchanged in the urine within 24 hours. Peak urine concentrations are approximately 1800 mcg/mL during the period following a single 500-mg oral dose. Increases in dosage generally produce a proportionate increase in cefadroxil monohydrate urinary concentration. The urine antibiotic concentration, following a 1-g dose, was maintained well above the MIC of susceptible urinary pathogens for 20 to 22 hours.

Microbiology

In vitro tests demonstrate that the cephalosporins are bactericidal because of their inhibition of cell-wall synthesis. Cefadroxil has been shown to be active against the following organisms both *in vitro* and in clinical infections (see **INDICATIONS AND USAGE**):

Beta-hemolytic streptococci

Staphylococci, including penicillinase-producing strains

Streptococcus (Diplococcus) pneumoniae

Escherichia coli

Proteus mirabilis

Klebsiella species

Moraxella (Branhamella) catarrhalis

Note: Most strains of *Enterococcus faecalis* (formerly *Streptococcus faecalis*) and *Enterococcus faecium* (formerly *Streptococcus faecium*) are resistant to cefadroxil monohydrate. It is not active against most strains of *Enterobacter species*, *Morganella morganii* (formerly *Proteus morganii*), and *P. vulgaris*. It has no activity against *Pseudomonas species* and *Acinetobacter calcoaceticus* (formerly *Mima* and *Herellea species*).

Susceptibility Tests

Diffusion Techniques:

The use of antibiotic disk susceptibility test methods which measure zone diameter give an accurate estimation of antibiotic susceptibility. One such standard procedure^{1,3} which has been recommended for use with disks to test susceptibility of organisms to cefadroxil uses the cephalosporin class (cephalothin) disk. Interpretation involves the correlation of the diameters obtained in the disk test with the minimum inhibitory concentration (MIC) for cefadroxil.

Reports from the laboratory giving results of the standard single-disk susceptibility test with a 30 mcg cephalothin disk should be interpreted according to the following criteria:

Interpretive Criteria for Enterobacteriaceae, and Staphylococcus spp.

Zone Diameter (mm)	Interpretation	MIC (mcg/mL)
≥18	Susceptible (S)	≤8
15 to 17	Intermediate (I)	-
≤14	Resistant (R)	≥32

A report of "Susceptible" indicates that the pathogen is likely to be inhibited by generally achievable blood levels. A report of "Intermediate" suggests that the organism would be susceptible if high dosage is used or if the infection is confined to tissue and fluids (e.g., urine) in which high antibiotic levels are attained. A report of "Resistant" indicates that achievable concentrations of the antibiotic are unlikely to be inhibitory and other therapy should be selected.

Standardized procedures require the use of laboratory control organisms. The 30 mcg cephalothin disk should give the following zone diameters:

Organism	Zone Diameter (mm)
<i>Staphylococcus aureus</i> ATCC 25923	29 to 37
<i>Escherichia coli</i> ATCC 25922	17 to 22

Dilution Techniques:

When using the CLSI agar dilution or broth dilution (including microdilution) method^{2,3} or equivalent, the MIC values should be interpreted according to the following criteria:

**Interpretive Criteria for
Enterobacteriaceae, and
Staphylococcus spp.**

MIC (mcg/mL)	Interpretation
≤8	Susceptible (S)
16	Intermediate (I)
≥32	Resistant (R)

As with standard diffusion methods, dilution procedures require the use of laboratory control organisms. Standard cephalothin powder should provide the following MIC values:

Microorganism		MIC (mcg/mL)
<i>Escherichia coli</i>	ATCC 25922	4 to 16
<i>Staphylococcus aureus</i>	ATCC 29213	0.12 to 0.5

INDICATIONS AND USAGE

Cefadroxil for oral suspension is indicated for the treatment of patients with infection caused by susceptible strains of the designated organisms in the following diseases:

Urinary tract infections caused by *E. coli*, *P. mirabilis*, and *Klebsiella* species.

Skin and skin structure infections caused by staphylococci and/or streptococci.

Pharyngitis and/or tonsillitis caused by *Streptococcus pyogenes* (Group A beta-hemolytic streptococci).

Note: Only penicillin by the intramuscular route of administration has been shown to be effective in the prophylaxis of rheumatic fever. Cefadroxil monohydrate is generally effective in the eradication of streptococci from the oropharynx. However, data establishing the efficacy of cefadroxil monohydrate for the prophylaxis of subsequent rheumatic fever are not available.

Note: Culture and susceptibility tests should be initiated prior to and during therapy. Renal function studies should be performed when indicated.

To reduce the development of drug-resistant bacteria and maintain the effectiveness of cefadroxil for oral suspension and other antibacterial drugs, cefadroxil for oral suspension should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

CONTRAINDICATIONS

Cefadroxil monohydrate is contraindicated in patients with known allergy to the cephalosporin group of antibiotics.

WARNINGS

BEFORE THERAPY WITH CEFADROXIL MONOHYDRATE IS INSTITUTED, CAREFUL INQUIRY SHOULD BE MADE TO DETERMINE WHETHER THE PATIENT HAS HAD PREVIOUS HYPERSENSITIVITY REACTIONS TO CEFADROXIL, CEPHALOSPORINS, PENICILLINS, OR OTHER DRUGS. IF THIS PRODUCT IS TO BE GIVEN TO PENICILLIN-SENSITIVE PATIENTS, CAUTION SHOULD BE EXERCISED BECAUSE CROSS-SENSITIVITY AMONG BETA-LACTAM ANTIBIOTICS HAS BEEN CLEARLY DOCUMENTED AND MAY OCCUR IN UP TO 10% OF PATIENTS WITH A HISTORY OF PENICILLIN ALLERGY.

IF AN ALLERGIC REACTION TO CEFADROXIL MONOHYDRATE OCCURS, DISCONTINUE THE DRUG. SERIOUS ACUTE HYPERSENSITIVITY REACTIONS MAY REQUIRE TREATMENT WITH EPINEPHRINE AND OTHER EMERGENCY MEASURES, INCLUDING OXYGEN, INTRAVENOUS FLUIDS, INTRAVENOUS ANTIHISTAMINES, CORTICOSTEROIDS, PRESSOR AMINES, AND AIRWAY MANAGEMENT, AS CLINICALLY INDICATED.

Clostridium difficile associated diarrhea (CDAD) has been reported with use of nearly all antibacterial agents, including cefadroxil monohydrate, and may range in severity from mild diarrhea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

C. difficile produces toxins A and B which contribute to the development of CDAD. Hypertoxin producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibiotic use. Careful medical history is necessary since CDAD has been reported to occur over two months after the administration of antibacterial agents.

If CDAD is suspected or confirmed, ongoing antibiotic use not directed against *C. difficile* may need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibiotic treatment of *C. difficile*, and surgical evaluation should be instituted as clinically indicated.

PRECAUTIONS

General

Cefadroxil monohydrate should be used with caution in the presence of markedly impaired renal function (creatinine clearance rate of less than 50 mL/min/1.73 m²). (See **DOSAGE AND ADMINISTRATION**). In patients with known or suspected renal impairment, careful clinical observation and appropriate laboratory studies should be made prior to and during therapy.

Prescribing cefadroxil for oral suspension in the absence of a proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

Prolonged use of cefadroxil monohydrate may result in the overgrowth of nonsusceptible organisms. Careful observation of the patient is essential. If superinfection occurs during therapy, appropriate measures should be taken.

Cefadroxil monohydrate should be prescribed with caution in individuals with history of gastrointestinal disease particularly colitis.

Information for Patients

Patients should be counseled that antibacterial drugs including cefadroxil for oral suspension should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When cefadroxil for oral suspension is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may (1) decrease the effectiveness of the immediate treatment and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by cefadroxil for oral suspension or other antibacterial drugs in the

future.

Diarrhea is a common problem caused by antibiotics which usually ends when the antibiotic is discontinued. Sometimes after starting treatment with antibiotics, patients can develop watery and bloody stools (with or without stomach cramps and fever) even as late as two or more months after having taken the last dose of the antibiotic. If this occurs, patients should contact their physician as soon as possible.

Drug / Laboratory Test Interactions

Positive direct Coombs' tests have been reported during treatment with the cephalosporin antibiotics. In hematologic studies or in transfusion cross-matching procedures when antiglobulin tests are performed on the minor side or in Coombs' testing of newborns whose mothers have received cephalosporin antibiotics before parturition, it should be recognized that a positive Coombs' test may be due to the drug.

Carcinogenesis, Mutagenesis, and Impairment of Fertility

No long-term studies have been performed to determine carcinogenic potential. No genetic toxicity tests have been performed.

Pregnancy

Teratogenic Effects:

Pregnancy Category B

Reproduction studies have been performed in mice and rats at doses up to 11 times the human dose and have revealed no evidence of impaired fertility or harm to the fetus due to cefadroxil monohydrate. There are, however, no adequate and well controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Labor and Delivery

Cefadroxil monohydrate has not been studied for use during labor and delivery. Treatment should only be given if clearly needed.

Nursing Mothers

Caution should be exercised when cefadroxil monohydrate is administered to a nursing mother.

Pediatric Use

(See **DOSAGE AND ADMINISTRATION**).

Geriatric Use

Of approximately 650 patients who received cefadroxil for the treatment of urinary tract infections in three clinical trials, 28% were 60 years and older, while 16% were 70 years and older. Of approximately 1000 patients who received cefadroxil for the treatment of skin and skin structure infection in 14 clinical trials, 12% were 60 years and older while 4% were 70 years and over. No overall differences in safety were observed between the elderly patients in these studies and younger patients. Clinical studies of cefadroxil for the treatment of pharyngitis or tonsillitis did not include sufficient numbers of patients 65 years and older to determine whether they respond differently from younger patients. Other reported clinical experience with cefadroxil has not identified differences in responses between elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

Cefadroxil is substantially excreted by the kidney, and dosage adjustment is indicated for patients with renal impairment (see **DOSAGE AND ADMINISTRATION: Renal Impairment**). Because elderly

patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

ADVERSE REACTIONS

Gastrointestinal

Onset of pseudomembranous colitis symptoms may occur during or after antibiotic treatment (see **WARNINGS**). Dyspepsia, nausea and vomiting have been reported rarely. Diarrhea has also occurred.

Hypersensitivity

Allergies (in the form of rash, urticaria, angioedema, and pruritus) have been observed. These reactions usually subsided upon discontinuation of the drug. Anaphylaxis has also been reported.

Other

Other reactions have included hepatic dysfunction including cholestasis and elevations in serum transaminase, genital pruritus, genital moniliasis, vaginitis, moderate transient neutropenia, fever. Agranulocytosis, thrombocytopenia, idiosyncratic hepatic failure, erythema multiforme, Stevens-Johnson syndrome, serum sickness, and arthralgia have been rarely reported.

In addition to the adverse reactions listed above which have been observed in patients treated with cefadroxil, the following adverse reactions and altered laboratory tests have been reported for cephalosporin-class antibiotics:

Toxic epidermal necrolysis, abdominal pain, superinfection, renal dysfunction, toxic nephropathy, aplastic anemia, hemolytic anemia, hemorrhage, prolonged prothrombin time, positive Coombs' test, increased BUN, increased creatinine, elevated alkaline phosphatase, elevated aspartate aminotransferase (AST), elevated alanine aminotransferase (ALT), elevated bilirubin, elevated LDH, eosinophilia, pancytopenia, neutropenia.

Several cephalosporins have been implicated in triggering seizures, particularly in patients with renal impairment, when the dosage was not reduced (see **DOSAGE AND ADMINISTRATION** and **OVERDOSAGE**). If seizures associated with drug therapy occur, the drug should be discontinued. Anticonvulsant therapy can be given if clinically indicated.

To report SUSPECTED ADVERSE REACTIONS, contact West-Ward Pharmaceuticals Corp. at 1-877-233-2001, or the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

OVERDOSAGE

A study of children under six years of age suggested that ingestion of less than 250 mg/kg of cephalosporins is not associated with significant outcomes. No action is required other than general support and observation. For amounts greater than 250 mg/kg, induce gastric emptying.

In five anuric patients, it was demonstrated that an average of 63% of a 1 g oral dose is extracted from the body during a 6 to 8 hour hemodialysis session.

DOSAGE AND ADMINISTRATION

Cefadroxil for oral suspension is acid-stable and may be administered orally without regard to meals. Administration with food may be helpful in diminishing potential gastrointestinal complaints occasionally associated with oral cephalosporin therapy.

Adults

Urinary Tract Infections

For uncomplicated lower urinary tract infections (i.e., cystitis) the usual dosage is 1 or 2 g per day in a single (q.d.) or divided doses (b.i.d.).

For all other urinary tract infections the usual dosage is 2 g per day in divided doses (b.i.d.).

Skin and Skin Structure Infections

For skin and skin structure infections the usual dosage is 1 g per day in single (q.d.) or divided doses (b.i.d.).

Pharyngitis and Tonsillitis

Treatment of group A beta-hemolytic streptococcal pharyngitis and tonsillitis— 1 g per day in single (q.d.) or divided doses (b.i.d.) for 10 days.

Children

For urinary tract infections, the recommended daily dosage for children is 30 mg/kg/day in divided doses every 12 hours. For pharyngitis, tonsillitis, and impetigo, the recommended daily dosage for children is 30 mg/kg/day in a single dose or in equally divided doses every 12 hours. For other skin and skin structure infections, the recommended daily dosage is 30 mg/kg/day in equally divided doses every 12 hours. In the treatment of beta-hemolytic streptococcal infections, a therapeutic dosage of cefadroxil for oral suspension should be administered for at least 10 days.

See chart for total daily dosage for children.

DAILY DOSAGE OF CEFADROXIL FOR ORAL SUSPENSION

Child's Weight			
lbs	kg	250 mg/5 mL	500 mg/5 mL
10	4.5	½ tsp	-
20	9.1	1 tsp	-
30	13.6	1½ tsp	-
40	18.2	2 tsp	1 tsp
50	22.7	2½ tsp	1¼ tsp
60	27.3	3 tsp	1½ tsp
70 & Above	31.8+	--	2 tsp

Renal Impairment

In patients with renal impairment, the dosage of cefadroxil monohydrate should be adjusted according to creatinine clearance rates to prevent drug accumulation. The following schedule is suggested. In adults, the initial dose is 1000 mg of cefadroxil monohydrate and the maintenance dose (based on the creatinine clearance rate [mL/min/1.73 m²]) is 500 mg at the time intervals listed below.

Creatinine Clearances	Dosage Interval
0 to 10 mL/min	36 hours
10 to 25 mL/min	24 hours
25 to 50 mL/min	12 hours

Patients with creatinine clearance rates over 50 mL/min may be treated as if they were patients having normal renal function.

Reconstitution Directions for Oral Suspension
--

Bottle Size	Reconstitution Directions
100 mL	Suspend in a total of 67 mL water. Method: Tap bottle lightly to loosen powder. Add 67 mL of water in two portions. Shake well after each addition.
75 mL	Suspend in a total of 51 mL water Method: Tap bottle lightly to loosen powder. Add 51 mL of water in two portions. Shake well after each addition.
50 mL	Suspend in a total of 34 mL water Method: Tap bottle lightly to loosen powder. Add 34 mL of water in two portions. Shake well after each addition.

After reconstitution, store in refrigerator. Shake well before using. Keep container tightly closed. Discard unused portion after 14 days.

HOW SUPPLIED

Cefadroxil for oral suspension, USP is an off-white to yellow powder with orange and pineapple flavor, and is supplied as follows:

250 mg/5 mL 50 mL Bottle
 100 mL Bottle
 500 mg/5 mL 75 mL Bottle
 100 mL Bottle

Prior to reconstitution: Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

After reconstitution: Store in refrigerator. Shake well before using. Keep container tightly closed. Discard unused portion after 14 days.

REFERENCES

1. Clinical and Laboratory Standards Institute, Approved Standard, Performance Standards for Antimicrobial Disk Susceptibility Tests; Approved Standard – Eleventh Edition, Vol. 32 (1): M02-A11, Wayne, PA, January, 2012.
2. Clinical and Laboratory Standards Institute, Approved Standard: Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria that Grow Aerobically; Approved Standard – Ninth Edition, Vol. 32 (2): M07-A9, Wayne, PA, January, 2012.
3. Clinical and Laboratory Standards Institute. Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Second Informational Supplement. CLSI Document M100-S22, Vol. 32, No. 3, CLSI, Wayne, PA, January, 2012.

Distributed By:


West-Ward Pharmaceuticals Corp.
 Eatontown, NJ 07724 USA.

Manufactured By:


Jazeera Pharmaceutical Industries (JPI)
 AL-Kharj Road
 P.O Box 106229
 Riyadh 11666
 Saudi Arabia.

An Affiliate of:
Hikma Pharmaceuticals
P.O Box 182400
Amman 11118-Jordan
Revised: March 2019

NDC 0143-9767-01
Cefadroxil for Oral
Suspension, USP
250 mg/5mL
Rx only
100 mL (When Mixed)

<p>Physician leaflet enclosed. Remove before dispensing. When mixed as directed, each 5 mL (approximately one teaspoonful) will contain: cefadroxil monohydrate equivalent to 250 mg cefadroxil.</p> <p>Usual Dosage: See package insert for complete dosage information.</p> <p>Directions for Reconstitution: Tap bottle lightly to loosen powder. Add 67 ml of water in two portions. Shake well after each addition.</p> <p>Prior to reconstitution: Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].</p>	<p>NDC 0143-9767-01</p> <hr/> <p>CEFADROXIL FOR ORAL SUSPENSION, USP</p> <hr/> <p>250 mg/5 mL</p> <p>Rx Only</p> <p>100 mL (When Mixed)</p> <hr/> <p>Distributed by: West-Ward Pharmaceutical Corp. Eatontown, NJ 07724 USA</p>	<p>SHAKE WELL BEFORE USING</p> <p>After Reconstitution: Store in refrigerator. Keep container tightly closed. Discard unused portion after 14 days.</p> <p>KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.</p> <p>Manufactured by: Jazeera Pharmaceutical Industries P.O. Box 106229, Riyadh 11666 Saudi Arabia An Affiliate of: HIKMA Pharmaceuticals P.O. Box 182400, Amman 11118 Jordan</p> <p>12/2012</p>	 <p>3 0143-9767-015</p> <p>Exp. Date: Batch no.:</p>
---	---	---	---

NDC 0143-9766-01
Cefadroxil for Oral
Suspension, USP
500 mg/5 mL
Rx only
100 mL (When Mixed)

<p>Physician leaflet enclosed. Remove before dispensing. When mixed as directed, each 5 mL (approximately one teaspoonful) will contain: cefadroxil monohydrate equivalent to 500 mg cefadroxil.</p> <p>Usual Dosage: See package insert for complete dosage information.</p> <p>Directions for Reconstitution: Tap bottle lightly to loosen powder. Add 67 ml of water in two portions. Shake well after each addition.</p> <p>Prior to reconstitution: Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].</p>	<p>NDC 0143-9766-01</p> <hr/> <p>CEFADROXIL FOR ORAL SUSPENSION, USP</p> <hr/> <p>500 mg/5 mL</p> <p>Rx Only</p> <p>100 mL (When Mixed)</p> <hr/> <p>Distributed by: West-Ward Pharmaceutical Corp. Eatontown, NJ 07724 USA</p>	<p>SHAKE WELL BEFORE USING</p> <p>After Reconstitution: Store in refrigerator. Keep container tightly closed. Discard unused portion after 14 days.</p> <p>KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.</p> <p>Manufactured by: Jazeera Pharmaceutical Industries P.O. Box 106229, Riyadh 11666 Saudi Arabia An Affiliate of: HIKMA Pharmaceuticals P.O. Box 182400, Amman 11118 Jordan</p> <p>12/2012</p>	 <p>3 0143-9766-018</p> <p>Exp. Date: Batch no.:</p>
---	---	---	---

CEFADROXIL

cefadroxil powder, for suspension

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0143-9767
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CEFADROXIL (UNII: 280111G160) (CEFADROXIL ANHYDROUS - UNII:Q525PA8JJB)	CEFADROXIL ANHYDROUS	250 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	WHITE (off-white to yellow)	Score	
Shape		Size	
Flavor	ORANGE, PINEAPPLE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0143-9767-01	100 mL in 1 BOTTLE; Type 0: Not a Combination Product	11/28/2012	
2	NDC:0143-9767-50	50 mL in 1 BOTTLE; Type 0: Not a Combination Product	11/28/2012	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091036	11/28/2012	

CEFADROXIL

cefadroxil powder, for suspension

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0143-9766	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
	Ingredient Name	Basis of Strength	Strength	
	CEFADROXIL (UNII: 280111G160) (CEFADROXIL ANHYDROUS - UNII:Q525PA8JJB)	CEFADROXIL ANHYDROUS	500 mg in 5 mL	
Inactive Ingredients				
	Ingredient Name		Strength	
	FD&C YELLOW NO. 6 (UNII: H77VEI93A8)			
	POLYSORBATE 80 (UNII: 6OZP39ZG8H)			
	SODIUM BENZOATE (UNII: OJ245FE5EU)			
	SUCROSE (UNII: C151H8M554)			
	XANTHAN GUM (UNII: TTV12P4NEE)			
Product Characteristics				
Color	WHITE (off-white to yellow)	Score		
Shape		Size		
Flavor	ORANGE, PINEAPPLE	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0143-9766-01	100 mL in 1 BOTTLE; Type 0: Not a Combination Product	11/28/2012	
2	NDC:0143-9766-75	75 mL in 1 BOTTLE; Type 0: Not a Combination Product	11/28/2012	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA091036	11/28/2012		

Labeler - Hikma Pharmaceuticals USA Inc. (001230762)

Revised: 3/2019

Hikma Pharmaceuticals USA Inc.