#### CHLORAPREP SWABSTICK- chlorhexidine gluconate and isopropyl alcohol solution CareFusion 213 LLC

-----

ChloraPrep<sup>™</sup> Swabsticks

### Active ingredients

Chlorhexidine gluconate (CHG), 2% w/v and Isopropyl alcohol (IPA), 70% v/v

### Purpose

Antiseptic

### Use

for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

### Warnings

### For external use only. Flammable, keep away from fire or flame.

do not use with electrocautery procedures

### Allergy alert:

This product may cause a severe allergic reaction.

Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

### Do not use

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

### When using this product,

keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a

doctor.

## Stop use and ask a doctor if

irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

## Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

# Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximal treatment area using
- one single swabstick applicator is approximately 2.5 in. x 2.5 in. (42 cm <sup>2</sup>)
- three swabsticks sequentially is approximately 5 in. x 5 in. (160 cm<sup>2</sup>)
- tear pouch at side notch to reveal applicator handles. Do not touch foam applicator tip. Place foam flat side down on the treatment area.
- completely wet the treatment area with antiseptic
- dry surgical sites (e.g., abdomen or arm): use repeated back-and-forth strokes for 30 seconds. (When using the triple swabstick applicators, use each swabstick sequentially within the 30 seconds). Allow the area to air dry for approximately 30 seconds. Do not blot or wipe away.
- moist surgical sites (e.g., inguinal fold): use repeated back-and-forth strokes for 2 minutes. (When using the triple swabstick applicators, use each swabstick sequentially within the 2 minutes). Allow the area to air dry for approximately 1 minute. Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

# Other information

- store between 15-30 °C (59-86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)


# Inactive ingredient

USP purified water


# Questions?

- www.chloraprep.com
- call 1-800-523-0502 (M-F 8 a.m.-5 p.m. CST)

### Package/Label Principal Display Panel



PRINCIPAL DISPLAY PANEL-CARTON BD ChloraPrep<sup>™</sup> Single Swabstick Chlorhexidine gluconate (CHG), 2% w/v and Isopropyl alcohol (IPA), 70% v/v Swab Patient Preoperative Skin Preparation Sterile Solution Applicator is sterile if package is intact 1.75 mL Applicators NDC 054365-401-28 Ref 930100



### PRINCIPAL DISPLAY PANEL-CARTON

BD ChloraPrep<sup>™</sup> Triple Swabstick

Chlorhexidine gluconate (CHG), 2% w/v

and Isopropyl alcohol (IPA), 70% v/v Swab

Patient Preoperative Skin Preparation

Sterile Solution

Applicator is sterile if package is intact

5.25 mL Applicators

NDC 054365-401-29

Ref 930103

CHLORAPREP SWABSTICK chlorhexidine gluconate and isopropyl alcohol solution							
Product Information							
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:54365-401				
Route of Administration	TOPICAL						

		Ingredient Name	Bac	is of Strength	Strength	
				0		
	ILORHEXIDI	NE GLUCONATE (UNII: MOR84MUD8E) (CHLORHEXIDINE - 2L)	RHEXIDINE ONATE	20 mg in 1 mL		
	<b>DPROPYL AI</b> III:ND2M4163	<b>COHOL</b> (UNII: ND2M416302) (ISOPROPYL ALCOHOL - 02)	ISOPF	ROPYL ALCOHOL	0.7 mL in 1 mL	
In	active In	gredients				
		Ingredient Name		Strength		
w	ATER (UNII: (	)59QF0KO0R)				
Pa	ackaging					
#	Item Package Description			Marketing Start Date	Marketing End Date	
1	NDC:54365- 401-28	48 in 1 CARTON		05/13/2024		
1		1 in 1 POUCH				
1		1.75 mL in 1 APPLICATOR; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)				
2	NDC:54365- 401-29	40 in 1 CARTON		05/13/2024		
2		3 in 1 POUCH				
~		5.25 mL in 1 APPLICATOR; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)				
2						
2	larketir					
2	l <b>arketir</b> Marketir Categor	ig Information Application Number or Monograph M	larketin Dat		keting End Date	

Labeler - CareFusion 213 LLC (826496312)

Registrant - Becton, Dickinson and Company (832696038)

Establishment						
Name	Address	ID/FEI	Business Operations			
CareFusion 213 LLC		826496312	manufacture (54365-401) , analysis (54365-401) , label (54365-401) , pack (54365-401) , sterilize (54365-401)			

Revised: 4/2024

CareFusion 213 LLC