

CHLORAPREP SWABSTICK- chlorhexidine gluconate and isopropyl alcohol solution
CareFusion 213 LLC

BD ChloroPrep™ Swabsticks

BD ChloroPrep™ Single Swabstick
1.75 mL Applicators
Active ingredients

Chlorhexidine gluconate (CHG), 2% w/v
and Isopropyl alcohol (IPA), 70% v/v

Purpose

Antiseptic

Use

for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame.

- do not use with electrocautery procedures

Allergy alert:

|This product may cause a severe allergic reaction.

|Symptoms may include:

- |wheezing/difficulty breathing
- shock
- |facial swelling
- |hives
- |rash

|If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product

keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if

irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximal treatment area for one applicator
- one single swabstick applicator is approximately 2.5 in. x 2.5 in. (42 cm²)
- tear pouch at side notch to reveal applicator handles. Do not touch foam applicator tip. Place foam flat side down on the treatment area.
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): use repeated back-and-forth strokes for 30 seconds. (When using the triple swabstick applicators, use each swabstick sequentially within the 30 seconds). Allow the area to air dry for approximately 30 seconds. Do not blot or wipe away.
- **moist surgical sites** (e.g., inguinal fold): use repeated back-and-forth strokes for 2 minutes. (When using the triple swabstick applicators, use each swabstick sequentially within the 2 minutes). Allow the area to air dry for approximately 1 minute. Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15-30 °C (59-86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredient

- USP purified water

Questions?

- www.chloraprep.com
- call **1-800-523-0502** (M-F 8 a.m.-5 p.m. CST)

BD ChloroPrep™ Triple Swabstick

5.25 mL Applicators

Active ingredients

Chlorhexidine gluconate 2% w/v

Isopropyl alcohol 70% v/v

Purposes

Antiseptic

Antiseptic

Use

for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame.

- do not use with electrocautery procedures

Allergy alert

This product may cause a severe allergic reaction.

Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product

keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a

doctor.

Stop use and ask a doctor if

irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximal treatment area using three swabsticks sequentially is approximately **5 in. x 5 in. (160 cm²)**
- tear pouch at side notch to reveal applicator handles. Do not touch foam applicator tip. Place foam flat side down on the treatment area.
- completely wet the treatment area with antiseptic
- **dry surgical sites** (e.g., abdomen or arm): using each swabstick sequentially, use repeated back-and-forth strokes for a total of 30 seconds. Allow the area to air dry for approximately 30 seconds. Do not blot or wipe away.
- **moist surgical sites** (e.g., inguinal fold): using each swabstick sequentially, use repeated back-and-forth strokes for a total of 2 minutes. Allow the area to air dry for approximately 1 minute. Do not blot or wipe away.
- discard the applicator after a single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.

Other information

- store between 15-30 °C (59-86 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

Inactive ingredient

USP purified water

Questions?

- www.chloraprep.com
- **call 1-800-523-0502** (M-F 8 a.m.-5 p.m. CST)

**BD Chloraprep™ Single Swabstick
Package/Label Principal Display Panel**

 **BD ChloroPrep™ Single Swabstick**

NDC 54365-401-28


Chlorhexidine gluconate (CHG), 2% w/v
and Isopropyl alcohol (IPA), 70% v/v Swab
Patient Preoperative Skin Preparation

Sterile Solution

Applicator is sterile if package is intact

1.75 mL Applicator

See Drug Facts for full labeling information



REF 930100



CareFusion 213, LLC,
El Paso, TX 79912, subsidiary
of Becton, Dickinson and Co.
1-800-523-0502
(M-F 8 a.m. - 5 p.m. CST)
6-930110 • C090623
Made in the USA of US and
imported parts • www.bd.com

↑ Tear Here ↓



WARNING. FLAMMABLE.
KEEP AWAY FROM FIRE OR FLAME.

Directions:

- Tear pouch at notch.
- Pull across package to expose the end of the swabstick.
- Do not touch foam when removing swabstick from pouch.
- Place foam flat side down on treatment area.



PRINCIPAL DISPLAY PANEL-CARTON

BD ChloroPrep™ Single Swabstick
Chlorhexidine gluconate (CHG), 2% w/v
and Isopropyl alcohol (IPA), 70% v/v Swab
Patient Preoperative Skin Preparation
Sterile Solution
Applicator is sterile if package is intact
1.75 mL Applicators
NDC 054365-401-28
Ref 930100



BD ChloraPrep™ Triple Swabsticks

BD ChloraPrep™ Triple Swabsticks


Chlorhexidine gluconate (CHG), 2% w/v
and Isopropyl alcohol (IPA), 70% v/v Swab
Patient Preoperative Skin Preparation

Sterile Solution
Applicator is sterile if package is intact

5.25 mL Applicator

See Drug Facts for full labeling information

NDC 54365-401-29



REF 930103



Drug Facts	
Active ingredients Chlorhexidine gluconate 2% w/v Isopropyl alcohol 70% v/v	Purposes Antiseptic Antisepsis
Use For the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infections.	
Warnings For external use only. Flammable, keep away from fire or flame. Do not use with electrocautery procedures.	
Allergy alert: This product may cause or worsen allergic reactions. Symptoms may include: • wheezing/coughing/shortness of breath • shock • facial swelling • hives • rash If an allergic reaction occurs, stop use and seek medical help right away.	
Do not use: • on patients allergic to chlorhexidine gluconate or any other ingredient in this product. • for lumbar puncture or in contact with the meninges. • on open skin wounds or on a general skin cleanse.	
When using this product: Keep out of eyes, ears, and mouth. May cause irritation or permanent injury if poured in or on eyes. If contact occurs, rinse with cold water right away and contact a doctor.	
Stop use and ask a doctor if: irritation, numbness, or allergic reactions occur. There may be signs of a serious condition.	
Drug Facts (continued) Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions • use with care in premature infants or infants under 2 months of age. • These products may cause irritation or chemical burns. • Apply 2 swabs only for one applicator as represented by the diagram. • Use pouch at the neck to reveal applicator handle. Do not touch foam applicator tip. Place foam that side down on the prepared area. • Completely wet the treatment area with one swab. • Dry targeted area (e.g., abdomen or wrist) using repeated back-and-forth strokes for 30 seconds. Allow the area to air dry for approximately 30 seconds. Do not blot or wipe away. • Repeat targeted area (e.g., inguinal fold) using repeated back-and-forth strokes for 30 seconds. Allow the area to air dry for approximately 1 minute. Do not blot or wipe away. • Alcohol-free applicator often is single use along with any portion of the solution not required to cover the prep area. It is not necessary to use the entire amount available.	
Other information • store between 15-30 °C (59-86 °F) avoid freezing and excessive heat above 40 °C (104 °F)	
Inactive ingredient USP purified water	
Questions? • www.bdcorp.com • call 1-800-523-4562 (M-F 8 a.m.-5 p.m. EST)	

Classification 213, LLC
El Paso, TX 79912, subsidiary of
Becton, Dickinson and Co.
Made in the USA of US and imported parts
www.bd.com

BD, the BD Logo, ChloraPrep, and the Sterile Solution design are the property of Becton, Dickinson and Company or its affiliates.
©2023 BD. All rights reserved.

CareFusion 213, LLC, El Paso, TX 79912,
subsidiary of Becton, Dickinson and Co.
1-800-523-0502 (M-F 8 a.m. - 5 p.m. CST)
6-930111 • C090623 • Made in the USA of US
and imported parts • www.bd.com

↑

Tear Here

↓



WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.

Directions:

- Tear pouch at notch.
- Pull across package to expose the ends of the swabsticks.
- Do not touch foam when removing swabsticks from pouch.
- Place foam flat side down on treatment area.



Chlorhexidine gluconate (CHG), 2% w/v and Isopropyl alcohol (IPA), 70% v/v Swab

Patient Preoperative Skin Preparation

Sterile Solution

Applicator is sterile if package is intact

5.25 mL Applicators

NDC 54365-401-29

REF 930103



CHLORAPREP SWABSTICK

chlorhexidine gluconate and isopropyl alcohol solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:54365-401
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CHLORHEXIDINE GLUCONATE (UNII: MOR84MUD8E) (CHLORHEXIDINE - UNII:R4K00DY52L)	CHLORHEXIDINE GLUCONATE	20 mg in 1 mL
ISOPROPYL ALCOHOL (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)	ISOPROPYL ALCOHOL	0.7 mL in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0K00R)	

Packaging

#	Item	Package Description	Marketing	Marketing
---	------	---------------------	-----------	-----------

#	Code	Package Description	Start Date	End Date
1	NDC:54365-401-28	48 in 1 CARTON	05/13/2024	
1		1 in 1 POUCH		
1		1.75 mL in 1 APPLICATOR; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)		
2	NDC:54365-401-29	40 in 1 CARTON	05/13/2024	
2		3 in 1 POUCH		
2		5.25 mL in 1 APPLICATOR; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA021555	05/13/2024	

Labeler - CareFusion 213 LLC (826496312)

Registrant - Becton, Dickinson and Company (832696038)

Establishment

Name	Address	ID/FEI	Business Operations
CareFusion 213 LLC		826496312	manufacture(54365-401) , analysis(54365-401) , label(54365-401) , pack(54365-401) , sterilize(54365-401)

Revised: 10/2025

CareFusion 213 LLC