# AURA BARRIER- glyceryl laurate and lactic acid solution AgroChem Inc

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

-----

Aura Barrier Teat Dip

AURA BARRIER DIP

**TEAT DIP** 

WARNING

CAUSES SERIOUS EYE IRRITATION.

FOR FARM AND INDUSTRIAL USE ONLY.

KEEP OUT OF REACH OF CHILDREN.

**KEEP OUT OF DIRECT SUNLIGHT.** 

AVOID FREEZING.

**PREVENTION:** Wash skin thoroughly after handling. Wear eye protection/ face protection.

FIRST AID / RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/ container to an approved waste disposal plant

EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494

### **INGREDIENTS:**

Active Ingredients:

**Glycerol Monolaurate - 1%** 

Lactic Acid - 3%

Emollients - 7.5%

Use Aura Barrier Dip effectively to reduce the spread of mastitis causing organisms. The specially selected emollients in Aura Barrier Dip soften teat skin and protect the surface of the teat by keeping it moistened and smooth. Aura Barrier Dip is suitable for use as a pre or post dip.

Directions:

General Use

Ready to use. Do not dilute. Do not mix product with water, any other teat dips, or other product.

Use a teat dip cup to apply. Clean dip cup just prior to use.

If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip.

If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.

If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

Teat Sprays can also be foamed or sprayed using recommended equipment.

Pre Dipping Use

Perform forestripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or premilking teat dip. Teats should then be dried with single-service towels

Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product.

Allow product to stand for 30 seconds.

Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

Post Dipping Use

Immediately after each milking, use product.

Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe.

If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking.

Dipping should be resumed at least twice daily one week before calving.

# **AURA BARRIER DIP**

### **TEAT DIP**

### WARNING

#### CAUSES SERIOUS EYE IRRITATION. FOR FARM AND INDUSTRIAL USE ONLY. KEEP OUT OF REACH OF CHILDREN. KEEP OUT OF DIRECT SUNLIGHT. AVOID FREEZING.

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.

FIRST AID / RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/ container to an approved waste disposal plant

EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494

EXP:

INGREDIENTS: Active Ingredients: Glycerol Monolaurate - 1% Lactic Acid - 3% Emollients - 7.5%

NET CONTENTS:

LOT#:



Use Aura Barrier Dip effectively to reduce the spread of mastitis causing organisms. The specially selected emollients in Aura Barrier Dip soften teat skin and protect the surface of the teat by keeping it moistened and smooth. Aura Barrier Dip is suitable for use as a pre or post dip.

Directions: General Use

Ready to use. Do not dilute. Do not mix product with water, any other teat dips, or other product.

Use a teat dip cup to apply. Clean dip cup just prior to use.

If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip.

If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.

If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

Teat Sprays can also be foamed or sprayed using recommended equipment.

#### Pre Dipping Use

Perform forestripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels

Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product. Allow product to stand for 30 seconds.

Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

Post Dipping Use

Immediately after each milking, use product. Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe.

If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

**Extended Use** 

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking. Dipping should be resumed at least twice daily one week before calving.





# **AURA BARRIER DIP**

### **TEAT DIP**

### WARNING

CAUSES SERIOUS EYE IRRITATION. FOR FARM AND INDUSTRIAL USE ONLY. KEEP OUT OF REACH OF CHILDREN. KEEP OUT OF DIRECT SUNLIGHT. AVOID FREEZING.

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.

FIRST AID / RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention. STORAGE: Store locked up.

DISPOSAL: Dispose of contents/ container to an approved waste disposal plant

EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494

EXP:

INGREDIENTS: Active Ingredients: Glycerol Monolaurate - 1% Lactic Acid - 3% Emollients - 7.5%

NET CONTENTS:

LOT#:



Use Aura Barrier Dip effectively to reduce the spread of mastitis causing organisms. The specially selected emollients in Aura Barrier Dip soften teat skin and protect the surface of the teat by keeping it moistened and smooth. Aura Barrier Dip is suitable for use as a pre or post dip.

Directions: General Use

Ready to use. Do not dilute. Do not mix product with water, any other teat dips, or other product.

Use a teat dip cup to apply. Clean dip cup just prior to use.

If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip.

If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.

If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

Teat Sprays can also be foamed or sprayed using recommended equipment.

Pre Dipping Use

Perform forestripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels

Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product. Allow product to stand for 30 seconds.

Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

Post Dipping Use

Immediately after each milking, use product. Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe. If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking. Dipping should be resumed at least twice daily one week before calving.



AURA BARRIER					
glyceryl laurate and lactic acid solu	ution				
Product Information					
Product Type	OTC ANIMAL DRUG	Item Code (Source) NDC:17307-3934		7307-3934	
Route of Administration	TOPICAL				
Active Ingredient/Active Moi	ety				
Ing	redient Name		Basis of Str	ength	Strength
GLYCERYL LAURATE (UNII: Y98611C087) (GLYCERYL LAURATE - UNII:Y98611C087)			GLYCERYL LAURATE 0.0025		0.0025 kg in 1 kg
LACTIC ACID (UNII: 33X04XA5AT) (LACTIC ACID - UNII:33X04XA5AT)			LACTIC ACID		0.03 kg in 1 kg
Inactive Ingredients					
Ingredient Name			Strength		
LANOLIN (UNII: 7EV65EAW6H)			0.0025 kg in 1 kg		
GLYCERIN (UNII: PDC6A3C0OX)			0.04 kg in 1 kg		
POLYSORBATE 20 (UNII: 7T1F30V5YH)			0.03 kg in 1 kg		

C10-16 PARETH-1 (UNII: 3Y76363WPB)	0.003 kg in 1 kg
C9-11 PARETH-3 (UNII: 8049740MU0)	0.003 kg in 1 kg
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	0.035 kg in 1 kg
XANTHAN GUM (UNII: TTV12P4NEE)	0.0033 kg in 1 kg
ALGINIC ACID (UNII: 8C3Z4148WZ)	0.0022 kg in 1 kg

### Packaging

	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17307-3934-1	19 kg in 1 CONTAINER		
2	NDC:17307-3934-2	57 kg in 1 DRUM		
3	NDC:17307-3934-3	208 kg in 1 DRUM		
4	NDC:17307-3934-4	1040 kg in 1 CONTAINER		

or Monograph Citation	Marketing Start Date	Marketing End Date
	10/01/2013	
	or Monograph Citation	51 5

## Labeler - AgroChem Inc (805427700)

## Establishment

Name	Address	ID/FEI	<b>Business Operations</b>
AgroChem Inc.		805427700	api manufacture, manufacture

Revised: 2/2018

AgroChem Inc