

BERKLEY AND JENSEN OMEPRAZOLE- omeprazole tablet, orally disintegrating, delayed release
BJWC

BJWC Omeprazole Drug Facts

Active ingredient (in each tablet)

Omeprazole 20 mg

Purpose

Acid reducer

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert:

- do not use if you are allergic to omeprazole.
- omeprazole may cause severe skin reactions. Symptoms may include:
 - skin reddening
 - blisters
 - rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss

- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- take 1 tablet before eating in the morning
- **do not crush or chew tablets**
- place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- do not take this medicine with alcohol

Repeated 14-Day Course (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.

- store at 20-25°C (68-77°F); keep product out of high heat and moisture
- take the tablet immediately after opening individual blister. Do not store the tablet outside the blister pack for future use.

Inactive ingredients

amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, crospovidone, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize maltodextrin, mannitol, microcrystalline cellulose, propylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

Questions or comments?

1-800-719-9260: weekdays 7:30 AM to 5:00 PM EST

Package/Label Principal Display Panel

Compare to Prilosec OTC®

3 PACK

THREE 14-day courses of treatment

berkley jensen®

MELTech®

Melts In Your Mouth

OMEPRAZOLE

DELAYED RELEASE ORALLY DISINTEGRATING TABLETS, 20 mg

ACID REDUCER

ACTUAL SIZE

MELTS IN YOUR MOUTH ● DISSOLVES WITHOUT WATER

DISSOLVE TABS

TREATS FREQUENT HEARTBURN!

24HR

42 TABLETS

THREE 14-DAY COURSES OF TREATMENT

MAY TAKE 1 TO 4 DAYS FOR FULL EFFECT

STRAWBERRY FLAVOR

Compare to
Prilosec OTC®*

3 PACK

THREE 14-day
courses of treatment



OMEPRAZOLE

DELAYED RELEASE ORALLY DISINTEGRATING TABLETS, 20 mg

ACID REDUCER

MELTS IN YOUR MOUTH • DISSOLVES WITHOUT WATER

DISSOLVE TABS



**42
TABLETS**

THREE 14-DAY COURSES OF TREATMENT
MAY TAKE 1 TO 4 DAYS FOR FULL EFFECT



TREATS FREQUENT
HEARTBURN!
24HR



STRAWBERRY FLAVOR

ACT 33 DS 03

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking



8 88670 09087 9

031424280

Drug Facts

Active ingredient (in each tablet) Omeprazole 20 mg

Use: To reduce frequent heartburn (acid reflux) 1 to 4 days a week for 14 days. To reduce frequent heartburn (acid reflux) 1 to 4 days a week for 14 days.

Warnings: Allergy alert: Do not use if you are allergic to omeprazole. Do not use if you are allergic to any of the ingredients. Do not use if you are allergic to any of the ingredients. Do not use if you are allergic to any of the ingredients.

Do not take if you have: Allergy alert: Do not use if you are allergic to omeprazole. Do not use if you are allergic to any of the ingredients. Do not use if you are allergic to any of the ingredients.

Ask a doctor before use if you have: Allergy alert: Do not use if you are allergic to omeprazole. Do not use if you are allergic to any of the ingredients. Do not use if you are allergic to any of the ingredients.

Other information: Allergy alert: Do not use if you are allergic to omeprazole. Do not use if you are allergic to any of the ingredients. Do not use if you are allergic to any of the ingredients.

Drug Facts (continued)

Directions: For adults 18 years of age and older: Take 1 tablet 1 to 4 days for heartburn, once a day for 14 days. For children 1 to 17 years of age: See your doctor for advice.

14-Day Course of Treatment: Take 1 tablet with breakfast in the morning. Do not eat for 1 hour after taking. Do not take more than 1 tablet a day.

Repeat 14-Day Course (if needed): If you are still having heartburn after 14 days, you may repeat the 14-day course. Do not take more than 14 days of treatment in a row without a doctor's advice.

Other information: Allergy alert: Do not use if you are allergic to omeprazole. Do not use if you are allergic to any of the ingredients. Do not use if you are allergic to any of the ingredients.

Inactive ingredients: Allergy alert: Do not use if you are allergic to omeprazole. Do not use if you are allergic to any of the ingredients. Do not use if you are allergic to any of the ingredients.

Safety Feature - Do not use if printed tablet blister is not open or torn.



Every Berkley Jensen product comes with our **100% Money-Back Guarantee**.

Made in Israel
Distributed by:
BI's Wholesale Club, Inc.
350 Campus Drive
Marlborough, MA 01752

*This product is not manufactured or distributed by Procter & Gamble, distributor of Prilosec OTC®.

BERKLEY AND JENSEN OMEPRAZOLE

omeprazole tablet, orally disintegrating, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68391-103
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OMEPRazole (UNII: KG60484QX9) (OMEPRazole - UNII:KG60484QX9)	OMEPRazole	20 mg

Inactive Ingredients

Ingredient Name	Strength
DIMETHYLAMINOETHYL METHACRYLATE - BUTYL METHACRYLATE - METHYL METHACRYLATE COPOLYMER (UNII: 905HNO1SIH)	
ASCORBIC ACID (UNII: PQ6CK8PD0R)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MANNITOL (UNII: 3OWL53L36A)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM STEARATE (UNII: QU7E2XA9TG)	
SODIUM STEARYL FUMARATE (UNII: 7CV7WJK4UI)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	RED (reddish)	Score	no score
Shape	ROUND	Size	9mm
Flavor	STRAWBERRY	Imprint Code	20
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68391-103-55	3 in 1 CARTON	03/09/2020	
1		14 in 1 CARTON		
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

NDA	NDA209400	03/09/2020	
-----	-----------	------------	--

Labeler - BJWC (159082692)

Revised: 3/2025

BJWC