# BERKLEY AND JENSEN OMEPRAZOLE- omeprazole tablet, orally disintegrating, delayed release

**BJWC** 

-----

# **BWJC Omeprazole Drug Facts**

# **Active ingredient (in each tablet)**

Omeprazole 20 mg

# Purpose

Acid reducer

## Use

- treats frequent heartburn (occurs <u>2 or more</u> days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

# **Warnings**

**Allergy alert:** do not use if you are allergic to omeprazole

# Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

# Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

# Ask a doctor or pharmacist before use if you are

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

# Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months

- you get diarrhea
- you develop a rash or joint pain

# If pregnant or breast-feeding,

ask a health professional before use.

# Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

# **Directions**

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

# **14-Day Course of Treatment**

- take 1 tablet before eating in the morning
- do not crush or chew tablets
- place the tablet on tongue; tablet disintegrates, with or without water. The tablets can also be swallowed whole with water.
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor
- do not take this medicine with alcohol

# Repeated 14-Day Course (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

## Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F); keep product out of high heat and moisture

# **Inactive ingredients**

amino methacrylate copolymer, ascorbic acid, cetyl alcohol, colloidal silicon dioxide, crospovidone, ferric oxide, flavor, hypromellose, hypromellose phthalate, maize maltodextrin, mannitol, microcrystalline cellulose, propylene glycol, silicon dioxide, sodium stearate, sodium stearyl fumarate, sorbitol, sucralose, sugar spheres, talc, titanium dioxide, triethyl citrate

# Questions or comments?

**1-800-719-9260:** weekdays 7:30 AM to 5:00 PM EST

# Package/Label Principal Display Panel

Compare to Prilosec OTC®

3 PACK

THREE 14-day courses of treatment

 $MELTech^{^{TM}}$ 

Melts In Your Mouth

**OMEPRAZOLE** 

DELAYED RELEASE ORALLY DISINTEGRATING TABLETS 20mg

**ACID REDUCER** 

TREATS FREQUENT HEARTBURN!

24 HR

**42 TABLETS** 

**ACTUAL SIZE** 

MELTS IN YOUR MOUTH

DISSOLVES WITHOUT WATER

THREE 14-DAY COURSES OF TREATMENT

STRAWBERRY FLAVOR

100% MONEY-BACK GUARANTEE

MAY TAKE 1 TO 4 DAYS FOR FULL EFFECT



Compare to Prilosec OTC®\*

**3 PACK** 

THREE 14-day courses of treatment





# **OMEPRAZOLE**

**DELAYED RELEASE ORALLY DISINTEGRATING TABLETS 20mg** 

# **ACID REDUCER**

TREATS FREQUENT HEARTBURN!

24 HR

**TABLETS** 



MELTS IN YOUR MOUTH DISSOLVES WITHOUT WATER





THREE 14-DAY COURSES OF TREATMENT MAY TAKE 1 TO 4 DAYS FOR FULL EFFECT



- Tips for Man aging Heartburn
   Do not lie flat or bend over after eating
   Do not wear tight-fitting dothing around the

- Do not wear tight-fitting dothing around the stomach
  Do not eatbefore bedtime
  Raise the head of your bed
  Avoid heartburn-causing foods such as rich, sploy, fatly or field book, chocolate, cafeine, alcohol and certain fruits and vegetables
  Eatslowly and avoid big meals

## DrugFacts

Active in gredie nt (in each tab let) On sprzota 20 mg.

mirash frequenthearthum/ ocure <u>2 ormar e</u> fays aweek) moth baded for immed as at later hearth umbhi s drug may bke 1004 days for fall effect

When ing a Along a bit doned used from an abord of the emporation before the efficiency and the effect of the effect and to be the effect of the effect of the effect of the effect about the effect of the effect of the effect of the effect mobile to the strippin with white or of the effect of the effect producing commands externationally all behandables. These may be sign of the effect of the effect of the effect These may be sign of the effect of the effect of the effect All stories of the end the effect of the effect of the effect All stories of the end the effect of the effect of the effect and the effect of the effect of the effect of the effect and the effect of the effect of the effect of the effect and the effect of the effect of the effect of the effect and the effect of the effect of the effect and the eff

# Abk ad outorb of one u selfy outh ave: whad heartburn over 3 months. This may be as ign of armore serbes

# Mination neutralines of condition and service of the condition of the cond

# emission and manufacture of the second process of the second part of t

## Drug Facts (continued)

Drug Face (continued)
(Procedions 5
(Procedi

ne tour h active ing and iest a anio metacryth polymus, south activally absoluted that it we desting any per tons, in the orde, and any other properties of the control of the control

Question sor comments? 1800-719-926 0: w



"This product is not manufactured or distributed by Procker & Gamble , distributor of Pribises OTC®.

# BERKLEY AND JENSEN OMEPRAZOLE

omeprazole tablet, orally disintegrating, delayed release

# **Product Information**

HUMAN OTC DRUG Product Type Item Code (Source) NDC:68391-103

Route of Administration ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
OMEPRAZOLE (UNII: KG60484QX9) (OMEPRAZOLE - UNII:KG60484QX9)	OMEPRAZOLE	20 mg	

Inactive Ingredients	
Ingredient Name	Strength
DIMETHYLAMINO ETHYL METHACRYLATE - BUTYL METHACRYLATE - METHYL METHACRYLATE COPOLYMER (UNII: 905HNO1SIH)	
ASCORBIC ACID (UNII: PQ6CK8PD0R)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
SILICON DIO XIDE (UNII: ETJ7Z6 XBU4)	
CROSPOVIDONE (15 MPA.S AT 5%) (UNII: 68401960 MK)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29 V3WO)	
MANNITOL (UNII: 3OWL53L36A)	
MICRO CRYSTALLINE CELLULO SE (UNII: OP1R32D61U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM STEARATE (UNII: QU7E2XA9TG)	
SODIUM STEARYL FUMARATE (UNII: 7CV7WJK4UI)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics			
Color	RED (reddish)	Score	no score
Shape	ROUND	Size	9 mm
Flavor	STRAWBERRY	Imprint Code	20
Contains			

F	Packaging			
#	Item Code	Package Description	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:68391-103-55	3 in 1 CARTON	03/09/2020	
1		14 in 1 CARTON		
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA209400	03/09/2020	

Revised: 3/2020 BJWC