

MEIJER EFFERVESCENT PAIN RELIEF- aspirin tablet, effervescent
Meijer Distribution Inc

Meijer Effervescent Pain Relief

Active ingredient

(in each effervescent tablet)

Buffered aspirin 325 mg **(NSAID)***

*Nonsteroidal anti-inflammatory drug

Purpose

(in each effervescent tablet)

Buffered aspirin 325 mg **(NSAID)***. Pain reliever/fever reducer

*Nonsteroidal anti-inflammatory drug

Uses

■ for the temporary relief of minor aches and pains due to:

■ headache ■ muscle pain ■ backache

■ toothache ■ menstrual pain ■ colds

■ minor arthritis pain

■ temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma ■ you have a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

Ask a doctor or pharmacist before use if you are taking a prescription drug for
■ diabetes ■ gout ■ arthritis

When using this product

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding

- feel faint ■ vomit blood ■ have bloody or black stools
- have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- pain lasts for more than 10 days or gets worse
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children.

In case of overdose

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

■ do not exceed recommended dosage

■ to avoid serious injury, completely dissolve tablets in 4 oz. of water before taking
adults and children 12 years and over: 2 tablets every 4 hours, or as directed by a doctor - do not exceed 8 tablets in 24 hours

60 years and over: 2 tablets every 4 hours, or as directed by a doctor - do not exceed 4 tablets in 24 hours

children under 12 years: consult a doctor

Take this product any time - morning, noon or night - when you need relief from headaches, backaches, body aches or muscle pain.

Place two (2) tablets in 4 oz. of water. Dissolve tablets completely.

You do not have to drink any of the residue that may be on the bottom of the glass. The medicines are in the water.

Other information

■ **each tablet contains:** sodium 524 mg

■ store at room temperature (59°-86°F). Avoid excessive heat.

Inactive ingredients

anhydrous citric acid, sodium bicarbonate

Questions or comments

Questions or comments? 1-888-228-6937



MEIJER EFFERVESCENT PAIN RELIEF

aspirin tablet, effervescent

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:79481-0073
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	

SODIUM BICARBONATE (UNII: 8MDF5V39QO)

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	25mm
Flavor		Imprint Code	L421
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79481-0073-6	18 in 1 CARTON	06/06/2024	
1		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	06/06/2024	

Labeler - Meijer Distribution Inc (006959555)

Revised: 6/2024

Meijer Distribution Inc