

CAREONE DIARRHEA CONTROL- loperamide hcl tablet, film coated
American Sales Company

American Sales Company Diarrhea Control Drug Facts

Active ingredient (in each caplet)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- store at 20°-25°C (68°-77°F)
- see end panel for lot number and expiration date

Inactive ingredients

anhydrous lactose, carnauba wax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

Questions or comments?

1-800-719-9260

Principal Display Panel

CAREone®

Compare to the active ingredient in Imodium[®] A-D Caplets

DIARRHEA CONTROL

Loperamide Hydrochloride Tablets, 2 mg

Anti Diarrheal

Anti Diarrheal

Controls the symptoms of diarrhea

Gluten Free

See New Warnings

Actual Size

OUR PHARMACISTS RECOMMEND

12 CAPLETS*

*Capsule-Shaped Tablets

22453 OF C3

Drug Facts

Active ingredient
(in each caplet)
Loperamide HCl 2 mg Anti-diarrheal

Use controls symptoms of diarrhea, including travelers' diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl
Heart alert: Taking more than directed can cause serious heart problems or death
Do not use if you have bloody or black stool
Ask a doctor before use if you have
■ fever
■ mucus in the stool
■ a history of liver disease
■ a history of abnormal heart rhythm
Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.
When using this product: redness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.
Stop use and ask a doctor if
■ symptoms get worse
■ diarrhea lasts for more than 2 days
■ you get abdominal swelling or bloating.
These may be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea
■ find right dose on chart, if possible, use weight to dose; otherwise, use age.
adults and children 12 years and over
2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)
1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)
1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)
ask a doctor
children under 2 years (up to 33 lbs)
do not use

Other information

■ store at 20°C (68°F)
■ see end panel for lot number and expiration date

Drug Facts (continued)

Inactive ingredients anhydrous lactose, canabawax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

Drug Facts (continued)

Questions or comments?
1-800-719-9280

DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN
**This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Imodium® A-D Caplets.
DISTRIBUTED BY: FOODHOLD U.S.A., LLC
LANDOVER, MD 20785 · 1-877-846-9949
©2019 Ahold Delhaize Licensing Sàrl
Quality guaranteed or your money back.

CAREone®


NDC 41520-813-53

Compare to the active ingredient in Imodium® A-D Caplets**

DIARRHEA CONTROL

Loperamide Hydrochloride Tablets, 2 mg
Anti Diarrheal





CAREONE DIARRHEA CONTROL

loperamide hcl tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41520-813
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	

Product Characteristics

Color	GREEN	Score	2 pieces
Shape	OVAL	Size	10mm
Flavor		Imprint Code	L2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41520-813-53	12 in 1 CARTON	09/14/2016	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:41520-813-62	24 in 1 CARTON	09/14/2016	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:41520-813-67	1 in 1 CARTON	09/14/2016	01/31/2022
3		48 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075232	09/14/2016	

Labeler - American Sales Company (809183973)

Revised: 12/2025

American Sales Company