

**DAYTIME SEVERE COLD AND FLU RELIEF- acetaminophen dextromethorphan  
hbr guaifenesin, phenylephrine hcl liquid  
Family Dollar (FAMILY WELLNESS)**

-----  
**Drug Facts**

**Active ingredients (in each 15 mL)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

**Purpose**

**Pain reliever/fever reducer**

Cough suppressant

Expectorant

Nasal decongestant

**Uses**

- temporarily relieves common cold and flu symptoms:
  - nasal congestion
  - sinus congestion and pressure
  - minor aches and pains
  - headache
  - fever
  - sore throat
  - reduces swelling of nasal passages
  - cough due to minor throat and bronchial irritation
  - temporarily restores freer breathing through the nose
  - promotes nasal and or sinus drainage
  - helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make cough more productive

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- adult take more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount
- child takes more than 4 doses (15 mL each) in 24 hours

- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blistere
- rash.

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if the user has**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- a sodium-restricted diet
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

### **Ask a doctor or pharmacist before use if user is**

taking the blood thinning drug warfarin

### **When using this product**

**do not take more than directed.**

### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present

- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause serious liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults and for children even if you do not notice any signs or symptoms

**Directions**

- **take only as directed (see overdose warning)**
- do not take more than 4 doses in any 24 hours
- measure only with dosing cup provided. Do not use any other dosing device
- mL = milliliter
- keep dosing cup with product

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children 4 to under 6 years	ask a doctor
children under 4 years	do not use

- **when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing**

**Other information**

- each 15 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients**

citric acid, FD&C Yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate sodium chloride, sodium citrate, sorbitol sucralose, xanthan gum

**Questions or comments?**

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

**Principal Display Panel**

Compare to the active ingredients in Vicks® DayQuil® Severe Cold & Flu

## **Severe**

## **Daytime**

## **Cold & Flu Relief**

Acetaminophen 325 mg

pain reliever/Fever reducer

Dextromethorphan HBr 10 mg

Cough suppressant

Guaifenesin 200 mg

Expectorant

Phenylephrine HCl 5 mg

Nasal decongestant

Relieves

- headache, fever, sore throat, minor aches & pains
- nasal/sinus congestion & sinus pressure
- cough
- Chest congestion

For ages 6 years and over

Max strength • Non-drowsy • Alcohol free

FL OZ (mL)

\*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and DayQuil® are registered trademarks of The Procter & Gamble Company.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.**

DISTRIBUTED BY:

## **Product Label**

NDC 55319-145-08  
\*Compare to the active ingredients in  
Vicks® DayQuil® Severe Cold & Flu

**Severe  
Daytime  
Cold & Flu Relief**

**Acetaminophen 325 mg**  
Pain reliever/Fever reducer

**Dextromethorphan HBr 10 mg**  
Cough suppressant

**Guaifenesin 200 mg**  
Expectorant

**Phenylephrine HCl 5 mg**  
Nasal decongestant

**Relieves**  
✓ Headache, fever, sore throat, minor aches & pains  
✓ Nasal/sinus congestion & Sinus pressure  
✓ Cough  
✓ Chest congestion

For ages 6 years and over  
Max strength • Non-drowsy • Alcohol free

Questions or comments?  
Call 1-877-753-3935  
Monday-Friday 9AM-5PM EST

DISTRIBUTED BY: MIDWOOD BRANDS, LLC  
500 VOLVO PKWY, CHESAPEAKE, VA 23320 USA

**NOT 100% SATISFIED?**  
Return within 30 days to the store of purchase for a refund (with receipt) or exchange.

**8 fl oz (237 mL)**

0 32251 14113 3  
500 99976 PLD-A340B LB009764

**Drug Facts**

**Active ingredients (in each 15 mL)**

Acetaminophen 325 mg.....Pain reliever/fever reducer  
Dextromethorphan HBr 10 mg.....Cough suppressant  
Guaifenesin 200 mg.....Expectorant  
Phenylephrine HCl 5 mg.....Nasal decongestant

**Purposes**

**Uses**

- temporarily relieves common cold/flu symptoms
- nasal congestion
- sinus congestion and pressure
- minor aches and pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- cough due to minor throat and bronchial irritation

\*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and DayQuil® are registered trademarks of The Procter & Gamble Company.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.**

**PARENTS:**  
Learn about how medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

PLD-A340B LB009764

**PEEL CORNER FOR MORE DRUG FACTS**

**Drug Facts (continued)**

- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if:

- adult takes more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount
- child takes more than 4 doses (15 mL each) in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

if a skin reaction occurs, stop use and seek medical help right away.

**PEEL CORNER FOR MORE DRUG FACTS**

**Drug Facts (continued)**

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if the user has**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- a sodium-restricted diet
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Ask a doctor or pharmacist before use if user is taking the blood thinning drug warfarin.**

**Drug Facts (continued)**

**When using this product, do not take more than directed.**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Drug Facts (continued)**

**Directions**

- take only as directed (see Overdose warning)
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL = milliliter
- keep dosing cup with product

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children 4 to under 6 years	ask a doctor
children under 4 years	do not use

when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

**Other information**

- each 15 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

**Inactive ingredients** citric acid, FD&C yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

**FAMILY WELLNESS Severe Daytime Cold & Flu Relief**

**DAYTIME SEVERE COLD AND FLU RELIEF**

acetaminophen dextromethorphan hbr guaifenesin, phenylephrine hcl liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55319-145
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 15 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

Inactive Ingredients	
Ingredient Name	Strength
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-145-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/31/2024	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/31/2024	

**Labeler** - Family Dollar (FAMILY WELLNESS) (024472631)