

**DAYTIME SEVERE COLD AND FLU RELIEF- acetaminophen dextromethorphan
hbr guaifenesin, phenylephrine hcl liquid
Family Dollar (FAMILY WELLNESS)**

Drug Facts

Active ingredients (in each 15 mL)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

- temporarily relieves common cold and flu symptoms:
 - nasal congestion
 - sinus congestion and pressure
 - minor aches and pains
 - headache
 - fever
 - sore throat
 - reduces swelling of nasal passages
 - cough due to minor throat and bronchial irritation
 - temporarily restores freer breathing through the nose
 - promotes nasal and or sinus drainage
 - helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make cough more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- adult take more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount
- child takes more than 4 doses (15 mL each) in 24 hours

- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blistere
- rash.

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if the user has

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if user is

taking the blood thinning drug warfarin

When using this product

do not take more than directed.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs

of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause serious liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults and for children even if you do not notice any signs or symptoms

Directions

- take only as directed (see overdose warning)
- do not take more than 4 doses in any 24 hours
- measure only with dosing cup provided. Do not use any other dosing device
- mL = milliliter
- keep dosing cup with product

| | |
|---------------------------------------|---------------------|
| adults and children 12 years and over | 30 mL every 4 hours |
| children 6 to under 12 years | 15 mL every 4 hours |
| children 4 to under 6 years | ask a doctor |
| children under 4 years | do not use |

- **when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing**

Other information

- each 15 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

citric acid, FD&C Yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate sodium chloride, sodium citrate, sorbitol sucralose, xanthan gum

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

COMPARE TO THE ACTIVE INGREDIENTS IN VICKS® DAYQUIL™ SEVERE COLD & FLU*

DAYTIME

COLD & FLU RELIEF

SEVERE

ACETAMINOPHEN 325 mg

pain reliever/Fever reducer

Dextromethorphan HBr 10 mg

Cough suppressant

Guaifenesin 200 mg

Expectorant

Phenylephrine HCl 5 mg

Nasal decongestant

Relieves

- Headache, fever, sore throat, minor aches & pains
- Chest congestion, thins & loosen mucus
- Nasal congestion, sinus pressure
- cough

- Alcohol-Free

FL OZ (mL)

*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and DayQuil™ are registered trademarks of The Procter & Gamble Company.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY: MIDWOOD BRANDS, LLC

510 VOLVO PKWY, CHESAPEAKE, VA 23320

Product Label

FAMILY Wellness™ COMPARE TO THE ACTIVE INGREDIENTS IN VICKS® DAYQUIL™ SEVERE COLD & FLU*

DAYTIME COLD & FLU RELIEF

SEVERE

ACETAMINOPHEN 325 mg

Pain reliever/Fever reducer
Dextromethorphan HBr 10 mg
Cough suppressant
Guaifenesin 200 mg - Expectorant
Phenylephrine HCl 5 mg - Nasal decongestant

RELIEVES

- Headache, fever, sore throat, minor aches & pains
- Chest congestion, thins & loosens mucus
- Nasal congestion, sinus pressure
- Cough

Alcohol-Free

Questions or comments?
Call 1-877-753-3935
Monday-Friday 9AM-5PM EST

DISTRIBUTED BY: MIDWOOD BRANDS, LLC
510 VOLVO PKWY, CHESAPEAKE, VA 23320 USA

NOT 100% SATISFIED? Return within 30 days to the store of purchase for a refund (with receipt) or exchange.

8 FL OZ (237 mL) NDC 55319-145-08

SKU 99076 PLD-B340C LB010679

Drug Facts

| Active ingredients (in each 15 mL) | Purposes |
|------------------------------------|-----------------------------|
| Acetaminophen 325 mg | Pain reliever/fever reducer |
| Dextromethorphan HBr 10 mg | Cough suppressant |
| Guaifenesin 200 mg | Expectorant |
| Phenylephrine HCl 5 mg | Nasal decongestant |

Uses

- temporarily relieves common cold/flu symptoms
- nasal congestion
- sinus congestion and pressure
- minor aches and pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- cough due to minor throat and bronchial irritation

*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and DayQuil™ are registered trademarks of The Procter & Gamble Company.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

PARENTS:
Learn about your medicine at
www.StopMedicineAbuse.org

PLD-B340C LB010679

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if:

- adult takes more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount
- child takes more than 4 doses (15 mL each) in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

PEEL CORNER FOR MORE DRUG FACTS

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if the user has

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if user is taking the blood thinning drug warfarin.

Drug Facts (continued)

When using this product, do not take more than directed.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- take only as directed (see Overdose warning)
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL = milliliter
- keep dosing cup with product

| | |
|---------------------------------------|---------------------|
| adults and children 12 years and over | 30 mL every 4 hours |
| children 6 to under 12 years | 15 mL every 4 hours |
| children 4 to under 6 years | ask a doctor |
| children under 4 years | do not use |

when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information

- each 15 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

citric acid, FD&C yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

FAMILY WELLNESS Daytime Cold & Flu Relief

DAYTIME SEVERE COLD AND FLU RELIEF

acetaminophen dextromethorphan hbr guaifenesin, phenylephrine hcl liquid

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:55319-145 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------------------|--------------------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 325 mg in 15 mL |
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 10 mg in 15 mL |
| GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ) | GUAIFENESIN | 200 mg in 15 mL |
| PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) | PHENYLEPHRINE HYDROCHLORIDE | 5 mg in 15 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| FD&C YELLOW NO. 6 (UNII: H77VEI93A8) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| WATER (UNII: 059QF0KO0R) | |
| SORBITOL (UNII: 506T60A25R) | |
| SACCHARIN SODIUM (UNII: SB8ZUX40TY) | |
| SODIUM BENZOATE (UNII: OJ245FE5EU) | |
| SODIUM CHLORIDE (UNII: 451W47IQ8X) | |
| SODIUM CITRATE (UNII: 1Q73Q2JULR) | |
| XANTHAN GUM (UNII: TTV12P4NEE) | |
| SUCRALOSE (UNII: 96K6UQ3ZD4) | |
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:55319-145-08 | 237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 07/31/2024 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M012 | 07/31/2024 | |

Revised: 10/2025

Family Dollar (FAMILY WELLNESS)