

MIRALAX- polyethylene glycol 3350 powder, for solution
Cardinal Health 107, LLC

MiraLAX®

Drug Facts

Active ingredient (in each dose) (Bottle Only)

Polyethylene Glycol 3350, 17 g (cap filled to line)

Purpose

Osmotic Laxative

Active ingredient (in each dose) (Packet Only)

Polyethylene Glycol 3350, 17 g

Purpose

Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert

Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor of pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or you nausea, bloating or cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions (Bottle Only)

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
 - o use once a day
 - o fill to top of white section in cap which is marked to indicate the correct dose (17 g)
 - o stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - o do not combine with starch-based thickeners used for difficulty swallowing
 - o ensure that the powder is fully dissolved before drinking
 - o do not drink if there are any clumps
 - o do not use no more than 7 days
- children 16 years of age or under: ask a doctor

Directions (Packet Only)

- **do not take more than directed unless advised by your doctor**
- adults and children 17 years of age and older:
 - o use once a day
 - o stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - o do not combine with starch-based thickeners used for difficulty swallowing
 - o ensure that the powder is fully dissolved before drinking
 - o do not drink if there are any clumps
 - o do not use more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20°-25°C (68°-77°F)

- tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken

Inactive ingredients

none

Questions or comments?

1-800-MiraLAX (1-800-647-2529)

.

Overbagged with 6 x 17g packets per bag, NDC 55154-5035-6

WARNING: This Unit Dose package is not child resistant and is Intended for Institutional Use Only. Keep this and all drugs out of the reach of children.

Distributed By:**Cardinal Health**

Dublin, OH 43017

L4532222-11025 / L4532222-21025

Principal Display Panel

NDC 55154-5035-6

MiraLAX®

Polyethylene Glycol 3350,

Powder for Solution, Osmotic Laxative

6 PACKETS NET WT 0.5OZ (17g)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55154-5035-6	6 in 1 BAG	02/23/2009	
1		17 g in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA022015	02/23/2009	

Labeler - Cardinal Health 107, LLC (118546603)

Revised: 11/2025

Cardinal Health 107, LLC