ESOMEPRAZOLE MAGNESIUM - esomeprazole magnesium tablet, delayed release WALGREEN CO.

Esomeprazole Magnesium Delayed-Release Tablets 20 mg*

Drug Facts

Active ingredient (in each tablet)

*Esomeprazole 20 mg (Each film-coated delayed-release tablet corresponds to 21.75 mg esomeprazole magnesium dihydrate USP)

Purpose

Acid reducer

Uses

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert:

- Do not use if you are allergic to esomeprazole.
- Esomeprazole may cause severe skin reactions. Symptoms may include:
 - skin reddening blisters rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness**, **sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain.

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn

- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are

• taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- may take 1 to 4 days for full effect

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- swallow whole. Do not crush or chew tablets.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20° to 25°C (68° to 77°F)

Inactive ingredients

colloidal silicon dioxide, crospovidone, hydroxy propyl cellulose, hypromellose, low substituted hydroxy propyl cellulose, magnesium carbonate, magnesium oxide, methacrylic acid and ethyl acrylate copolymer dispersion (which contains copolymer based on ethyl acrylate and methacrylic acid, polysorbate 80 and sodium lauryl sulfate), microcrystalline cellulose, mono and di-glycerides, polyethylene glycol, polysorbate 80, red iron oxide, sodium stearyl fumarate, sugar spheres [which contains liquid glucose, starch (maize) and sucrose], talc, titanium dioxide, triethyl citrate and yellow iron oxide.

Questions or comments? call **1-855-274-4122** (Monday – Friday 8:30 AM to 5:00 PM EST)

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

DISTRIBUTED BY: WALGREEN CO.
DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com © 2024 Walgreen Co.

MADE IN INDIA

PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 20 mg (14 Tablets Bottle) Walgreens

Esomeprazole Magnesium
DELAYED-RELEASE TABLETS, 20 mg* / ACID REDUCER

24 Hour

14 TABLETS

ONE 14-DAY COURSE OF TREATMENT
MAY TAKE 1 TO 4 DAYS FOR FULL EFFECT
SEE NEW WARNING INFORMATION

Walgreens

Esomeprazole Magnesium

DELAYED-RELEASE TABLETS, 20 mg* / ACID REDUCER

24 Hour

14 TABLETS ONE 14-DAY COURSE OF TREATMENT MAY TAKE 1 TO 4 DAYS FOR FULL EFFECT SEE NEW WARNING INFORMATION

Do not use if seal imprinted with SEALED for YOUR PROTECTION under the bottle cap is broken or missing.

KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMAT DISTRIBUTED BY: WALGREEN CO. DEERFIELD, IL 60015 100% SATISFACTION GUARANTEED walgreens.com ©2024 Walgreen Co. MADE IN INDIA

MADE IN INDIA
Walgreens Pharmacist Recommended.
Our pharmacists recommend the Walgreens brand.
We invite you to compare to national brands.
ITEM 622889 W00000-0000-0
W3ORG0624-F

W3ORG0624-F

Active ingredient (in each tablet)
*Esomeprazole 20 mg

Purpose .Acid reducer (Each film-coated delayed-release tablet corresponds to 21.75 mg esomeprazole magnesium dihydrate USP)

Uses ■ treats frequent P1434591 heartburn (occurs 2 or more days a week) ■ not intended!

take 1 to 4 days for full

Unvarnished Zone (dotted line not for printing) 27 x 13 mm

Allergy alert: ■ Do not use if you are allergic to esomeprazole. ■ Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening blisters rank. If an allergic reaction occurs, stop use and seek medical help right away. Do not use if you have: trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath: sweating; pain spreading to arms, neck or shoulders; or lightheadedness frequent chest pain. These may be signs of a serious condition. See your doctor. Ask a

doctor before use if you have 🔳 had heartburn over 3 months. This may be a sign of a more serious condition. 🗷 frequent wheezing, particularly with heartburn unexplained weight loss nausea or vomiting stomach pain. Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs. Stop use and ask a doctor if your heartburn continues or worsens you need to take this product for more than 14 days 📫 you need to take more than 1 course of treatment every 4 months 💻 you get diarrhea 💻 you develop a rash or joint pain. If pregnant or breast-feeding, ask a health

1

professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Directions ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ may take 1 to 4 days for full effect. 14-Day Course of Treatment

swallow 1 tablet with a glass of water before eating in the morning

take every day for 14 days

do not take more than 1 tablet a day ■ swallow whole. Do not crush or chew tablets. ■ do not use for more than 14 days unless directed by your doctor Repeated 14-Day Courses (if needed) = you may repeat a 14-day course every 4 months = do not take for more than 14 days or more often than every 4 months unless directed by a doctor = children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition. Other information ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20° to 25°C (68° to 77°F) Questions or comments? call 1-855-274-4122 (Monday - Friday 8:30 AM to 5:00 PM EST) NDC 0363-0361-14 P1434591 LM-5854

PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 20 mg (42 Tablets Container Carton Label)

Walgreens

Compare to the active ingredient in Nexium[®] 24 HR^{††}

Esome prazole Magnesium **DELAYED-RELEASE TABLETS, 20 mg* / ACID REDUCER**

24 Hour **Tablets**

Treats frequent heartburn

42 TABLETS (3 BOTTLES OF 14 EACH) THREE 14-DAY COURSES OF TREATMENT MAY TAKE 1 TO 4 DAYS FOR FULL EFFECT

SEE NEW WARNING INFORMATION



ESOMEPRAZOLE MAGNESIUM

HYDROXYPROPYL CELLULOSE (20000 WAMW) (UNII: KZQ570MOA5)

esomeprazole magnesium tablet, delayed release					
Product Information					
Product Type	HUMAN OTC DRUG Item Code (Source) NDC:0363-0			-0361	
Route of Administration	ORAL				
Active Ingredient/Active Moiety					
Ingredient Name Basis of Strength					Strength
ESOMEPRAZOLE MAGNESIUM DIHYDRATE (UNII: 36H71644EQ) (ESOMEPRAZOLE - UNII:N3PA6559FT) ESOMEPRAZOLE				20 mg	
Inactive Ingredients					
	Ingredient Nam	ne			Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)					
SILICON DIOXIDE (ONII. LT)/20XD	IU4)				

HYPROMELLOSE 2910 (3 MPA.S) (UNII: 0VUT3PMY82) HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ 8WG20P6) LOW-SUBSTITUTED HYDROXYPROPYL CELLULOSE (11% HYDROXYPROPYL; 130000 MW) (UNII: 7773C1ROEU) MAGNESIUM CARBONATE (UNII: 0E53J927NA) MAGNESIUM OXIDE (UNII: 3A3U0GI71G) METHACRYLIC ACID - ETHYL ACRYLATE COPOLYMER (1:1) TYPE A (UNII: NX76LV5T8J) POLYSORBATE 80 (UNII: 60ZP39ZG8H) **SODIUM LAURYL SULFATE** (UNII: 368GB5141J) MICROCRYSTALLINE CELLULOSE 301 (UNII: W7YXH6D4BD) MICROCRYSTALLINE CELLULOSE 302 (UNII: 91B875MM4H) GLYCERYL MONO AND DIPALMITOSTEARATE (UNII: KC98RO82HJ) POLYETHYLENE GLYCOL 6000 (UNII: 30IQX730WE) FERRIC OXIDE RED (UNII: 1K09F3G675) **SODIUM STEARYL FUMARATE** (UNII: 7CV7WK4UI) DEXTROSE, UNSPECIFIED FORM (UNII: 1Y9XDZ35W2) STARCH, CORN (UNII: O8232NY3SJ) SUCROSE (UNII: C151H8M554) TALC (UNII: 7SEV7J4R1U) **TITANIUM DIOXIDE** (UNII: 15FIX9V2JP) TRIETHYL CITRATE (UNII: 8Z96QXD6UM) FERRIC OXIDE YELLOW (UNII: EX43802MRT)

Product Characteristics			
Color	PINK (Light pink)	Score	no score
Shape	OVAL (Oblong, Biconvex)	Size	13mm
Flavor		Imprint Code	K;12
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0361- 42	3 in 1 CARTON	07/08/2024	
1		14 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA214473	07/08/2024	

Labeler - WALGREEN CO. (008965063)

Establishment				
Name	Address	ID/FEI	Business Operations	
Aurobindo Pharma Limited		650381903	ANALYSIS(0363-0361), MANUFACTURE(0363-0361)	

Revised: 7/2024 WALGREEN CO.