

THERATEARS- carboxymethylcellulose sodium gel
MEDTECH PRODUCTS INC

TheraTears Liquid Night PF 58790-003

Drug Facts

Active ingredient (In each unit dose)

sodium carboxymethylcellulose 1%

Purpose

Eye lubricant

Uses

- As a lubricant to relieve dryness of the eye.
- As a protectant against further irritation of the eye.
- For temporary relief of burning, irritation, and discomfort including exposure to wind or sun.

Warnings

For external use only

- To avoid contamination do not touch tip of opened container to any surface. Do not reuse. Once opened discard. Use individual vials within 90 days of opening foil pouch.
- **This product contains no preservatives.** Any solution not used immediately after opening should be discarded. Re-use of this single-use product may lead to inflammation of the eye and/or discomfort, based on potential contamination during handling.

Do not use

- If solution changes color or becomes cloudy.

Stop use and ask a doctor if

- You experience eye pain, changes in vision, continued redness or irritation.
- Condition worsens or persists for more than 72 hours.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center immediately.

Directions

- To open, ***twist*** tab completely off.
- Instill 1 or 2 drops in the affected eye(s) as needed.

Other information

- Store between 15C-25C (59F-77F)
- Do not touch unit-dose tip to eye.

Inactive ingredients

boric acid, calcium chloride, magnesium chloride, potassium chloride, sodium bicarbonate, sodium borate, sodium chloride, sodium phosphate, water

Questions or comments? 1-800-579-8327

Principal Display Panel Text for Carton Label:

PRESERVATIVE

FREE

RECOMMENDED

DOCTOR

CREATED

thera

tears®

THERAPY FOR YOUR EYES®

Liquid Gel

nighttime

dry eye therapy

LUBRICANT EYE GEL

SOOTHING

OVERNIGHT

RELIEF

30 STERILE

Single-Use Vials* 0.60 FL OZ (18.0 mL) TOTAL

Drug Facts	
Active ingredient (in each unit-dose) Sodium carboxymethylcellulose 1%	Purpose Eye Lubricant
Uses	
<ul style="list-style-type: none"> As a lubricant to relieve dryness of the eye. As a protectant against further irritation of the eye. For temporary relief of burning, irritation, and discomfort including exposure to wind or sun. 	
Warnings	
For external use only.	
<ul style="list-style-type: none"> To avoid contamination, do not touch tip of opened container to any surface. Do not reuse. Once opened discard. Use individual vials within 90 days of opening for pouches. This product contains no preservatives. Any solution not used immediately after opening should be discarded. Re-use of this single-use product may lead to inflammation of the eye and/or discomfort, based on potential contamination during handling. 	
Do not use	
<ul style="list-style-type: none"> If solution changes color or becomes cloudy. 	
Stop use and ask a doctor if	
<ul style="list-style-type: none"> You experience eye pain, changes in vision, continued redness or irritation. Condition worsens or persists for more than 72 hours. 	
Keep out of reach of children, if swallowed, get medical help or contact a Poison Control Center right away.	
Directions	
<ul style="list-style-type: none"> To open, tear tab completely off. Instill 1 or 2 drops in the affected eye(s) as needed. 	
Other information	
<ul style="list-style-type: none"> Store between 15°C - 25°C (59°F - 77°F) Do not touch unit-dose tip to eye. 	
Inactive ingredients	
boric acid, calcium chloride, magnesium chloride, potassium chloride, sodium borate, sodium carbonate, sodium chloride, sodium phosphate, water	
Questions or comments? 1-800-579-0327	

Nighttime is the time when your eyes naturally rest and recover. But I have seen hundreds of frustrated dry-eye patients who told me their eyes still felt dry and irritated in the morning. That is why I developed TheraTears®. Try TheraTears® at bedtime and you will feel the difference in the morning.

Jeffrey P. Gilbard, MD
Jeffrey P. Gilbard, MD
Ophthalmologist

Part of the TheraTears® Dry Eye Therapy Line Formulated for Dry Eye Symptoms

Keep out of reach of children, if swallowed, get medical help or contact a Poison Control Center right away.

TAMPER EVIDENT:
Use only if foil pouch is sealed and single-use container is intact.

theratears.com

MADE IN FRANCE.
Dist. by Mediatech Products Inc., Tarrytown, NY 10591
A Prestige Consumer Healthcare company
©2024 TheraTears is owned by Mediatech Products Inc. All rights reserved. 71722001-04

PRESERVATIVE FREE

RECOMMENDED DOCTOR CREATED

thera tears®

RESTORES EYES NATURAL BALANCE™

Liquid Gel nighttime dry eye therapy

Sodium Carboxymethylcellulose 1%

Liquid Gel nighttime dry eye therapy

WITH OSMO-CORRECTION® HYPOTONIC & ELECTROLYTE BALANCED

Clinically Tested Formula
Replicates healthy tears

Preservative free for sensitive eyes

SOOTHING OVERNIGHT RELIEF

30 STERILE Single-Use Vials

0.6 FL OZ (18 mL) TOTAL

QP701848

Principal Display Panel Text for Carton Label:

PRESERVATIVE

FREE

RECOMMENDED

DOCTOR

CREATED

thera

tears®

THERAPY FOR YOUR EYES®

Liquid Gel

nighttime

dry eye therapy

LUBRICANT EYE GEL

SOOTHING

OVERNIGHT

RELIEF

4 Single-Use Vials 0.08 FL OZ (2.4 mL) TOTAL

PRESERVATIVE FREE

thera tears
THERAPY FOR YOUR EYES.
Liquid Gel
nighttime
dry eye therapy
LUBRICANT EYE GEL

SOOTHING OVERNIGHT RELIEF

STERILE
4 Single-Use Vials 0.08 FL OZ (2.4mL) TOTAL

Drug Facts

Active ingredient Purpose
Sodium carboxymethylcellulose 1%.....eye lubricant

Uses ■ As a lubricant to relieve dryness of the eye. ■ As a protectant against further irritation of the eye. ■ For temporary relief of burning, irritation, and discomfort including exposure to wind or sun. ▶

Drug Facts (continued)

Warnings For external use only
■ To avoid contamination do not touch tip of opened container to any surface. Do not reuse. Once opened discard. Use individual vials within 90 days of opening foil pouch.
Do not use if solution changes color or becomes cloudy. **Stop use and ask a doctor** if you experience eye pain, changes in vision, continued redness or irritation, or if condition worsens or persists for more than 72 hours. ▶

Drug Facts (continued)

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center immediately.

Directions ■ To open, twist tab completely off.
■ Instill 1 or 2 drops in the affected eye(s) as needed.

Other information ■ Use only if foil pouch is sealed and single-use container is intact.
■ Do not touch unit-dose tip to eye. ▶

Drug Facts (continued)

Inactive ingredients Borate buffers, calcium-chloride, magnesium chloride, potassium chloride, purified water, sodium bicarbonate, sodium chloride and sodium phosphate

Questions or comments? 1-800-579-8327

MADE IN USA theratears.com
© 2017 Alcon Consumer Health | A Division of Alcon, Inc.
Ann Arbor, MI 48105 T16-CA-01

THERATEARS

carboxymethylcellulose sodium gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58790-002
Route of Administration	OPHTHALMIC		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
carboxymethylcellulose sodium, unspecified form (UNII: K679OBS311) (carboxymethylcellulose - UNII:05JZ17B19X)	carboxymethylcellulose sodium, unspecified form	10 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
boric acid (UNII: R57Z HV85D4)	

sodium borate (UNII: 91MBZ8H3QO)	
calcium chloride (UNII: M4IOD6VV5M)	
magnesium chloride (UNII: 02F3473H9O)	
potassium chloride (UNII: 660YQ98I10)	
water (UNII: 059QF0KO0R)	
sodium bicarbonate (UNII: 8MDF5V39QO)	
sodium chloride (UNII: 451W47IQ8X)	
sodium phosphate (UNII: SE337SVY37)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58790-002-28	7 in 1 CARTON	12/01/2002	
1		0.6 mL in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M018	12/01/2002	

THERATEARS

carboxymethylcellulose sodium gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58790-003
Route of Administration	OPHTHALMIC		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
carboxymethylcellulose sodium, unspecified form (UNII: K679OBS311) (carboxymethylcellulose - UNII:05JZ17B19X)	carboxymethylcellulose sodium, unspecified form	10 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
boric acid (UNII: R57ZHV85D4)	
sodium borate (UNII: 91MBZ8H3QO)	
calcium chloride (UNII: M4IOD6VV5M)	
magnesium chloride (UNII: 02F3473H9O)	
potassium chloride (UNII: 660YQ98I10)	
water (UNII: 059QF0KO0R)	
sodium bicarbonate (UNII: 8MDF5V39QO)	

sodium chloride (UNII: 451W47IQ8X)

sodium phosphate (UNII: SE337SVY37)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58790-003-30	6 in 1 CARTON	12/07/2018	
1		5 in 1 POUCH		
1		0.6 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M018	12/07/2018	

Labeler - MEDTECH PRODUCTS INC (114707784)

Revised: 10/2025

MEDTECH PRODUCTS INC