NICOTINE POLACRILEX- nicotine polacrilex gum, chewing Rite Aid Corporation

Nicotine Polacrilex Gum Mint 2 mg and 4 mg - Walgreens

Drug Facts

Active ingredient (in each chewing piece) - 2 mg

Nicotine polacrilex 2 mg (nicotine)

Purpose

Stop smoking aid

Use

reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets.

Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions - 2 mg

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6 Weeks 7 to 9 Weeks 10 to 12

1 piece every	1 piece every	1 piece every
1 to 2 hours	2 to 4 hours	4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider

Other information - 2 mg

- each piece contains: calcium 115 mg, sodium 8 mg
- store at 20 25°C (68 77°F)
- protect from light

Inactive ingredients - 2 mg

acacia, acesulfame potassium, carnauba wax, flavors, gum base, hydroxypropyl cellulose,magnesium oxide, sodium bicarbonate, sodium carbonate, talc, titanium dioxide, xylitol.

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

PRINCIPAL DISPLAY PANEL - 2 mg

Compare to the active ingredient of Nicorette[®] Gum*

FREE | SUGAR

FROM | FREE

Nicotine Gum

Nicotine Polacrilex Gum USP,

2 mg (nicotine

Stop Smoking Aid

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP. If you smoke your first cigarette <u>WITHIN</u> 30 MINUTES of waking up, use Nicotine Polacrilex Gum USP, 4 mg

sugar free coated nicotine gum

TO INCREASE YOUR SUCCESS IN QUITTING:

- 1. You must be motivated to quit.
- 2. Use Enough Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
- 3. Use Long Enough Use Nicotine Polacrilex Gum for the full 12 weeks.
- 4. Use with a support program as directed in the enclosed User's Guide.

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.

- NOT FOR SALE TO THOSE UNDER 18 YEARS OF AGE.
- PROOF OF AGE REQUIRED.
- NOT FOR SALE IN VENDING MACHINES OR FROM ANY SOURCE WHERE PROOF OF AGE CANNOT BE VERIFIED.

*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® Gum.

DISTRIBUTED BY:

RITE AID, 30 HUNTER LANE,

CAMP HILL, PA 17011

Package label

TO INCREASE YOUR SUCCESS IN QUITTING:

FREE

FREM

SUGAR

You must be motivated to quit.
Use Enough - Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
Use Long Enough - Use Nicotine Polacrilex Gum for the full 12 weeks.

4. Use With a Support Program as directed in the enclosed User's Guide







NDC 11822-3325-5

mo

ACTUAL SIZE COATED MINT FLAVOR

ma

Directions

Compare to the active ingredient of Nicorette[®] Gum*

Nicotine Gum

Nicotine Polacrilex Gum USP, 2 mg (nicotine) Stop Smoking Aid

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP

If you smoke your first cigarette <u>WITHIN</u> 30 MINUTES of waking up, use Nicotine Polacrilex Gum USP, 4 mg

sugar free coated nicotine gum



PIECES 2 mg EACH

FREE SUGAR FREE FROM

NDC 11822-3325-5 Compare to the active ingredient of Nicorette[®] Gum^{*}

If you make the set of a gap, and a doctor before use a logar use and the regional regional controlled a logar use and der ingolaan radium fahrmation a logar use and der ingolaan radium fahrmation and you was under to gap and the set of a logar between a logar use and the radium and the set of a logar and you was and the radium and the set and the radium and the set of a logar and you was and the radium and the set and the radium and the set of a logar and the radium and the radium and the set of a logar and the set of a l

ware from Control Center (1-600-222-1222) right away.

event year must entrook to accord acted and entropy of the or the or the order of t

icotine Gum Nicotine Polacrilex Gum USP, 2 mg (nicotine) Stop Smoking Aid



moo.bieetin.www

DISTRIBUTED BY: RITE AID, 30 HUNTER LANE. САМР НІLL, РА 17011

This product is not manufactured or distributed by GiaxoSmithMine Consumer Healthcare, L.P., distributor of Nicoretta# Gum.

CSI 1-811-723-3832 Morday-Friday See-577-778-1165 Cluestions of comments?

flavors, gum base, hydroxypropy cellulose, magnesium oxide, aodium bicarbonate, aodium carbonate, talo, titanium dixode, xylitici



ABRA BAT SOMALLEWRUS THEFT



NIHTIM strengts strift not store up 1	Step use and sak a doctor if	Inactive ingredien	oq amiñuesce, acesuñiane po	Assium, cameuba wax,	
FOR THOSE WHO SMOKE THEIR FIRST CIGRAFETTE MORE THAN 30 MINUTES AFTER WHO IN 30 MINUTES	Asis a decior et prementa parte en en la construction de la construction encourse de la construction encourse en la construction encourse en la construction encourse en la construction encourse en la construction encourse encour	Other information a each piece contains: c a store at 20 -25°C (66 -7 a store at 20 -25°C (66 -7 a store at 20 -25°C (64 -7 a store at 20 -25°C (65 -7) a store at 20 -25°C (7) a store at 20 -25°	ê mulbee .gm ði t muisles	Su	
Sugar Sugar Free Materials	An a back reform an a from two is additional teach from the from	Satern 8 Satern 8 Satern 8 Satern 9 Satern 9 Sat	ucy, "sgriksto fineupati no g evi visuoretino da no da no da suan "muchanati, storuciali veb a second tali da un da no your h dali da da da no your h	scelq brocces a sev yem verthors rafts sceld erro scroth size sher with ro sou scroth and seven un a not mug ent seu or been un	commo lo variado aureo aviaro dago aureo aviaro dago (social dago) (social dago (social dago) (social dago (social dago) (social dago
ing (nicosing) bing prisoning bing prisoning qosi bing prisoning dosi bing prisoning	instances on the start from the second start of the second start o	cpewing a piece a do not est or drive for 1 drum. Whom the tinge is a cheek the process until machine Whom the tinge is mathing machine tingen machine machi	d II. Ansy next. Lengues. Then see privents rilood, cores ones si olgnö soft to ison i priverto ended selturin: 2.	patheen your cheek and in, until the tingle returns.	NINDERO KRAD JAN KITARI TANDANAN NINDERO KRAD JAN KITARI TANDANAN NINDERO KRAD JAN KITARI TANDANAN
Vicotine Polacrilex Gum USP,	cuiting amount including nicolae craving associated with a reduces withdrawal symptome, including nicolae craving, associated with USB	a oz 1 zarewy 7 pres every 7 pres a mus anticcin m 7 pres a mus anticcin m	Veeks 7 by 9 7 piece every 2 by 4 hours 2 by	ZT of DT askey t piece every 6 to 8 hours 6 to 8 hours 6 hours 7 hours	
Aicotine	aeoqru9 (aceida gniwento trase ini) traeliaargene word. Inome policater 2 mg (nootin) (m 2shonland enimelia	mug endooin gm S eau	griwoliol arti of gritmooe.	In the second se	
	stoen pund	пицисо) яроеу блид	(pan		



	m, chewing					
Product Informat	tion					
Product Type	HUMA	AN OTC DRUG	ltem Code	(Source)	NDC:11	822-3325
Route of Administra	tion BUCC	CAL				
Active Ingredient	Active Moie/	ty				
	Ingredient	Name		Basis of Stre	ength	Strength
NICOTINE (UNII: 6M3C89	9ZY6R) (NICOTINI	E - UNII:6M3C89ZY6	R)	NICOTINE		2 mg
Inactive Ingredie						
3						
	Ir	ngredient Name	•			Strength
ACACIA (UNII: 5C5403N2		ngredient Name	•			Strength
	260)		•			Strength
ACESULFAME POTASSI	260) IUM (UNII: 230V7		9			Strength
ACESULFAME POTASSI CARNAUBA WAX (UNII:	260) IUM (UNII: 230V7 R12CBM0EIZ)	73Q5G9)				Strength
ACESULFAME POTASSI CARNAUBA WAX (UNII: HYDROXYPROPYL CELI MAGNESIUM OXIDE (UN	260) IUM (UNII: 230V7 R12CBM0EIZ) LULOSE, UNSPE VII: 3A3U0GI71G)	73Q5G9) CIFIED (UNII: 9XZ8)				Strength
ACESULFAME POTASSI CARNAUBA WAX (UNII: HYDROXYPROPYL CELI MAGNESIUM OXIDE (UN SODIUM BICARBONATI	260) IUM (UNII: 230V7 R12CBM0EIZ) LULOSE, UNSPE NII: 3A3U0GI71G) E (UNII: 8MDF5V3	23Q5G9) CIFIED (UNII: 9XZ8) 9QO)				Strength
ACESULFAME POTASSI CARNAUBA WAX (UNII: HYDROXYPROPYL CELI MAGNESIUM OXIDE (UN SODIUM BICARBONATE SODIUM CARBONATE (260) IUM (UNII: 230V7 R12CBM0EIZ) LULOSE, UNSPE VII: 3A3U0GI71G) E (UNII: 8MDF5V3 UNII: 45P3261C7	23Q5G9) CIFIED (UNII: 9XZ8) 9QO)				Strength
ACESULFAME POTASSI CARNAUBA WAX (UNII: HYDROXYPROPYL CELI MAGNESIUM OXIDE (UN SODIUM BICARBONATE SODIUM CARBONATE (TALC (UNII: 7SEV7J4R1U)	260) IUM (UNII: 230V7 R12CBM0EIZ) LULOSE, UNSPE NII: 3A3U0GI71G) E (UNII: 8MDF5V3 UNII: 45P3261C7	23Q5G9) CIFIED (UNII: 9XZ8) 9QO)				Strength
ACACIA (UNII: 5C5403N2 ACESULFAME POTASSI CARNAUBA WAX (UNII: HYDROXYPROPYL CELI MAGNESIUM OXIDE (UN SODIUM BICARBONATE (TALC (UNII: 7SEV7J4R1U) TITANIUM DIOXIDE (UN	260) IUM (UNII: 230V7 R12CBM0EIZ) LULOSE, UNSPE VII: 3A3U0GI71G) E (UNII: 8MDF5V3 UNII: 45P3261C7) III: 15FIX9V2JP)	23Q5G9) CIFIED (UNII: 9XZ8) 9QO)				Strength
ACESULFAME POTASSI CARNAUBA WAX (UNII: HYDROXYPROPYL CELI MAGNESIUM OXIDE (UN SODIUM BICARBONATE SODIUM CARBONATE (TALC (UNII: 7SEV7J4R1U) TITANIUM DIOXIDE (UN	260) IUM (UNII: 230V7 R12CBM0EIZ) LULOSE, UNSPE VII: 3A3U0GI71G) E (UNII: 8MDF5V3 UNII: 45P3261C7) III: 15FIX9V2JP)	23Q5G9) CIFIED (UNII: 9XZ8) 9QO)				Strength
ACESULFAME POTASSI CARNAUBA WAX (UNII: HYDROXYPROPYL CELI MAGNESIUM OXIDE (UN SODIUM BICARBONATE SODIUM CARBONATE (TALC (UNII: 7SEV7J4R1U)	260) IUM (UNII: 230V7 R12CBM0EIZ) LULOSE, UNSPE NII: 3A3U0GI71G) E (UNII: 8MDF5V3 UNII: 45P3261C7) III: 15FIX9V2JP) (Q1E)	23Q5G9) CIFIED (UNII: 9XZ8) 9QO)				Strength
ACESULFAME POTASSI CARNAUBA WAX (UNII: HYDROXYPROPYL CELI MAGNESIUM OXIDE (UN SODIUM BICARBONATE SODIUM CARBONATE (TALC (UNII: 7SEV7J4R1U) TITANIUM DIOXIDE (UN XYLITOL (UNII: VCQ006K	260) IUM (UNII: 230V7 R12CBM0EIZ) LULOSE, UNSPE NII: 3A3U0GI71G) E (UNII: 8MDF5V3 UNII: 45P3261C7) III: 15FIX9V2JP) (Q1E)	23Q5G9) CIFIED (UNII: 9XZ8) 9QO)				
ACESULFAME POTASSI CARNAUBA WAX (UNII: HYDROXYPROPYL CELI MAGNESIUM OXIDE (UN SODIUM BICARBONATE SODIUM CARBONATE (TALC (UNII: 75EV7J4R1U) TITANIUM DIOXIDE (UN XYLITOL (UNII: VCQ006K	260) 10M (UNII: 230V7 R12CBM0EIZ) LULOSE, UNSPE VII: 3A3U0GI71G) E (UNII: 8MDF5V3 UNII: 45P3261C7) III: 15FIX9V2JP) (Q1E) Pristics	73Q5G9) CIFIED (UNII: 9XZ8 9QO) T)			no score 14mm	

Packaging			
# Item Code	Package Description	Marketing Start Date	Marketing End Date
NDC:11822- 3325-5	100 in 1 CARTON	12/27/2019	
1	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing	Information		
Marketing Marketing Category	Information Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

Labeler - Rite Aid Corporation (014578892)

Revised: 11/2023

Rite Aid Corporation