

VICKS DAYQUIL AND VICKS NYQUIL FOR PEOPLE WITH DIABETES COLD AND FLU- acetaminophen, dextromethorphan hbr, doxylamine succinate
The Procter & Gamble Manufacturing Company

VICKS® DayQuil™ NyQuil™ FOR PEOPLE WITH DIABETES COLD & FLU
Liquid Convenience Pack

NyQuil FOR PEOPLE WITH DIABETES COLD & FLU

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg
Dextromethorphan HBr 30 mg
Doxylamine succinate 12.5 mg

Purpose

Pain reliever/ fever reducer
Cough suppressant
Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever
- runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 30 mL dose contains:**sodium 28.23 mg
- store at no greater than 25°C (77°F) and do not refrigerate

Inactive ingredients

citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

Questions?

1-800-362-1683

TAMPER EVIDENT:

DO NOT USE IF PRINTED SHRINKBAND SEAL AROUND THE NECK IS BROKEN OR MISSING.

DIST. BY: PROCTER & GAMBLE,

CINCINNATI, OH 45202

DayQuil FOR PEOPLE WITH DIABETES COLD & FLU

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg
Dextromethorphan HBr 20 mg

Purpose

Pain reliever/Fever reducer
Cough suppressant

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation

- sore throat
- headache
- minor aches & pains
- fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions.

Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist .
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain or cough get worse or last more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 30 mL contains:** sodium 29 mg
- store at no greater than 25 °C (77 °F) and do not refrigerate

Inactive ingredients

citric acid, FD&C yellow no. 6, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

Questions?

1-800-362-1683

TAMPER EVIDENT: DO NOT USE IF PRINTED SHRINKBAND SEAL AROUND THE NECK IS BROKEN OR MISSING

**DIST. BY: PROCTER & GAMBLE,
CINCINNATI, OH 45202**

PRINCIPAL DISPLAY PANEL - Kit Carton

SUGAR FREE

VALUE PACK

Vicks DayQuil and NyQuil FOR PEOPLE WITH DIABETES

COLD & FLU

Vicks DayQuil FOR PEOPLE WITH DIABETES

COLD & FLU

THE DAYTIME, COUGHING, ACHING, SORE THROAT, FEVER, POWER THROUGH YOUR DAY, MEDICINE.

Acetaminophen, Dextromethorphan HBr

Headache, Fever, Sore Throat, Minor Aches & Pains

Cough

Non-Drowsy

Alcohol Free

Vicks NyQuil FOR PEOPLE WITH DIABETES

COLD & FLU

THE NIGHTTIME, SNIFFLING, SNEEZING, COUGHING, ACHING, FEVER, BEST SLEEP WITH A COLD, MEDICINE.

Acetaminophen, Doxylamine Succinate, Dextromethorphan HBr

Headache, Fever, Sore Throat, Minor Aches & Pains

Sneezing, Runny Nose

Cough

Nighttime Relief

Alcohol Free

ALSO SOLD AS DAYQUIL/NYQUIL HIGH BLOOD PRESSURE

2 BOTTLES - 1 DAYQUIL/1 NYQUIL 12 FL OZ (354 mL) EACH; TOTAL 24 FL OZ (708 mL)



VICKS DAYQUIL AND VICKS NYQUIL FOR PEOPLE WITH DIABETES COLD AND FLU

acetaminophen, dextromethorphan hbr, doxylamine succinate kit

Product Information				
Product Type		HUMAN OTC DRUG	Item Code (Source)	NDC:69423-845
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69423-845-24	1 in 1 PACKAGE; Type 0: Not a Combination Product	07/09/2024	
Quantity of Parts				
Part #	Package Quantity		Total Product Quantity	
Part 1	1 BOTTLE, PLASTIC		354 mL	
Part 2	1 BOTTLE, PLASTIC		354 mL	

Part 1 of 2

VICKS DAYQUIL FOR PEOPLE WITH DIABETES COLD AND FLU

acetaminophen, dextromethorphan hbr liquid

Product Information

Item Code (Source)	NDC:69423-843
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
XANTHAN GUM (UNII: TTV12P4NEE)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	orange	Score	
Shape		Size	
Flavor	CITRUS	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69423-843-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/09/2024	

Part 2 of 2

VICKS NYQUIL FOR PEOPLE WITH DIABETES COLD AND FLU

acetaminophen, dextromethorphan hbr, doxylamine succinate liquid

Product Information

Item Code (Source)	NDC:69423-844
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
SUCRALOSE (UNII: 96K6UQ3ZD4)	
SORBITOL (UNII: 506T60A25R)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
XANTHAN GUM (UNII: TTV12P4NEE)	
GLYCERIN (UNII: PDC6A3C0OX)	

Product Characteristics

Color	red	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	

Contains**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69423-844-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/09/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/09/2024	

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Revised: 5/2024

The Procter & Gamble Manufacturing Company