

PHENOBARBITAL- phenobarbital tablet
WINDER LABORATORIES, LLC

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

PHENOBARBITAL TABLETS, USP CIV

Rx Only


Rev. Jan 2022

DESCRIPTION

The barbiturates are nonselective central nervous system (CNS) depressants primarily used as sedative-hypnotics. In subhypnotic doses, they are also used as anticonvulsants. The barbiturates and their sodium salts are subject to control under the Federal Controlled Substances Act.

Phenobarbital is a barbituric acid derivative for oral administration and occurs as a white, odorless, slightly bitter powder that is soluble in chloroform, freely soluble in alcohol or ether, and slightly soluble in water. Its saturated solution has a pH of about 5.6. Chemically, it is 5-ethyl-5-phenylbarbituric acid with the molecular formula $C_{12}H_{12}N_2O_3$ (232.24).

The structural formula is as follows:



Each Phenobarbital Tablet, USP contains 15 mg, 16.2 mg, 30 mg, 32.4 mg, 60 mg, 64.8 mg, 97.2 mg, or 100 mg of phenobarbital, USP.

Inactive ingredients are as follows:

Microcrystalline Cellulose; Lactose Monohydrate; Sodium Starch Glycolate, Type-A; Colloidal Silicon Dioxide; Magnesium Stearate

CLINICAL PHARMACOLOGY

Phenobarbital, a long-acting barbiturate, is a central nervous system depressant. In ordinary doses, the drug acts as a sedative and anticonvulsant. Its onset of action occurs within 30 minutes, and the duration of action ranges from 5 to 6 hours. It is detoxified in the liver.

INDICATIONS AND USAGE

Phenobarbital Tablets, USP are indicated for use as a sedative or anticonvulsant.

CONTRAINDICATIONS

Phenobarbital is contraindicated in patients who are hypersensitive to barbiturates. In such patients, severe hepatic damage can occur from ordinary doses and is usually associated with dermatitis and involvement of parenchymatous organs. A personal or familial history of acute intermittent porphyria represents one of the few absolute contraindications to the use of barbiturates. Phenobarbital is also contraindicated in patients with marked impairment of liver function, or respiratory disease in which dyspnea or obstruction is evident. It should not be administered to persons with known previous addiction to the sedative/hypnotic group, since ordinary doses may be ineffectual and may contribute to further addiction.

WARNINGS

In small doses, the barbiturates may increase the reaction to painful stimuli. Taken by themselves, the barbiturates cannot be relied upon to relieve pain or even to produce sedation or sleep in the presence of severe pain.

PRECAUTIONS

General Precautions

Barbiturates induce liver microsomal enzyme activity. This accelerates the biotransformation of various drugs and is probably part of the mechanism of the tolerance encountered with barbiturates.

Phenobarbital, therefore, should be used with caution in patients with decreased liver function. This drug should also be administered cautiously to patients with a history of drug dependence or abuse (see **DRUG ABUSE AND DEPENDENCE**).

Phenobarbital may decrease the potency of coumarin anticoagulants; therefore, patients receiving such concomitant therapy should have more frequent prothrombin determinations. As with other sedatives and hypnotics, elderly or debilitated patients may react to barbiturates with marked excitement, depression, or confusion.

The systemic effects of exogenous hydrocortisone and endogenous hydrocortisone (cortisol) may be diminished by phenobarbital. Thus, this product should be administered with caution to patients with borderline hypoadrenal function, regardless of whether it is of pituitary or of primary adrenal origin.

Information for Patients

Phenobarbital may impair the mental and/or physical abilities required for the performance of potentially hazardous tasks, such as driving a car or operating machinery. The patient should be cautioned accordingly.

Drug Interactions

Phenobarbital in combination with alcohol, tranquilizers, and other central nervous system depressants has additive depressant effects, and the patient should be so advised. Patients taking this drug should be warned not to exceed the dosage recommended by their physician. Toxic effects and fatalities have occurred following overdoses of phenobarbital alone and in combination with other central nervous system depressants. Caution should be exercised in prescribing unnecessarily large amounts of phenobarbital for patients who have a history of emotional disturbances or suicidal ideation or who have misused alcohol and other CNS drugs (see **OVERDOSAGE**).

Usage in Pregnancy

Pregnancy Category B

Reproduction studies have been performed in animals and have revealed no evidence of impaired fertility or harm to the fetus due to phenobarbital. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed and under the supervision of a physician .

Nursing Mothers

Caution should be exercised when phenobarbital is administered to a nursing woman.

ADVERSE REACTIONS

The following adverse reactions have been reported:

CNS Depression:

Residual sedation or "hangover," drowsiness, lethargy, and vertigo. Emotional disturbances and phobias may be accentuated. In some persons, barbiturates such as phenobarbital repeatedly produce excitement rather than depression, and the patient may appear to be inebriated. Like other nonanalgesic hypnotic drugs, barbiturates, such as phenobarbital, when given in the presence of pain, may cause restlessness, excitement, and even delirium. Rarely, the use of barbiturates results in localized or diffused myalgic, neuralgic, or arthritic pain, especially in psychoneurotic patients with insomnia. The pain may appear in paroxysms, is most intense in the early morning hours, and is most frequently located in the region of the neck, shoulder girdle, and upper limbs. Symptoms may last for days after the drug is discontinued.

Respiratory/Circulatory:

Respiratory depression, apnea, circulatory collapse.

Allergic:

Acquired hypersensitivity to barbiturates consists chiefly in allergic reactions that occur especially in persons who tend to have asthma, urticaria, angioedema, and similar

conditions. Hypersensitivity reactions in this category include localized swelling, particularly of the eyelids, cheeks, or lips, and erythematous dermatitis. Rarely, exfoliative dermatitis (e.g., Stevens-Johnson syndrome and toxic epidermal necrolysis) may be caused by phenobarbital and can prove fatal. The skin eruption may be associated with fever, delirium, and marked degenerative changes in the liver and other parenchymatous organs. In a few cases, megaloblastic anemia has been associated with the chronic use of phenobarbital.

Other:

Nausea and vomiting; headache.

To report SUSPECTED ADVERSE REACTIONS, contact Winder Laboratories, LLC at 1-770-307-0703, or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG ABUSE AND DEPENDENCE

Controlled Substance

Phenobarbital is a Schedule IV drug.

Dependence

Prolonged, uninterrupted use of barbiturates (particularly the short-acting drugs), even in therapeutic doses, may result in psychic and physical dependence. Withdrawal symptoms due to physical dependence following chronic use of large doses of barbiturates may include delirium, convulsions, and death.

OVERDOSAGE

The signs and symptoms of barbiturate poisoning are referable especially to the central nervous system and the cardiovascular system. Moderate intoxication resembles alcoholic inebriation. In severe intoxication, the patient is comatose, the level of reflex activity conforming in a general way to the intensity of the central depression. The deep reflexes may persist for some time despite coexistent coma. The Babinski sign is often positive. The EEG may be of the "burst-suppression" type, with brief periods of electrical silence. The pupils may be constricted and react to light, but late in the course of barbiturate poisoning they may show hypoxic paralytic dilatation. Respiration is affected early.

Breathing may be either slow or rapid and shallow; Cheyne-Stokes rhythm may be present. Respiratory minute volume is diminished, and hypoxia and respiratory acidosis may develop. The blood pressure falls, owing partly to depression of medullary vasomotor centers; partly to a direct action of the drug on the myocardium, sympathetic ganglia, and vascular smooth muscle; partly to hypoxia.

The patient thus develops a typical shock syndrome, with a weak and rapid pulse, cold and clammy skin, and a rise in the hematocrit. Respiratory complications (atelectasis, pulmonary edema, and bronchopneumonia) and renal failure are much dreaded and not infrequent concomitant of severe barbiturate poisoning. There is usually hypothermia, sometimes with temperatures as low as 32°C.

Treatment

General management should consist of symptomatic and supportive therapy, including gastric lavage, administration of intravenous fluids, and maintenance of blood pressure, body temperature and adequate respiratory exchange. Dialysis will increase the rate of removal of barbiturates from the body fluids. Antibiotics may be required to control pulmonary complications.

DOSAGE AND ADMINISTRATION

Oral Sedative Dose, Adults: 30 to 120 mg daily in 2 or 3 divided doses.

Children: 6 mg/kg of body weight daily in 3 divided doses.

Oral Hypnotic Dose, Adults: 100 to 320 mg.

Oral Anticonvulsant Dose, Adults: 50 to 100 mg 2 or 3 times daily.

Children: 15 to 50 mg 2 or 3 times daily.

HOW SUPPLIED

Phenobarbital Tablets, USP 15 mg:

White, Round Tablet; Debossed WL on one side and 136 on the other side.

Bottles of 90 tablets NDC 75826-136-90
Bottles of 100 tablets NDC 75826-136-10
Bottles of 500 tablets NDC 75826-136-50
Bottles of 1000
tablets NDC 75826-136-00

Phenobarbital Tablets, USP 16.2 mg ($\frac{1}{4}$ grain):

White, Round, Scored Tablet; Debossed WL on one side and 137 on the other side.

Bottles of 90 tablets NDC 75826-137-90
Bottles of 100 tablets NDC 75826-137-10
Bottles of 500 tablets NDC 75826-137-50
Bottles of 1000
tablets NDC 75826-137-00

Phenobarbital Tablets, USP 30 mg:

White, Round, Scored Tablet; Debossed WL on one side and 138 on the other side.

Bottles of 90 tablets NDC 75826-138-90
Bottles of 100 tablets NDC 75826-138-10
Bottles of 500 tablets NDC 75826-138-50

Bottles of 1000 tablets NDC 75826-138-00

Phenobarbital Tablets, USP 32.4 mg (½ grain):

White, Round, Scored Tablet; Debossed WL on one side and 139 on the other side.

Bottles of 90 tablets NDC 75826-139-90
Bottles of 100 tablets NDC 75826-139-10
Bottles of 500 tablets NDC 75826-139-50
Bottles of 1000 tablets NDC 75826-139-00

Phenobarbital Tablets, USP 60 mg:

White, Round Tablet; Debossed WL on one side and 140 on the other side.

Bottles of 90 tablets NDC 75826-140-90
Bottles of 100 tablets NDC 75826-140-10
Bottles of 500 tablets NDC 75826-140-50
Bottles of 1000 tablets NDC 75826-140-00

Phenobarbital Tablets, USP 64.8 mg (1 grain):

White, Round, Scored Tablet; Debossed WL on one side and 141 on the other side.

Bottles of 90 tablets NDC 75826-141-90
Bottles of 100 tablets NDC 75826-141-10
Bottles of 500 tablets NDC 75826-141-50
Bottles of 1000 tablets NDC 75826-141-00

Phenobarbital Tablets, USP 97.2 mg (1½ grain):

White, Round, Scored Tablet; Debossed WL on one side and 142 on the other side.

Bottles of 90 tablets NDC 75826-142-90
Bottles of 100 tablets NDC 75826-142-10
Bottles of 500 tablets NDC 75826-142-50
Bottles of 1000 tablets NDC 75826-142-00

Phenobarbital Tablets, USP 100 mg:

White, Round, Scored Tablet; Debossed WL on one side and 143 on the other side.

Bottles of 90 tablets NDC 75826-143-90
Bottles of 100 tablets NDC 75826-143-10
Bottles of 500 tablets NDC 75826-143-50
Bottles of 1000 tablets NDC 75826-143-00

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. Protect from light and moisture.

Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure.

Manufactured by:
Winder Laboratories, LLC.
Winder, GA 30680

RLS.135.99-1.0
Updated: Jan 2022

PRINCIPAL DISPLAY PANEL - 15 mg Tablet Bottle Label

NDC 75826-136-10
winder[®]
LABS

Phenobarbital
Tablets, USP
CIV

15 mg

WARNING: May be habit forming.

Each tablet contains:
Phenobarbital, USP
15 mg

Rx Only
100 Tablets

NDC 75826-136-10 **winder[®]**
LABS

Phenobarbital
Tablets, USP **CIV**

15 mg

WARNING: May be habit forming.

Each tablet contains:
Phenobarbital, USP..... 15 mg

Rx Only **100 Tablets**

USUAL ADULT DOSAGE: See accompanying product literature for complete information.
Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. Protect from light and moisture. Keep container tightly closed.
Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure.
KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN.

© 2018 Winder Laboratories, LLC.
® Marks are the property of Winder Laboratories, LLC.
Manufactured by:
Winder Laboratories, LLC., Winder, GA 30680
RLS.136.10 08/18 50301US10

GTIN 00375826136104
N 3 75826 13610 4

PRINCIPAL DISPLAY PANEL - 16.2 mg Tablet Bottle Label

NDC 75826-137-10

winder[®] LABS

Phenobarbital Tablets,
USP CIV

16.2 mg

WARNING: May be habit forming.

Each tablet contains:
Phenobarbital, USP
16.2 mg

Rx Only
100 Tablets

NDC 75826-137-10 **winder[®]**
LABS

Phenobarbital
Tablets, USP **CIV**

16.2 mg (1/4 grain)

WARNING: May be habit forming.

Each tablet contains:
Phenobarbital, USP..... 16.2 mg (1/4 grain)

Rx Only **100 Tablets**

Serialization & Lot Coding Area
14.7mm x 34mm

USUAL ADULT DOSAGE: See accompanying product literature for complete information.
Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. Protect from light and moisture. Keep container tightly closed.
Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure.
KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN.

© 2018 Winder Laboratories, LLC.
® Marks are the property of Winder Laboratories, LLC.
Manufactured by:
Winder Laboratories, LLC., Winder, GA 30680
RLS.137.10 05/18 50302US10

GTIN 00375826137101
N 3 75826 13710 1

PRINCIPAL DISPLAY PANEL - 30 mg Tablet Bottle Label

NDC 75826-138-10

winder[®]
LABS

Phenobarbital
Tablets, USP
CIV

30 mg

WARNING: May be habit forming.

Each tablet contains:
Phenobarbital, USP
30 mg

Rx Only
100 Tablets

NDC 75826-138-10

winder[®]
LABS

Phenobarbital
Tablets, USP

30 mg

WARNING: May be habit forming.

Each tablet contains:
Phenobarbital, USP..... 30 mg

Rx Only **100 Tablets**

USUAL ADULT DOSAGE: See accompanying product literature for complete information.
Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. Protect from light and moisture. Keep container tightly closed.
Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure.
KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN.

© 2018 Winder Laboratories, LLC.
® Marks are the property of Winder Laboratories, LLC.
Manufactured by:
Winder Laboratories, LLC., Winder, GA 30680
RLS.138.10 08/18 50303US10

GTIN 00375826138108

75826 13810 8

N 3

PRINCIPAL DISPLAY PANEL - 32.4 mg Tablet Bottle Label

NDC 75826-139-10
winder[®] LABS

Phenobarbital Tablets,
USP CIV

32.4 mg

WARNING: May be habit forming.

Each tablet contains:
Phenobarbital, USP
32.4 mg

Rx Only
100 Tablets

NDC 75826-139-10

winder
LABS

Phenobarbital
Tablets, USP

32.4 mg (1/2 grain)

WARNING: May be habit forming.

Each tablet contains:
Phenobarbital, USP.....32.4 mg (1/2 grain)

Rx Only **100 Tablets**

Serialization & Lot Coding Area
14.7mm x 34mm

USUAL ADULT DOSAGE: See accompanying product literature for complete information.
Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. Protect from light and moisture. Keep container tightly closed.
Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure.
KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN.

© 2018 Winder Laboratories, LLC.
© Marks are the property of Winder Laboratories, LLC.
Manufactured by:
Winder Laboratories, LLC., Winder, GA 30680
PLS.139.10 05/18 50304US10

GTIN 00375826139105



N 3 75826 13910 5

PRINCIPAL DISPLAY PANEL - 60 mg Tablet Bottle Label

NDC 75826-140-10

winder®
LABS

Phenobarbital
Tablets, USP
CIV

60 mg

WARNING: May be habit forming.

Each tablet contains:
Phenobarbital, USP
60 mg

Rx Only
100 Tablets

NDC 75826-140-10

winder
LABS

Phenobarbital
Tablets, USP

60 mg

WARNING: May be habit forming.

Each tablet contains:
Phenobarbital, USP..... 60 mg

Rx Only **100 Tablets**

USUAL ADULT DOSAGE: See accompanying product literature for complete information.
Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. Protect from light and moisture. Keep container tightly closed.
Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure.
KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN.

© 2018 Winder Laboratories, LLC.
© Marks are the property of Winder Laboratories, LLC.
Manufactured by:
Winder Laboratories, LLC. RLS.140.10 08/18
Winder, GA 30680 50405US10

GTIN 00375826140101



N 3 75826 14010 1

PRINCIPAL DISPLAY PANEL - 64.8 mg Tablet Bottle Label

NDC 75826-141-10

winder[®] LABS

Phenobarbital Tablets,
USP CIV

64.8 mg

WARNING: May be habit forming.

Each tablet contains:
Phenobarbital, USP
64.8 mg

Rx Only
100 Tablets

NDC 75826-141-10

winder[®]
LABS

Phenobarbital
Tablets, USP CIV

64.8 mg (1 grain)

WARNING: May be habit forming.

Each tablet contains:
Phenobarbital, USP.....64.8 mg (1 grain)

Rx Only **100 Tablets**

Serialization & Lot Coding Area
At Least 18mm x 58mm

USUAL ADULT DOSAGE: See accompanying product literature for complete information.
Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. Protect from light and moisture. Keep container tightly closed.
Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure.
KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN.
© 2018 Winder Laboratories, LLC.
® Marks are the property of Winder Laboratories, LLC.
Manufactured by:
Winder Laboratories, LLC.
Winder, GA 30680

GTIN 00375826141108
N 3 75826 14110 8
RLS.141.10 05/18
50406US10

PRINCIPAL DISPLAY PANEL - 100 mg Tablet Bottle Label

NDC 75826-143-10

winder[®]
LABS

Phenobarbital
Tablets, USP
CIV

100 mg

WARNING: May be habit forming.

Each tablet contains:
Phenobarbital, USP
100 mg

Rx Only
100 Tablets

NDC 75826-143-10

winder[®]
LABS

Phenobarbital
Tablets, USP CIV

100 mg

WARNING: May be habit forming.

Each tablet contains:
Phenobarbital, USP.....100 mg

Rx Only **100 Tablets**

USUAL ADULT DOSAGE: See accompanying product literature for complete information.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. Protect from light and moisture. Keep container tightly closed.

Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure.

KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN.

© 2018 Winder Laboratories, LLC.
® Marks are the property of Winder Laboratories, LLC.

Manufactured by:
Winder Laboratories, LLC. RLS.143.10 08/18
Winder, GA 30680 50408US10

GTIN 00375826143102

7582614310

N 3

PRINCIPAL DISPLAY PANEL - 97.2 mg Tablet Bottle Label

NDC 75826-142-10
winder[®] LABS

Phenobarbital Tablets,
USP CIV

97.2 mg

WARNING: May be habit forming.

Each tablet contains:
Phenobarbital, USP
97.2 mg

Rx Only
100 Tablets

NDC 75826-142-10

winder
LABS

Phenobarbital
Tablets, USP 

97.2 mg (1½ grain)

WARNING: May be habit forming.

Each tablet contains:
Phenobarbital, USP.....97.2 mg (1½ grain)

Rx Only

100 Tablets

Serialization &
Lot Coding Area
At Least 18mm x 58mm

USUAL ADULT DOSAGE: See accompanying product literature for complete information.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. Protect from light and moisture. Keep container tightly closed.

Dispense in a tight, light-resistant container as defined in the USP using a child-resistant closure.

KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN.

© 2018 Winder Laboratories, LLC.
® Marks are the property of Winder Laboratories, LLC.

Manufactured by:
Winder Laboratories, LLC.
Winder, GA 30680

RLS.142.10 05/18
50407US10



PHENOBARBITAL

phenobarbital tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:75826-136
Route of Administration	ORAL	DEA Schedule	CIV

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENOBARBITAL (UNII: YQE403BP4D) (PHENOBARBITAL - UNII:YQE403BP4D)	PHENOBARBITAL	15 mg

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	WL;136
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:75826-136-10	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/01/2018	
2	NDC:75826-136-50	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/01/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		10/01/2018	

PHENOBARBITAL

phenobarbital tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:75826-137
Route of Administration	ORAL	DEA Schedule	CIV

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENOBARBITAL (UNII: YQE403BP4D) (PHENOBARBITAL - UNII:YQE403BP4D)	PHENOBARBITAL	16.2 mg

Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	6mm
Flavor		Imprint Code	W;L;137
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:75826-	100 in 1 BOTTLE, PLASTIC; Type 0: Not a	06/11/2018	

1	137-10	Combination Product	06/11/2019	
2	NDC:75826-137-00	1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/11/2019	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		06/11/2019	

PHENOBARBITAL				
phenobarbital tablet				
Product Information				
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:75826-138	
Route of Administration	ORAL	DEA Schedule	CIV	
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
PHENOBARBITAL (UNII: YQE403BP4D) (PHENOBARBITAL - UNII:YQE403BP4D)		PHENOBARBITAL	30 mg	
Inactive Ingredients				
Ingredient Name			Strength	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)				
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)				
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
Product Characteristics				
Color	white	Score	2 pieces	
Shape	ROUND	Size	6mm	
Flavor		Imprint Code	W;L;138	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:75826-138-10	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/01/2018	
2	NDC:75826-138-50	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/01/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		10/01/2018	

PHENOBARBITAL

phenobarbital tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:75826-139
Route of Administration	ORAL	DEA Schedule	CIV

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENOBARBITAL (UNII: YQE403BP4D) (PHENOBARBITAL - UNII:YQE403BP4D)	PHENOBARBITAL	32.4 mg

Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	6mm
Flavor		Imprint Code	W;L;139
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:75826-139-10	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/11/2019	
2	NDC:75826-139-00	1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/11/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		06/11/2019	

PHENOBARBITAL

phenobarbital tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:75826-140
Route of Administration	ORAL	DEA Schedule	CIV

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENOBARBITAL (UNII: YQE403BP4D) (PHENOBARBITAL - UNII:YQE403BP4D)	PHENOBARBITAL	60 mg

Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	WL;140
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:75826-140-10	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/01/2018	
2	NDC:75826-140-50	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/01/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

unapproved drug other		10/01/2018	
--------------------------	--	------------	--

PHENOBARBITAL

phenobarbital tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:75826-141
Route of Administration	ORAL	DEA Schedule	CIV

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENOBARBITAL (UNII: YQE403BP4D) (PHENOBARBITAL - UNII:YQE403BP4D)	PHENOBARBITAL	64.8 mg

Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	8mm
Flavor		Imprint Code	W;L;141
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:75826-141-10	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/11/2019	
2	NDC:75826-141-00	1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/11/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		06/11/2019	

PHENOBARBITAL

phenobarbital tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:75826-142
Route of Administration	ORAL	DEA Schedule	CIV

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENOBARBITAL (UNII: YQE403BP4D) (PHENOBARBITAL - UNII:YQE403BP4D)	PHENOBARBITAL	97.2 mg

Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	W;L;142
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:75826-142-10	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/11/2019	
2	NDC:75826-142-00	1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/11/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		06/11/2019	

PHENOBARBITAL

phenobarbital tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:75826-143
Route of Administration	ORAL	DEA Schedule	CIV

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENOBARBITAL (UNII: YQE403BP4D) (PHENOBARBITAL - UNII:YQE403BP4D)	PHENOBARBITAL	100 mg

Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	W;L;143
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:75826-143-10	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	10/01/2018	
2	NDC:75826-143-50	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/01/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		10/01/2018	

Labeler - WINDER LABORATORIES, LLC (965195170)

Establishment

Name	Address	ID/FEI	Business Operations
WINDER LABORATORIES, LLC		965195170	manufacture(75826-136, 75826-137, 75826-138, 75826-139, 75826-140, 75826-141, 75826-142, 75826-143)

Revised: 5/2022

WINDER LABORATORIES, LLC