

# **ACETAMINOPHEN- acetaminophen tablet**

## **Meijer**

-----

### **Acetaminophen Red EZ Tablets**

#### **Pain Reliever/Fever Reducer FOR ADULTS**

#### **Active Ingredient (in each coated tablet)**

Acetaminophen USP, 500mg

#### **Purpose**

Pain reliever/fever reducer

#### **Uses**

□ temporarily relieves minor aches and pains due to:

- the common cold □ headache
- backache □ minor pain of arthritis
- toothache □ muscular aches
- premenstrual and menstrual cramps

□ temporarily reduces fever

#### **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening □ blisters □ rash

If a skin reaction occurs, stop use and seek medical help right away

#### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have**

liver disease

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition

**If pregnant or breast-feeding,**

ask a health professional before use

**Keep out of reach of children**

Keep out of reach of children

**Overdose warning:**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed (see overdose warning)**

adults and children 12  take 2 tablets every 6 hours while symptoms last  
years and over  do not take more than 6 tablets in 24 hours, unless directed by a  
doctor

- do not use for more than 10 days unless directed by a doctor

children under  ask a doctor  
12 years

**Other information**

- store at 20°-25°C (68°-77°F). See USP Controlled Room Temperature.

□ see bottom of the label for lot number and expiration date

## Inactive ingredients

FD&C red no. 40, FD&C yellow no. 6, lecithin, polyethylene glycol, polyvinyl alcohol, povidone k-30, pregelatinized starch, stearic acid, sucralose, talc, titanium dioxide

## Questions or comments?

call **1-877-770-3183** Mon-Fri 8:00 AM EST to 5:00 PM PST

## PDP

**meijer.**  
Extra Strength  
**Pain Relief**

NDC 79481-5380-2

COMPARE TO EXTRA STRENGTH  
TYLENOL® ACTIVE INGREDIENT\*

**ACETAMINOPHEN TABLETS USP 500 mg**

Pain Reliever/Fever Reducer  
Contains No Aspirin  
FOR ADULTS

225 COATED TABLETS  
500 MG EACH

Actual Size

IMPORTANT: READ THE DIRECTIONS AND WARNINGS BEFORE USE  
TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

**Drug Facts**  
**Active ingredient (in each coated tablet)**  
Acetaminophen USP, 500 mg... Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - minor pain of arthritis
  - toothache
  - muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**  
**Liver warnings:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- hives
- skin rash
- blisters

If a skin reaction occurs, stop use and seek medical help right away

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if you have liver disease**

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

**Other information**

- store at 20°-25°C (68°-77°F). See USP Controlled Room Temperature.
- see bottom of the label for lot number and expiration date

**Inactive ingredients**  
FD&C red no. 40, FD&C yellow no. 6, lecithin, polyethylene glycol, polyvinyl alcohol, povidone k-30, pregelatinized starch, stearic acid, sucralose, talc, titanium dioxide

**Questions or comments?**  
call 1-877-770-3183 Mon-Fri 8:00 AM EST to 5:00 PM PST

This product is not manufactured or distributed by Kenvue Inc., owner of Extra Strength Tylenol®.

**Drug Facts (continued)**  
These could be signs of a serious condition:

- pregnant or breast-feeding, ask a health professional before use
- Keep out of reach of children
- Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away, (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed (see overdose warning)
- adults and children 12 years and over:
  - take 2 tablets every 6 hours while symptoms last
  - do not take more than 6 tablets in 24 hours, unless directed by a doctor
  - do not use for more than 10 days unless directed by a doctor
- children under 12 years:
  - ask a doctor

**Other information**

- store at 20°-25°C (68°-77°F). See USP Controlled Room Temperature.
- see bottom of the label for lot number and expiration date

**Inactive ingredients**  
FD&C red no. 40, FD&C yellow no. 6, lecithin, polyethylene glycol, polyvinyl alcohol, povidone k-30, pregelatinized starch, stearic acid, sucralose, talc, titanium dioxide

**Questions or comments?**  
call 1-877-770-3183 Mon-Fri 8:00 AM EST to 5:00 PM PST

This product is not manufactured or distributed by Kenvue Inc., owner of Extra Strength Tylenol®.

how2recycle.info  
PLASTIC BOTTLE

LOT  
200000005302  
700000003887

Exp.  
MADE IN INDIA

SEE HERE

www.meijer.com

FD-506450  
1928379235

simulate a printed label,  
fold along dotted line.

Inside (adhesive side)

**Drug Facts (continued)**  
These could be signs of a serious condition:

- pregnant or breast-feeding, ask a health professional before use
- Keep out of reach of children
- Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away, (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed (see overdose warning)
- adults and children 12 years and over:
  - take 2 tablets every 6 hours while symptoms last
  - do not take more than 6 tablets in 24 hours, unless directed by a doctor
  - do not use for more than 10 days unless directed by a doctor
- children under 12 years:
  - ask a doctor

**Other information**

- store at 20°-25°C (68°-77°F). See USP Controlled Room Temperature.
- see bottom of the label for lot number and expiration date

**Inactive ingredients**  
FD&C red no. 40, FD&C yellow no. 6, lecithin, polyethylene glycol, polyvinyl alcohol, povidone k-30, pregelatinized starch, stearic acid, sucralose, talc, titanium dioxide

**Questions or comments?**  
call 1-877-770-3183 Mon-Fri 8:00 AM EST to 5:00 PM PST

This product is not manufactured or distributed by Kenvue Inc., owner of Extra Strength Tylenol®.

## ACETAMINOPHEN

acetaminophen tablet

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:79481-5380
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
POVIDONE K30 (UNII: U725QWY32X)	
STARCH, CORN (UNII: O8232NY35J)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TALC (UNII: 7SEV7J4R1U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

**Product Characteristics**

<b>Color</b>	red	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	11mm
<b>Flavor</b>		<b>Imprint Code</b>	L;1
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79481-5380-2	225 in 1 BOTTLE; Type 0: Not a Combination Product	09/09/2024	

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M013	09/09/2024	

**Labeler** - Meijer (006959555)