

**CRATIV SUNSCREEN- ethylhexyl methoxycinnamate,titanium dioxide, homosalate, ethylhexyl, salicylate, octocrylene, bis-ethylhexyloxyphenolmethoxyphenyltriazine,4-methylbenzylidene camphor, dimethicone,butyl methoxydibenzoylmethane lotion
Foshan Xinhai Jiayu Import and Export Trade Co., Ltd**

CraTiv Sunscreen

Ethylhexyl methoxycinnamate 10%
Titanium dioxide 4.5%
Homosalate 6%
Ethylhexyl salicylate 4.5%
Octocrylene 2.9%
Bis-ethylhexyloxyphenolmethoxyphenyltriazine 2%
4-methylbenzylidene camphor 0.5%
DIMETHICONE 1%
Butyl methoxydibenzoylmethane 1%

Protecting the skin from the damage caused by ultraviolet (UV) radiation. UV rays can lead to skin sunburn, sunburn erythema, skin cancer, and other issues. By forming a protective layer on the skin surface, sunscreen blocks, absorbs, or reflects UV radiation, thus reducing its harmful effects on the skin.

Protecting the skin from the damage caused by ultraviolet (UV) radiation. UV rays can lead to skin sunburn, sunburn erythema, skin cancer, and other issues. By forming a protective layer on the skin surface, sunscreen blocks, absorbs, or reflects UV radiation, thus reducing its harmful effects on the skin.

For external use only

5. Do not use when the skin has abnormal conditions. This product has passed the quality test, but it does not mean that very few people will not have adverse reactions such as allergies. If anything abnormal occurs during use, discontinue use immediately and consult your doctor if necessary.

1. Do not place in sunlight and high temperatures, do not swallow. 2. Keep out of reach of the baby. 3. Open the bag and use it. 4. In case of contact with eyes, rinse immediately with water.

5. Do not use when the skin has abnormal conditions. This product has passed the quality test, but it does not mean that very few people will not have adverse reactions such as allergies. If anything abnormal occurs during use, discontinue use immediately and consult your doctor if necessary.

Keep out of reach of the baby.

Step 1: Cleanse. Start with clean and dry skin. Wash your face and any exposed areas gently to remove dirt and sweat.

Step 2: Apply Before Sun Exposure. Apply sunscreen generously at least 15 minutes before going out in the sun. This allows your skin to absorb the product properly.

Step 3: Amount Matters. Use enough sunscreen to cover all exposed skin. As a general guide, apply about a teaspoon (5ml) to your face and a shot glass worth for your body.

Step 4: Even Application. Spread the su-nscreen evenly over your skin. Don't for-get easily overlooked areas like ears, ne-ck, and the back of your hands.

Step 5: Reapply. Reapply sunscreen eve-ry two hours, or more often if you're sw-eating heavily or swimming. Even water-resistant' sunscreen can wear off.

1. Do not place in sunlight and high temperatures, do not swallow. 2. Keep out of reach of the baby.

water

propylene glycol

C12-15 Alcohol Benzoate

cyclopentasiloxane

Caprylic/capric triglyceride

PEG-30 Dipolyhydroxystearate

Sodium chloride

MONTMORILLONITE

Sorbitan oleate

Phenoxyethanol

.ALPHA.-TOCOPHEROL

SILICON DIOXIDE

Methylparaben

CETYL PHOSPHATE

Propylparaben

Ethylhexylglycerin

Sodium hyaluronate



CRATIV SUNSCREEN

ethylhexyl methoxycinnamate, titanium dioxide, homosalate, ethylhexyl, salicylate, octocrylene, bis-ethylhexyloxyphenolmethoxyphenyltriazine, 4-methylbenzylidene camphor, dimethicone, butyl methoxydibenzoylmethane lotion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:84301-001
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTOCRYLENE (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	2.9 g in 100 mL
BEMOTRIZINOL (UNII: PWZ1720CBH) (BEMOTRIZINOL - UNII:PWZ1720CBH)	BEMOTRIZINOL	2 g in 100 mL
AVOBENZONE (UNII: G63QQF2NOX) (AVOBENZONE - UNII:G63QQF2NOX)	AVOBENZONE	1 g in 100 mL
HOMOSALATE (UNII: V06SV4M95S) (HOMOSALATE - UNII:V06SV4M95S)	HOMOSALATE	6 g in 100 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	4.5 g in 100 mL
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	10 g in 100 mL
OCTISALATE (UNII: 4X49Y0596W) (OCTISALATE - UNII:4X49Y0596W)	OCTISALATE	4.5 g in 100 mL
ENZACAMENE (UNII: 8I3XWY40L9) (ENZACAMENE - UNII:8I3XWY40L9)	ENZACAMENE	0.5 g in 100 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	1 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
MONTMORILLONITE (UNII: A585MN1H2L)	
CETYL PHOSPHATE (UNII: VT07D6X67O)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
HYALURONATE SODIUM (UNII: YSE9PPT4TH)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
WATER (UNII: 059QF0KO0R)	
ALKYL (C12-15) BENZOATE (UNII: A9EJ3J61HQ)	
SORBITAN MONOOLEATE (UNII: 06XEA2VD56)	
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)	
ETHYLHEXYLGLYCERIN (UNII: 147D247K3P)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PEG-30 DIPOLYHYDROXYSTEARATE (UNII: 9713Q0S7FO)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84301-001-01	50 mL in 1 TUBE; Type 0: Not a Combination Product	05/11/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	05/11/2024	

Labeler - Foshan Xinhai Jiayu Import and Export Trade Co., Ltd (412478748)

Establishment

Name	Address	ID/FEI	Business Operations
Foshan Xinhai Jiayu Import and Export Trade Co., Ltd		412478748	manufacture(84301-001)

Revised: 5/2024

Foshan Xinhai Jiayu Import and Export Trade Co., Ltd