

ASPIRIN ENTERIC COATED LOW DOSE- aspirin tablet, delayed release
Walgreen Company

Walgreens 44-600A

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- take more or for a longer time than directed
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product
- take a blood thinning (anticoagulant) or steroid drug

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
 - feel faint
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- see end flap for expiration date and lot number
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients

corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-426-9391

Principal Display Panel

Walgreens

• **WALGREENS** •

PHARMACIST RECOMMENDED[†]

Compare to the active ingredient in
Bayer[®] Low Dose Aspirin^{††}

Aspirin 81

81 mg / PAIN RELIEVER (**NSAID**)

Low Dose Enteric-Coated

**HEART
HEALTH****

**200
TABLETS**

ACTUAL SIZE

****Talk to your doctor or other healthcare
provider before using this product for your heart.**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

NDC 0363-0563-06

[†]Our pharmacists recommend the Walgreens brand.
We invite you to compare to national brands.

^{††}This product is not manufactured or distributed by
Bayer AG, owner of the registered trademark
Bayer[®] Low Dose Aspirin.

50844 REV0122A60006

DISTRIBUTED BY: **WALGREEN CO.**

DEERFIELD, IL 60015
Walgreens
100% SATISFACTION
GUARANTEED
walgreens.com
 ©2024 Walgreen Co.

Do not print Lot & Exp Only

3 1917 18593 4

NDC 0363-063-06
 ITEM 945684 W00000-0000-0

TAMPER EVIDENT- DO NOT USE IF IMPAIRED
 SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Walgreens
 Compare to the active ingredient in
 Bayer® Low Dose Aspirin™

Aspirin 81
 81 mg / PAIN RELIEVER (NSAID)

Low Dose Enteric-Coated

200
 TABLETS

ACTUAL SIZE

*Talk to your doctor or other healthcare provider before using this product for your heart.

how2recycle.info

B-2201-600A-06-HR2
 REV10122A60006

HEART HEALTH

*Talk to your doctor or other healthcare provider before using this product for your heart.

Our pharmacists recommend the Walgreens brand.
 We invite you to compare to national brands.
 This product is not manufactured or distributed by
 Bayer AG, owner of the registered trademark
 Bayer® Low Dose Aspirin.

50844 REV10122A60006

DISTRIBUTED BY: WALGREEN CO.
 DEERFIELD, IL 60015

Walgreens
100% SATISFACTION
GUARANTEED

walgreens.com
 ©2024 Walgreen Co.

Walgreens 44-600A

ASPIRIN ENTERIC COATED LOW DOSE

W30RG1221-F
 REV-0924

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking a prescription drug for:

- gout
- diabetes
- arthritis

Stop use and ask a doctor if:

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
 - feel faint
- have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 3 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- see end flap for expiration date and lot number
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients corn starch, D&C yellow #10, FD&C yellow #6, hypromellose, methacrylic acid, microcrystalline cellulose, polyethylene glycol, shellac wax, silica, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments? 1-800-426-9391

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredient (in each tablet)
 Aspirin 81 mg (NSAID)¹ Pain reliever
¹nonsteroidal anti-inflammatory drug

Purpose
 Pain reliever

Uses For the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

Warnings
 Reye's syndrome: Child or teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product when using this product. If changes behavior with this product, stop use and ask a doctor. In rare cases, these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take more or for a longer time than directed
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product
- take a blood thinning (anticoagulant) or steroid drug

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

Walgreens
 Compare to the active ingredient in
 Bayer® Low Dose Aspirin™

Aspirin 81
 81 mg / PAIN RELIEVER (NSAID)

Low Dose Enteric-Coated

200
 TABLETS

ACTUAL SIZE

*Talk to your doctor or other healthcare provider before using this product for your heart.

Walgreens
 Compare to the active ingredient in
 Bayer® Low Dose Aspirin™

Aspirin 81
 81 mg / PAIN RELIEVER (NSAID)

Low Dose Enteric-Coated

200
 TABLETS

ACTUAL SIZE

*Talk to your doctor or other healthcare provider before using this product for your heart.

aspirin tablet, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0563
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 (UNII: 355W5USQ3G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
METHACRYLIC ACID (UNII: 1CS02G8656)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SHELLAC (UNII: 46N107B71O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	L
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0563-06	1 in 1 CARTON	05/01/2011	
1		200 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

NDC:0363

2	NDC:0363-0563-17	1 in 1 CARTON	05/01/2011	
2		300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:0363-0563-27	1 in 1 CARTON	05/01/2011	
3		32 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
4	NDC:0363-0563-14	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/01/2011	
5	NDC:0363-0563-32	1 in 1 CARTON	05/01/2011	
5		120 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
6	NDC:0363-0563-63	2 in 1 CARTON	05/01/2011	
6		120 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
7	NDC:0363-0563-62	1 in 1 CARTON	07/19/2023	02/28/2026
7		15 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
8	NDC:0363-0563-22	1 in 1 CARTON	05/01/2011	08/14/2021
8		48 in 1 BOTTLE; Type 0: Not a Combination Product		
9	NDC:0363-0563-90	1 in 1 CARTON	05/01/2011	03/30/2022
9		150 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/01/2011	

Labeler - Walgreen Company (008965063)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(0363-0563)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(0363-0563)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(0363-0563)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(0363-0563)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(0363-0563)

Revised: 1/2026

Walgreen Company