

PHYSICIANS CARE NON-DROWSY COLD AND COUGH- acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride tablet, film coated
Acme United

Physicians Care Non-Drowsy Cold & Cough

Drug Facts

Drug Facts

Active Ingredient

Active ingredient (in each caplet)

Acetaminophen 325 mg

Dextromethorphan Hydrobromide 10 mg

Guaifenesin 100 mg

Phenylephrine Hydrochloride 5 mg

Purpose

Purpose

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

Uses

Temporarily relieves these cold symptoms

- cough
- sore throat
- minor aches and pains
- headache
- nasal congestion
- sinus congestion and pressure
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make cough more productive

Temporarily reduces fever.

Warnings

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough that lasts as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

Stop use and ask a doctor if

- new symptoms occur
- redness or swelling is present
- pain, cough or nasal congestion gets worse or lasts for more than 7 days
- fever gets worse or lasts more than 3 days
- nervousness, dizziness, or sleeplessness occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Directions

- do not take more than directed

Adults and children: (12 years and older)

Take 2 caplets every 4 hours

Do not to take more than 10 caplets in 24 hours

Children under 12 years: ask a doctor

Other information

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- tamper evident sealed packets

- do not use any opened or torn packets

Inactive ingredients

Inactive ingredients

corn starch, crospovidone, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions

Questions? 1-800-835-2263

Physicians Care Cold & Cough

91462

PhysiciansCare®

Non-Drowsy

Cold & Cough

Cold, Cough and Flu Relief

Pull

to Open

COATED CAPLET

Individually packaged for workplace use

Tamper-Evident Packets of 2 Caplets

Do not use if packet is open or torn.

Not Actual Size

100 Caplets (50 Packets, 2 caplets each)

91462

Retain carton for complete product information

91462

91462

Also Available from PhysiciansCare® in individual dose packets:

- PhysiciansCare® Aspirin
- PhysiciansCare® Ibuprofen
- PhysiciansCare® Non-Aspirin (Acetaminophen)
- PhysiciansCare® Antacid
- PhysiciansCare® Sinus
- PhysiciansCare® Allergy

PhysiciansCARE

Non-Drowsy COLD & COUGH

Cold, Cough and Flu Relief

COATED CAPLET

Individually packaged for workplace use

Tamper-Evident Packets of 2 Caplets
Do not use if packet is open or torn.

Not Actual Size

100 Caplets (50 Packets, 2 caplets each)

Drug Facts

Active ingredient in each caplet: Acetaminophen 325 mg, Dextromethorphan Hydrobromide 10 mg, Phenylephrine Hydrochloride 5 mg.

Purpose
Temporarily relieves these cold symptoms: cough, sore throat, minor aches and pains, headache, nasal congestion, helps loosen phlegm (mucus) and thin bronchial secretions to make coughing more productive. Temporarily reduces fever.

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
• more than 4,000 mg of acetaminophen in 24 hours
• with other drugs containing acetaminophen
• or more alcoholic drinks every day while using this product.
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
• skin redness • hives • rash
If a skin reaction occurs, stop use and seek medical help right away.
See throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, tinnitus or vomiting, consult a doctor promptly.

Do not use
• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
• if you have ever had an allergic reaction to this product or any of its ingredients.
• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
• liver disease • heart disease • high blood pressure • kidney disease • diabetes • difficulty in urination due to enlargement of the prostate gland • cough that occurs with the mucus (phlegm) • persistent or chronic cough that lasts as occurs with smoking, asthma, chronic bronchitis or emphysema.

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are:
• taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if:
• new symptoms occur
• redness or swelling is present
• pain, cough or nasal congestion gets worse or lasts for more than 7 days
• fever gets worse or lasts more than 3 days
• nervousness, dizziness, or sleeplessness occur
• cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Directions
• Do not take more than directed.

Adults and children (12 years and older) Take 2 caplets every 4 hours. Do not take more than 10 caplets in 24 hours.

Children under 12 years: Ask a doctor.

Other information
• Store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
• tamper evident sealed packets
• Do not use any opened or torn packets.

Inactive ingredients
corn starch, crospovidone, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide.

Questions? 1-800-835-2263

Contact Us 1-800-835-2263 www.FirstAidOnly.com

100 Caplets (50 Packets, 2 caplets each)

PHYSICIANS CARE NON-DROWSY COLD AND COUGH

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride
tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0924-0142
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
CROSPVIDONE (UNII: 2S7830E561)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
POVIDONE (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
MAGNESIUM STEARATE (UNII: 70097M6130)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	orange	Score	no score
Shape	CAPSULE ((Caplet))	Size	17mm
Flavor		Imprint Code	44;546
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0924-0142-01	50 in 1 BOX	01/19/2023	

1	NDC:0924-0142-00	2 in 1 PACKET; Type 0: Not a Combination Product		
----------	------------------	--	--	--

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/19/2023	

Labeler - Acme United (001180207)

Registrant - Unifirst First Aid Corportaion (832947092)

Establishment

Name	Address	ID/FEI	Business Operations
Prestige Packaging		080667761	pack(0924-0142)

Establishment

Name	Address	ID/FEI	Business Operations
Medique Products		086911794	pack(0924-0142)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International		832867837	manufacture(0924-0142)

Revised: 9/2025

Acme United