

**AXIV NIGHT TIME - LIQUID- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate liquid**  
**VIVUNT PHARMA LLC**

-----  
**AXIV Night Time Cold & Flu - Liquid**

**Drug Facts**

<b>Active ingredients (in each 30 mL dose cup)</b>	<b>Purpose</b>
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 30 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine

**Uses**

Temporarily relieves common cold/flu symptoms:

- sore throat
- minor aches & pains
- runny nose & sneezing
- headache
- fever
- cough due to minor throat & bronchial irritation

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if:

- adults and children over 12 years of age take more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning**

If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- to make a child sleep

**Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

**When using this product**

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**Keep out of reach of children.****If pregnant or breast-feeding**

ask a health professional before use.

**Overdose warning**

Taking more than the recommended dose (overdose) could cause serious health problems, including liver damage. In case of overdose, get medical help or contact a

Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms

### **Directions**

- take only as directed (see overdose warning)
- measure only with dosage cup provided and keep dosage cup with product
- mL = milliliter
- do not exceed 4 doses per 24 hours
- when using DayTime and NightTime products, limit total to 4 doses per 24 hours

---

adults and children 12 years and over  
children under 12 years

30 mL every 6 hours  
do not use

---

### **Other information**

- Each 30 mL dose cup contains: sodium 10 mg
- Store at room temperature 15 - 30°C (59 - 89°F) and do not refrigerate
- Tamper evident: Do not use if shrink band is missing or broken

### **Inactive ingredients**

Citric acid, FD&C Blue #1, FD&C Red #40, flavor, glycerin, menthol, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum

### **Distributed by:**

**VIVUNT PHARMA LLC**  
**8950 SW 74th. Court. Suite 1901**

**Miami, FL 33156-3178**

**Made in USA**

**[www.vivunt.live](http://www.vivunt.live)**

### **PRINCIPAL DISPLAY PANEL**

Compare to VICKS<sup>®</sup> NyQuil<sup>®</sup>

Cold & Flu active ingredients\*

NDC 82706-023-01

AXIV - NightTime

COLD&FLU

Multi-Symptom Relief

- Pain Reliever
- Fever Reducer
- Cough Suppresant

- Antihistamine

## Nighttime Relief

Acetaminophen, Dextromethorphan HBr, Doxylamine Succinate

12 FL OZ (355 mL)

\*This product is not manufactured or distributed by The Procter & Gamble Company, owner of registered trademarks Vicks® NyQuil®

**Compare to Vicks® NyQuil® Cold & Flu active ingredients\* NDC 82706-023-01**

**axiv**

**NightTime**

**COLD & FLU**

**Multi-Symptom Relief**

- Pain Reliever
- Fever Reducer
- Cough Suppressant
- Antihistamine

**NIGHTTIME RELIEF**

Acetaminophen HBr  
Doxylamine Succinate

CHERRY FLAVOR  
ALCOHOL FREE

**12 FL OZ (355 mL)**

**Drug Facts**

**Active ingredients (in each 30 mL dose cup)**

Acetaminophen 650 mg	Pain Reliever-Fever Reducer
Dextromethorphan HBr 30 mg	Cough Suppressant
Doxylamine Succinate 12.5 mg	Antihistamine

**Purpose**

Temporarily relieves common cold/flu symptoms: ■ sore throat ■ minor aches and pains ■ runny nose and sneezing ■ headache ■ fever ■ cough due to minor throat and bronchial irritation.

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if: ■ adults and children over 12 years of age take more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ to make a child sleepy.

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts or as occur with smoking, asthma, chronic bronchitis or emphysema ■ trouble urinating due to an enlarged prostate gland.

**Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers** ■ taking the blood thinning drug warfarin.

**When using this product** ■ do not use more than directed ■ avoid alcoholic drinks ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness.

**Stop use and ask a doctor if** ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new

**Drug Facts (continued)**

symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**Keep out of reach of children. If pregnant or breast-feeding, ask a health professional before use.**

**Overdose warning:** Taking more than the recommended dose (overdose) could cause serious health problems, including liver damage. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ take only as directed (see overdose warning) ■ measure only with dosage cup provided and keep dosage cup with product ■ mL= milliliter ■ do not exceed 4 doses per 24 hours

adults and children 12 years and over	30 mL every 6 hours
children under 12 years	do not use

■ when using DayTime and NightTime products, limit total to 4 doses per 24 hours.

**Other information** ■ Each 30 mL dose cup contains: sodium 10 mg ■ store at room temperature 15 - 30°C (59-86 °F) and do not refrigerate ■ Tamper-evident: Do not use if shrink band is missing or broken.

**Inactive ingredients** Citric acid, FD&C Blue #1, FD&C Red #40, flavor, glycerin, menthol, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum.

**VIVUNT™ 30 YEARS**

Distributed by:  
VIVUNT PHARMA LLC  
8850 SW 74th Court, Suite 1901  
Miami, FL 33156-3178  
Made in USA

www.vivunt.live

8 50047 1178064 5

This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® NyQuil® Cold & Flu.

## AXIV NIGHT TIME - LIQUID

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:82706-023
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RT919KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	

<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>MENTHOL, UNSPECIFIED FORM</b> (UNII: L7T10EIP3A)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SORBITOL SOLUTION</b> (UNII: 8KW3E207O2)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

### Product Characteristics

<b>Color</b>	red	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82706-023-01	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/14/2024	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/14/2024	

**Labeler** - VIVUNT PHARMA LLC (045829437)

Revised: 9/2025

VIVUNT PHARMA LLC