

AQUAPRIME LIQUID SODIUM SALICYLATE- sodium salicylate liquid Blendpack, Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

AquaPrime Liquid Sodium Salicylate

Indications

For use in the drinking water of poultry and swine as an aid in reducing pain, fever and inflammation

CALF LABEL CLAIM

Indications For Use, Calves: Supportive treatment of pyrexia in acute respiratory disease, in combination with appropriate (e.g., anti-infective) therapy if necessary.

Contraindications, Calves: Do not use Sodium Salicylate in neonates or calves less than 2 weeks of age

Dosage, Calves: 40 mg Sodium Salicylate per kg body weight once daily, for 1-3 days. Administer orally in drinking water or milk (replacer).

Recommended withdrawal period for all food animals following use at "usual dosages": meat 24 hours.

Warnings

Not for use in lactating dairy cattle.

A withdrawal period of 24 hours is recommended in meat.

Not for human use.

Keep container tightly closed.

Keep out of reach of children.

Store at room temperature.

Warning: Do not administer concentrated solution by direct oral administration — gastrointestinal irritation or overdose may occur. Do not use in piglets less than 4 weeks of age. Store at controlled room temperature between 15°-30° C (59°-86° F).

Directions for use

For Analgesic and Antipyretic Use

Water Proportioner Use: Add 8 ounces (236 ml) of Sodium Salicylate 48.6% Concentrate to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 11.3 mg/b (25 mg/kg) body weight daily.

Livestock Tank Use: 8 ounces (473 ml) of Sodium Salicylate 48.6% Concentrate to 128 gallons of drinking water. This will achieve a target dose of 11.3 mg/b (25 mg/kg) body weight daily.

For Anti-Inflammatory/Anti-Prostaglandin Use

Water Proportioner Use: Add 16 ounces (473 ml) of Sodium Salicylate 48.6% Concentrate to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 22.7 mg/lb (50 mg/kg) body weight daily.

Livestock Tank Use: Add 16 ounces (473 ml) of Sodium Salicylate 48.6% Concentrate to 128 gallons of drinking water. This will achieve a target dose of 22.7 mg/lb (50 mg/kg) body weight daily.

Following Days

Water Proportioner Use: Add 10 ounces (296 ml) of Sodium Salicylate 48.6% Concentrate to make 1 gallon of stock solution and administer through a mediator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 13.6 mg/b (30 mg/kg) body weight daily.

Livestock Tank Use: Add 10 ounces (296 ml) of Sodium Salicylate 48.6% Concentrate to 128 gallons of drinking water. This will achieve a target dose of 13.6 mg/lb (30 mg/kg) body weight daily.

Prepare fresh solutions daily. Repeat as necessary.

Liquid Sodium Salicylate for Drinking Water Solutions

AquaPrime

Liquid Sodium Salicylate

48.6%

ANALGESIC, ANTIPYRETIC, AND ANTI-INFLAMMATORY

NEOGEN

Item Number: 486256/1

Net Contents: 1 gal. (3.79L)

7 89703 00612 9

INDICATIONS: For use in the drinking water of poultry and swine as an aid in reducing pain, fever and inflammation.

CALF LABEL CLAIM
Indications For Use, Calfes: Supportive treatment of pyrexia in acute respiratory disease, in combination with appropriate (e.g., antimicrobial) therapy if necessary.
Contraindications, Calfes: Do not use Sodium Salicylate in neonates or calves less than 2 weeks of age.
Dosage, Calfes: 40 mg Sodium Salicylate per kg body weight once daily for 1-3 days. Administer orally in drinking water or milk replacer.
Resume normal withdrawal period for all food animals following use at "usual dosages"; max. 24 hours.

ACTIVE INGREDIENT:
Sodium Salicylate 48.6% w/v

DIRECTIONS FOR USE:
For Analgesic and Antipyretic Use
Water Proportioner Use: Add 8 ounces (226 ml) of Sodium Salicylate 48.6% Concentrate to make 1 gallon of stock solution and administer through a mediator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 11.3 mg/b (25 mg/kg) body weight daily.
Livestock Tank Use: 8 ounces (473 ml) of Sodium Salicylate 48.6% Concentrate to 128 gallons of drinking water. This will achieve a target dose of 11.3 mg/b (25 mg/kg) body weight daily.
For Anti-Inflammatory/Anti-Prostaglandin Use
Water Proportioner Use: Add 16 ounces (473 ml) of Sodium Salicylate 48.6% Concentrate to make 1 gallon of stock solution and administer through a mediator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 22.7 mg/lb (50 mg/kg) body weight daily.
Livestock Tank Use: Add 16 ounces (473 ml) of Sodium Salicylate 48.6% Concentrate to 128 gallons of drinking water. This will achieve a target dose of 22.7 mg/lb (50 mg/kg) body weight daily.

Following Days
Water Proportioner Use: Add 10 ounces (296 ml) of Sodium Salicylate 48.6% Concentrate to make 1 gallon of stock solution and administer through a mediator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 13.6 mg/b (30 mg/kg) body weight daily.
Livestock Tank Use: Add 10 ounces (296 ml) of Sodium Salicylate 48.6% Concentrate to 128 gallons of drinking water. This will achieve a target dose of 13.6 mg/lb (30 mg/kg) body weight daily.
Prepare fresh solutions daily. Repeat as necessary.

WARNING: Do not administer concentrated solutions by direct oral administration - gastro-intestinal irritation or overdose may occur. Do not use in poultry less than 4 weeks of age. Store as concentrated in a temperature between 15-30°C (59-86°F).

CAUTION: Keep container closed when not in use. Pail at may solidify at cold storage temperatures. Place container in room temperature storage, which will thaw the solution or place in warm water. Gently invert container to ensure uniformity of product.

Good animal care will not affect product stability.

Manufactured by: **Blue Ridge Elements**
3730 Old Tasso Rd.
Cleveland, TN, 37312
423-863-0808

Add image transcription here...

AQUAPRIME LIQUID SODIUM SALICYLATE
sodium salicylate liquid

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:86050-2486
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM SALICYLATE (UNII: WQ1H85SYP) (SALICYLIC ACID - UNII:O414PZ4LPZ)	SODIUM SALICYLATE	486 g in 1 L

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:86050-2486-1	3.79 L in 1 JUG		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		07/27/2023	

Labeler - Blendpack, Inc. (043362034)

Establishment

Name	Address	ID/FEI	Business Operations
Blendpack, Inc.		043362034	manufacture

Revised: 7/2023

Blendpack, Inc.