

**CHILDRENS PAIN RELIEVER AND FEVER REDUCER CHERRY-
acetaminophen suspension
DeMoulas Market Basket**

Children's Pain Relief Oral Suspension, Cherry

Drug Facts

Active ingredient

(in each 5 mL = 1 teaspoonful)

Acetaminophen 160 mg Pain reliever/fever reducer

Purpose

Acetaminophen Pain reliever/fever reducer

Uses temporarily • reduces fever

- relieves minor aches and pains due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Warnings

Liver warning

This product contains acetaminophen.

Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Keep out of reach of children.

Directions

- **this product does not contain directions or complete warnings for adult use.**
- **shake well before using**
- mL = milliliter; tsp = teaspoonful
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor

Weight (lb)	Age (yr)	Dose (mL or tsp) *
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL (1 tsp)
36-47	4-5 years	7.5 mL (1½ tsp)
48-59	6-8 years	10 mL (2 tsp)
60-71	9-10 years	12.5 mL (2½ tsp)
72-95	11 years	15 mL (3 tsp)

*or as directed by a doctor

Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL (1 tsp) contains: sodium 3 mg
- store between 20-25°C (68-77°F)
- do not refrigerate
- **Keep carton for full directions for use.**

Inactive ingredients

anhydrous citric acid, butylparaben, FD&C red #40, flavor, glycerin, high fructose corn syrup, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

Children's Pain Relief Oral Suspension, Cherry - PRINCIPAL DISPLAY PANEL

MARKET BASKET®

"MORE FOR YOUR DOLLAR"

Children's

Compare

to the Active

Pain &

Ingredients in Children's

Fever

Tylenol® Oral Suspension

Acetaminophen

Oral Suspension

Cherry Flavor

4 FL OZ (118 mL)

**DISTRIBUTED BY
DEMOULAS SUPERMARKETS INC.
875 East Street
Tewksbury, MA 01876**

DO NOT USE IF PRINTED NECK BAND IS BROKEN OR MISSING

This product is not manufactured or distributed by McNeil Consumer Healthcare,
distributor of Children's Tylenol®

Oral Suspension.

Carton

MARKET BASKET (100% WHOLESALE)
BX-176H

GU_3353 K 254769

DO NOT USE IF PRINTED NECK BAND IS
BROKEN OR MISSING

Drug Facts
(in each 5 mL = 1 teaspoonful)
Active Ingredient Acetaminophen 160 mg **Purpose** Pain reliever/fever reducer

Uses Temporarily reduces fever
relieves minor aches and pains due to:
the common cold flu headache sore throat
toothache

Warnings
Liver warning
This product contains acetaminophen. Severe liver damage may occur if your child takes:
more than 5 doses in 24 hours, which is the maximum daily amount
with other drugs containing acetaminophen
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
skin reddening blisters rash
If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning
If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use
with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has liver disease.
Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product do not exceed recommended dose (see overdose warning).

Stop use and ask a doctor if
pain gets worse or lasts more than 5 days
fever gets worse or lasts more than 3 days
new symptoms occur
redness or swelling is present
These could be signs of a serious condition.

Keep out of reach of children.
Overdose warning
Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is

Drug Facts (continued)

critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use.
- shake well before using
- mL = milliliter; tsp = teaspoonful
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor.

Weight (lb)	Age (yr)	Dose (mL or tsp)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL (1 tsp)
36-47	4-5 years	7.5 mL (1½ tsp)
48-59	6-8 years	10 mL (2 tsp)
60-71	9-10 years	12.5 mL (2½ tsp)
72-85	11 years	15 mL (3 tsp)

*or as directed by a doctor
Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information

- each 5 mL (1 tsp) contains: sodium 3 mg
- store between 20°-25°C (68°-77°F)
- do not refrigerate
- Keep cartons for full directions for use.

Inactive ingredients anhydrous citric acid, butylparaben, FD&C red #40, flavor, glycerin, high fructose corn syrup, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Oral Suspension.

DISTRIBUTED BY: DEMOULAS SUPERMARKETS INC.
875 EAST STREET, TEWKSBURY, MA 01876

@marketbasket
Please visit us at:
shopmarketbasket.com



MARKET BASKET
MADE FOR YOUR DOLLAR

Children's Pain & Fever

Acetaminophen Oral Suspension

Compare to the Active Ingredient in CHILDREN'S TYLENOL® Oral Suspension*

Children's Pain & Fever

Acetaminophen Oral Suspension

PAIN RELIEVER FEVER REDUCER

- Alcohol-Free
- Ibuprofen Free
- Aspirin Free
- For Ages 2 to 11

Cherry Flavor

4 FL OZ (118 mL)
160 mg per 5 mL

MARKET BASKET
MADE FOR YOUR DOLLAR

Children's Pain & Fever

Acetaminophen Oral Suspension

Compare to the Active Ingredient in CHILDREN'S TYLENOL® Oral Suspension*

Children's Pain & Fever

Acetaminophen Oral Suspension

PAIN RELIEVER FEVER REDUCER

- Alcohol-Free
- Ibuprofen Free
- Aspirin Free
- For Ages 2 to 11

Cherry Flavor

4 FL OZ (118 mL)
160 mg per 5 mL



Bottle

MARKET BASKET
MADE FOR YOUR DOLLAR

Children's Pain & Fever

Acetaminophen Oral Suspension

PAIN RELIEVER / FEVER REDUCER

- Alcohol-Free • Ibuprofen Free
- Aspirin Free • For Ages 2 to 11

Cherry Flavor

4 FL OZ (118 mL) 160 mg per 5 mL

DO NOT USE IF PRINTED NECK BAND IS BROKEN OR MISSING

Active Ingredient (in each 5 mL = 1 teaspoonful) Acetaminophen 160 mg **Purpose** Pain reliever/fever reducer

Uses Temporarily reduces fever relieves minor aches and pains due to:
the common cold flu headache sore throat toothache

Warnings: Liver warning This product contains acetaminophen. Severe liver damage may occur if your child takes:
more than 5 doses in 24 hours, which is the maximum daily amount
with other drugs containing acetaminophen
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
skin reddening blisters rash
If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

Do not use: with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
if your child is allergic to acetaminophen or any of the inactive ingredients in this product.
Ask a doctor before use if your child has liver disease.
Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product do not exceed recommended dose (see overdose warning).

Stop use and ask a doctor if:
pain gets worse or lasts more than 5 days
fever gets worse or lasts more than 3 days
new symptoms occur
redness or swelling is present
These could be signs of a serious condition.

Keep out of reach of children.
Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is

Directions

- this product does not contain directions or complete warnings for adult use.
- shake well before using
- mL = milliliter; tsp = teaspoonful
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor.

Weight (lb)	Age (yr)	Dose (mL or tsp)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL (1 tsp)
36-47	4-5 years	7.5 mL (1½ tsp)
48-59	6-8 years	10 mL (2 tsp)
60-71	9-10 years	12.5 mL (2½ tsp)
72-85	11 years	15 mL (3 tsp)

*or as directed by a doctor
Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Other information
each 5 mL (1 tsp) contains: sodium 3 mg
store between 20°-25°C (68°-77°F)
do not refrigerate
Keep cartons for full directions for use.

Inactive ingredients anhydrous citric acid, butylparaben, FD&C red #40, flavor, glycerin, high fructose corn syrup, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Tylenol® Oral Suspension.

DISTRIBUTED BY: DEMOULAS SUPERMARKETS INC.
875 EAST STREET, TEWKSBURY, MA 01876

@marketbasket
Please visit us at:
shopmarketbasket.com

for as directed by a doctor
Attention: use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device. **Other information** each 5 mL (1 tsp) contains: sodium 3 mg store between 20°-25°C (68°-77°F) do not refrigerate

Active ingredients: anhydrous citric acid, butylparaben, FD&C red #40, flavor, glycerin, high fructose corn syrup, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum

Questions? Call 1-877-798-5944

LOT:
DISTRIBUTED BY: DEMOULAS SUPERMARKETS INC.
875 EAST STREET, TEWKSBURY, MA 01876

LR-200
EXP:

CHILDRENS PAIN RELIEVER AND FEVER REDUCER CHERRY

acetaminophen suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53942-525
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
BUTYLPARABEN (UNII: 3QP11U3FV8)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	red	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53942-525-24	1 in 1 CARTON	01/01/2026	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/01/2026	

Labeler - DeMoulas Market Basket (007869647)

Registrant - Aptapharma Inc. (790523323)

Establishment

Name	Address	ID/FEI	Business Operations
Aptapharma Inc.		790523323	manufacture(53942-525)

Revised: 5/2026

DeMoulas Market Basket