

EXTRA STRENGTH ASPIRIN- aspirin tablet
Health Pharma USA LLC

Extra Strength Aspirin

Drug facts

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/ Fever reducer

Uses

Temporarily relieves minor aches and pains due to:

- headache
- muscular aches
- backache
- minor pain of arthritis
- the common cold
- toothache
- premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- shock
- facial swelling
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin, or any other pain reliever/fever reducer

Ask a doctor before use if

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- under a doctor's care for any serious condition
- taking any other drug

Stop use and ask a doctor if:

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- an allergic reaction occurs.

Seek medical help right away.

- ringing in the ears or loss of hearing occurs.
- pain gets worse or lasts for more than 10 days.
- fever gets worse or lasts for more than 3 days.
- redness or swelling present in the painful area.
- new symptoms occur.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- drink a full glass of water with each dose
- **adults and children 12 years of age and over:**
 - take 1 to 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
- **children under 12 years:**ask a doctor

Other information

- store at room temperature 15-30°C (59-86°F)
- read all product information before using
- **TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

Inactive ingredients

Starch, Povidone, Hydroxy propyl methyl cellulose, Polyethylene glycol, Talc, Titanium dioxide.

Questions or comments?

Call toll free 1-844-832-1138

Mon - Fri 9AM - 4PM EST

PRINCIPAL DISPLAY PANEL

Extra Strength Aspirin 325 mg - 100ct, 300ct, 500ct

Compare to Bayer® EXTRA STRENGTH active ingredients*

This product is not manufactured or distributed by Bayer HealthCare LLC., owner of the registered trademark Bayer® Low Dose Aspirin.



Drug Facts

Ask a doctor if you have: stomach pain, you have a fever, or kidney disease, or you are taking any other medicine.

Ask a doctor if you are taking any other medicine, or if you are taking any other medicine, or if you are taking any other medicine.

Stop use and ask a doctor if you experience: bleeding, blood, or hives, or if you experience any other allergic reaction.

Seek medical help if you experience: ringing in ears, pain, fever, or redness, or if you experience any other symptoms.

If pregnant or breastfeeding, ask a health professional before use. Do not use as directed if you are pregnant or breastfeeding.

Directions: Drink a full glass of water with each dose. Adults and children 12 years and older: take 1 to 2 tablets every 4 to 6 hours, not to exceed 12 tablets in 24 hours. Children 6 to 11 years: take 1 to 2 tablets every 4 to 6 hours, not to exceed 12 tablets in 24 hours. Children under 6 years: do not use unless directed by a doctor.

Other information: Store at room temperature (20°C to 25°C). Excipients permitted between 15-30°C (59-86°F). **TAMPER EVIDENT:** DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

Inactive ingredients: starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide.

NDC 71679-115-01



Compare to Bayer® EXTRA STRENGTH active ingredients*

Extra Strength Aspirin

Aspirin 325 mg

Pain Reliever / Fever Reducer (NSAID)
Fast, Safe Pain Relief

100 COATED TABLETS

Drug Facts

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)* Pain reliever/Fever reducer

Nonsteroidal anti-inflammatory drug

Uses: temporarily relieves minor aches and pains due to: headache, muscular aches, backache, minor pain of arthritis, the common cold, toothache, premenstrual and menstrual cramps.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, it changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include: hives, shock, facial swelling, asthma (wheezing). Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed.

Do not use if you are allergic to aspirin, or any other pain reliever/fever reducer.

Directions: Drink a full glass of water with each dose. Adults and children 12 years and older: take 1 to 2 tablets every 4 to 6 hours, not to exceed 12 tablets in 24 hours. Children 6 to 11 years: take 1 to 2 tablets every 4 to 6 hours, not to exceed 12 tablets in 24 hours. Children under 6 years: do not use unless directed by a doctor.

Other information: Store at room temperature 25°C (77°F). Excursions permitted between 15-30°C (59-86°F). **TAMPER EVIDENT:** DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

Inactive ingredients: starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide.

Questions or comments? Call toll free 1-844-832-1138 Monday through Friday 9AM-5PM EST

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Bayer® Low Dose Aspirin.

Mfg./Dist. by: Health Pharma, Rahway, NJ 07065, www.health-pharma.us

Lot No.: 71679 11501

Exp. Date:

PEEL HERE



MADE IN THE USA

NDC 71679-115-03



Compare to Bayer® EXTRA STRENGTH active ingredients*

Extra Strength Aspirin

Aspirin 325 mg

Pain Reliever / Fever Reducer (NSAID)
Fast, Safe Pain Relief

300 COATED TABLETS

Drug Facts

Active ingredient (in each caplet)
Aspirin 325 mg (NSAID)* Pain reliever/Fever reducer

Nonsteroidal anti-inflammatory drug

Uses: temporarily relieves minor aches and pains due to: headache, muscular aches, backache, minor pain of arthritis, the common cold, toothache, premenstrual and menstrual cramps.

Warnings: Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if you experience behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: hives, facial swelling, asthma (wheezing), shock. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach ulcers or bleeding problems. The chance is higher if you: are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinning (anticoagulant) or steroid drug, take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed.

Do not use if you are allergic to aspirin, or any other pain reliever/fever reducer. **Ask a doctor before use** if you have a history of stomach problems, such as heartburn, or you have high blood pressure, heart disease, liver cirrhosis or kidney disease. **Do not use** if you are taking a diuretic.

Ask a doctor or pharmacist before use if you are taking a prescription drug for diabetes, gout, or arthritis. **Ask a doctor** if you are under a doctor's care for any serious condition.

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint, vomit blood, or experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better, pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, redness or swelling is present, ringing in the ears or a loss of hearing occurs.

If pregnant or breastfeeding, ask a health professional before use. **It is especially important** not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Do not use** if you are pregnant or breastfeeding, or if you are taking a diuretic.

Directions: Drink a full glass of water with each dose. Adults and children 12 years and older: take 1 to 2 tablets every 4 to 6 hours, not to exceed 12 tablets in 24 hours. Children 6 to 11 years: take 1 to 2 tablets every 4 to 6 hours, not to exceed 12 tablets in 24 hours. Children under 6 years: do not use unless directed by a doctor.

Other information: Store at room temperature 25°C (77°F). Excursions permitted between 15-30°C (59-86°F). **TAMPER EVIDENT:** DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

Inactive ingredients: starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide.

Questions or comments? Call toll free 1-844-832-1138 Monday through Friday 9AM-5PM EST

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Bayer® Extra Strength.

Mfg./Dist. by: Health Pharma, Rahway, NJ 07065, www.health-pharma.us

Lot No.: 71679 11503

Exp. Date:



MADE IN THE USA

NDC 71679-115-05



Compare to Bayer® EXTRA STRENGTH active ingredients*

Extra Strength Aspirin

Aspirin 325 mg

Pain Reliever / Fever Reducer (NSAID)
Fast, Safe Pain Relief

500 TABLETS

Drug Facts

Active ingredient (in each caplet)
Aspirin 325 mg (NSAID)* Pain reliever/Fever reducer

Nonsteroidal anti-inflammatory drug

Uses: temporarily relieves minor aches and pains due to: headache, muscular aches, backache, minor pain of arthritis, the common cold, toothache, premenstrual and menstrual cramps.

Warnings: Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if you experience behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: hives, facial swelling, asthma (wheezing), shock. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach ulcers or bleeding problems. The chance is higher if you: are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinning (anticoagulant) or steroid drug, take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed.

Do not use if you are allergic to aspirin, or any other pain reliever/fever reducer. **Ask a doctor before use** if you have a history of stomach problems, such as heartburn, or you have high blood pressure, heart disease, liver cirrhosis or kidney disease. **Do not use** if you are taking a diuretic.

Ask a doctor or pharmacist before use if you are taking a prescription drug for diabetes, gout, or arthritis. **Ask a doctor** if you are under a doctor's care for any serious condition.

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint, vomit blood, or experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better, pain gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, redness or swelling is present, ringing in the ears or a loss of hearing occurs.

If pregnant or breastfeeding, ask a health professional before use. **It is especially important** not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Do not use** if you are pregnant or breastfeeding, or if you are taking a diuretic.

Directions: Drink a full glass of water with each dose. Adults and children 12 years and older: take 1 to 2 tablets every 4 to 6 hours, not to exceed 12 tablets in 24 hours. Children 6 to 11 years: take 1 to 2 tablets every 4 to 6 hours, not to exceed 12 tablets in 24 hours. Children under 6 years: do not use unless directed by a doctor.

Other information: Store at room temperature 25°C (77°F). Excursions permitted between 15-30°C (59-86°F). **TAMPER EVIDENT:** DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

Inactive ingredients: corn starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide.

Questions or comments? Call toll free 1-844-832-1138 Monday through Friday 9AM-5PM EST

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Bayer® Extra Strength.

Mfg./Dist. by: Health Pharma, Rahway, NJ 07065, www.healthpharma.us

Lot No.: 71679 11505

Exp. Date:



MADE IN THE USA

EXTRA STRENGTH ASPIRIN

aspirin tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:71679-115
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE (UNII: FZ989GH94E)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	ASP325
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71679-115-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	10/15/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	10/15/2023	

Labeler - Health Pharma USA LLC (080804485)

Establishment

Name	Address	ID/FEI	Business Operations
Health Pharma USA LLC		080804485	manufacture(71679-115)

Revised: 5/2024

Health Pharma USA LLC