

**CORID TYPE A MEDICATED ARTICLE- amprolium granule
Huvepharma, Inc**

CORID® 25%

(amprolium)

Type A Medicated Article

Coccidiostat

**To be used in the preparation of medicated feed to aid
in the prevention and treatment of coccidiosis in calves.**

ACTIVE DRUG INGREDIENT: Amprolium 25%

INGREDIENTS: Corn Gluten Feed and Soybean Oil

SEE DIRECTIONS ON BACK PANEL

STORAGE: Store at or below 77°F (25°C), excursions permitted to 104°F (40°C).
Avoid prolonged exposure to high humidity.

Approved by FDA under NADA # 012-350

®CORID is a trademark of Huvepharma EOOD.

Net Wt.: 50 lb (22.68 kg)

CORID® 25%

(amprolium)

Type A Medicated Article

INDICATIONS FOR USE:

An aid in the prevention and treatment of coccidiosis caused by *Eimeria bovis* and *E. zurnii* in calves.

WITHDRAWAL PERIODS AND RESIDUE WARNINGS:

**Withdraw 24 hours before slaughter. A withdrawal period has not been
established for this
product in pre-ruminating calves. Do not use in calves to be processed for
veal.**

USER SAFETY WARNING:

Not for human use. Keep out of reach of children.

Contact Huvepharma Inc. at 1-877-994-4883 or <http://www.huvepharma.us>. For
additional information about
reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or
<http://www.fda.gov/reportanimalae>.

CAUTION: For satisfactory diagnosis, a microscopic examination of the feces should be
done by a veterinarian or
diagnostic laboratory before treatment. When treating outbreaks, drug should be
administered promptly after
diagnosis is determined.

**Do not use in feeds containing bentonite.
Use as the sole source of amprolium.**

Restricted Drug (California) - Use only as directed.

MIXING DIRECTIONS: CORID 25% (amprolium) Type A medicated article should be thoroughly and evenly mixed in the feed. CORID 25% may be used to manufacture a Type B medicated feed with a concentration of 2.5% amprolium or a Type C medicated feed in the concentration range of 0.0125% to 1.25% amprolium.

- Use 200 pounds of CORID 25% per ton of feed to produce Type B medicated feed containing 2.5% amprolium.
- Use 1 pound of CORID 25% per ton of feed to produce Type C medicated feed containing 0.0125% amprolium.*
- Use 100 pounds of CORID 25% per ton of feed to produce Type C medicated feed containing 1.25% amprolium.
- Use 10 pounds of 2.5% Type B per ton of feed to produce Type C medicated feed containing 0.0125% amprolium.
- Use 1000 pounds of 2.5% Type B per ton of feed to produce Type C medicated feed containing 1.25% amprolium.

AMOUNT AND DIRECTIONS FOR USE:

Prevention: 227 mg amprolium /100 lb (5 mg/kg) body weight per day for 21 days during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard.

Treatment: 454 mg amprolium/100 lb (10 mg/kg) bodyweight per day for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed and all calves in the group should be treated.

*To aid in even distribution of CORID 25% in the finished feed, prepare a mixture of CORID 25% in a portion of complete feed ingredient before mixing into the finished ration. Blend the mixture with the remainder of the finished feed and mix thoroughly.

The proper amount of amprolium supplement is mixed with the amount of ration consumed in one day. The tables below give suggested examples for using Type C medicated feed.

21 DAY PREVENTIVE PROGRAM:

DMI (as % Body Weight)		4	3	2
Body Weight (lb)	Amprolium per Day (mg)	0.0125% Amprolium	0.0167% Amprolium	0.025% Amprolium
		Amount of Feed per Day in Pounds		
100	227	4	3	2
200	454	8	6	4
300	681	12	9	6
400	908	16	12	8
500	1134	20	15	10
600	1361	24	18	12

5 DAY TREATMENT PROGRAM:

DMI (as % Body Weight)		4	3	2
Body Weight (lb)	Amprolium per Day (mg)	0.025% Amprolium	0.033% Amprolium	0.05% Amprolium
		Amount of Feed per Day in Pounds		
100	454	4	3	2
200	908	8	6	4
300	1362	12	9	6
400	1816	16	12	8
500	2268	20	15	10
600	2722	24	18	12



Distributed by:
 Huvepharma, Inc.
 Peachtree City, GA 30269

CORID[®] 25%
 (amprolium)
 Type A medicated article

INDICATIONS FOR USE:
 For the prevention and treatment of coccidiosis caused by *Coccidia* feed and *E. coli* in calves.

USE DIRECTIONS:
 For calves, use 25 mg of amprolium per pound of body weight per day for 21 days depending on the severity of the infection. For additional information about reporting side effects to animal drugs, contact FDA at 1-888-564-6362 or FDA.gov/medwatch.

CAUTION: For coccidiosis diagnosis, a microscopic examination of the feces should be done by a veterinarian or diagnostic laboratory before treatment. When testing outcome, drug should be discontinued promptly after diagnosis is determined.

Do not use in feeds containing tetracyclines.
 Use in the same ration as a tetracycline.

Medicated feed (Coccidiostat) - Use only as directed.

MIXING DIRECTIONS: CORID 25% (amprolium) Type A medicated article should be thoroughly and evenly mixed in the feed. CORID 25% may be used to medicate a Type B medicated feed with a concentration of 2.5% amprolium or a Type C medicated feed with a concentration of 0.025% to 1.25% amprolium.

- Use 700 pounds of CORID 25% per ton of feed to produce Type B medicated feed containing 2.5% amprolium.
- Use 1 pound of CORID 25% per ton of feed to produce Type C medicated feed containing 0.025% amprolium.
- Use 100 pounds of CORID 25% per ton of feed to produce Type C medicated feed containing 1.25% amprolium.
- Use 10 pounds of 2.5% Type B per ton of feed to produce Type C medicated feed containing 0.025% amprolium.
- Use 1000 pounds of 2.5% Type B per ton of feed to produce Type C medicated feed containing 1.25% amprolium.

ADMINISTRATIVE INFORMATION:
 Formulation: 222 mg amprolium (100%) (5 mg/kg) body weight per day for 21 days depending on the severity of the infection. For additional information about reporting side effects to animal drugs, contact FDA at 1-888-564-6362 or FDA.gov/medwatch.

Directions:
 450 mg amprolium (100%) (10 mg/kg) body weight per day for 21 days. Use one level scoop only when one or more calves show signs of coccidiosis. It is likely that the rest of the group has been exposed and all calves in the group should be treated.

To aid in the identification of CORID 25% in the medicated feed, prepare a solution of CORID 25% in a portion of complete feed. Ingestible before mixing into the finished ration. Read the directions with the remainder of the finished feed and mix thoroughly. The proper amount of amprolium supplement is related to the amount of ration usually consumed in one day. The table below gives suggested amounts for using Type C medicated feed.

21 DAY PREVENTION PROGRAM	CORID 25% (amprolium)			21 DAY TREATMENT PROGRAM	CORID 25% (amprolium)		
	4	3	2		4	3	2
Body Weight (lb)	150	222	4	150	454	4	3
Amprolium (mg)	4	3	2	454	4	3	2
Body Weight (kg)	68	101	1.8	68	206	1.8	1.4
Amprolium (mg)	1.8	1.4	0.8	206	1.8	1.4	0.8
Body Weight (kg)	300	405	6	300	908	6	4
Amprolium (mg)	6	4	2	908	6	4	2
Body Weight (kg)	300	405	6	300	908	6	4
Amprolium (mg)	6	4	2	908	6	4	2
Body Weight (kg)	600	908	12	600	1816	12	8
Amprolium (mg)	12	8	4	1816	12	8	4
Body Weight (kg)	600	908	12	600	1816	12	8
Amprolium (mg)	12	8	4	1816	12	8	4
Body Weight (kg)	1200	1816	24	1200	3632	24	16
Amprolium (mg)	24	16	8	3632	24	16	8

TAKE TIME
 OBSERVE LABEL
 DIRECTIONS

HUVEPHARMA

Distributed by:
 Huvepharma, Inc.
 Peachtree City, GA 30269

KCP-500014
 Version: 02
 Rev: 06/2022

Type A medicated article

CORID[®] 25%
 (amprolium)

CORID[®] 25%
 (amprolium)
 Type A medicated article
Coccidiostat

To be used in the preparation of medicated feed to aid in the prevention and treatment of coccidiosis in calves.

ACTIVE DRUG INGREDIENT: Amprolium.....25%

INGREDIENTS: Corn Gluten Feed and Soybean Oil

SEE DIRECTIONS ON BACK PANEL

STORAGE: Store at or below 77°F (25°C), excursions permitted to 104°F (40°C). Avoid prolonged exposure to high humidity.

Approved by FDA under NADA # 012-350

©CORID is a trademark of Huvepharma EOOD.

Net Wt.: 50 lb (22.68 kg)

HUVEPHARMA

CORID[®] 25%
 (amprolium)
 Type A medicated article

LOT NO. & EXPIRE DATE

CORID TYPE A MEDICATED ARTICLE			
amprolium granule			
Product Information			
Product Type	OTC TYPE A MEDICATED ARTICLE ANIMAL DRUG	Item Code (Source)	NDC:23243-9704
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
amprolium (UNII: 95CO6N199Q) (AMPROLIUM ION - UNII:H2T307KMZR)	amprolium	250 mg in 0.001 kg	
Inactive Ingredients			

Ingredient Name		Strength		
AMINO ACIDS, CORN GLUTEN (UNII: 0540V8ZD7V)				
SOYBEAN OIL (UNII: 241ATL177A)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:23243-9704-1	22.68 kg in 1 BAG		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
NADA	NADA012350	11/30/2017		

Labeler - Huvepharma, Inc (619153559)

Registrant - Huvepharma EOOD (552691651)

Establishment

Name	Address	ID/FEI	Business Operations
Huvepharma, Inc.		883128204	medicated animal feed manufacture, analysis, pack, label, manufacture

Revised: 12/2022

Huvepharma, Inc