

**CORID TYPE A MEDICATED ARTICLE- amprolium granule
Huvepharma, Inc.**

CORID® 25%

(amprolium)

Type A Medicated Article

Coccidiostat

**To be used in the preparation of medicated feed to aid
in the prevention and treatment of coccidiosis in calves.**

ACTIVE DRUG INGREDIENT: Amprolium 25%

INGREDIENTS: Corn Gluten Feed and Soybean Oil

SEE DIRECTIONS ON BACK PANEL

STORAGE: Store at or below 77°F (25°C), excursions permitted to 104°F (40°C).
Avoid prolonged exposure to high humidity.

Approved by FDA under NADA # 012-350

®CORID is a trademark of Huvepharma EOOD.

Net Wt.: 50 lb (22.68 kg)

CORID® 25%

(amprolium)

Type A Medicated Article

INDICATIONS FOR USE:

An aid in the prevention and treatment of coccidiosis caused by *Eimeria bovis* and *E. zurnii* in calves.

WITHDRAWAL PERIODS AND RESIDUE WARNINGS:

Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

USER SAFETY WARNING:

Not for human use. Keep out of reach of children.

Contact Huvepharma Inc. at 1-877-994-4883 or <http://www.huvepharma.us>. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

CAUTION: For satisfactory diagnosis, a microscopic examination of the feces should be done by a veterinarian or diagnostic laboratory before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

Do not use in feeds containing bentonite.

Use as the sole source of amprolium.

Restricted Drug (California) - Use only as directed.

MIXING DIRECTIONS: CORID 25% (amprolium) Type A medicated article should be thoroughly and evenly mixed in the feed. CORID 25% may be used to manufacture a Type B medicated feed with a concentration of 2.5% amprolium or a Type C medicated feed in the concentration range of 0.0125% to 1.25% amprolium.

- Use 200 pounds of CORID 25% per ton of feed to produce Type B medicated feed containing 2.5% amprolium.
- Use 1 pound of CORID 25% per ton of feed to produce Type C medicated feed containing 0.0125% amprolium.*
- Use 100 pounds of CORID 25% per ton of feed to produce Type C medicated feed containing 1.25% amprolium.
- Use 10 pounds of 2.5% Type B per ton of feed to produce Type C medicated feed containing 0.0125% amprolium.
- Use 1000 pounds of 2.5% Type B per ton of feed to produce Type C medicated feed containing 1.25% amprolium.

AMOUNT AND DIRECTIONS FOR USE:

Prevention: 227 mg amprolium /100 lb (5 mg/kg) body weight per day for 21 days during periods of exposure or when experience indicates that coccidiosis is likely to be a hazard.

Treatment: 454 mg amprolium/100 lb (10 mg/kg) bodyweight per day for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed and all calves in the group should be treated.

*To aid in even distribution of CORID 25% in the finished feed, prepare a mixture of CORID 25% in a portion of complete feed ingredient before mixing into the finished ration. Blend the mixture with the remainder of the finished feed and mix thoroughly.

The proper amount of amprolium supplement is mixed with the amount of ration consumed in one day. The tables below give suggested examples for using Type C medicated feed.

21 DAY PREVENTIVE PROGRAM:

DMI (as % Body Weight)		4	3	2
Body Weight (lb)	Amprolium per Day (mg)	0.0125% Amprolium	0.0167% Amprolium	0.025% Amprolium
		Amount of Feed per Day in Pounds		
100	227	4	3	2
200	454	8	6	4
300	681	12	9	6
400	908	16	12	8
500	1134	20	15	10
600	1361	24	18	12

5 DAY TREATMENT PROGRAM:

DMI (as % Body Weight)		4	3	2
Body Weight (lb)	Amprolium per Day (mg)	0.025% Amprolium	0.033% Amprolium	0.05% Amprolium
		Amount of Feed per Day in Pounds		
100	454	4	3	2
200	908	8	6	4
300	1362	12	9	6
400	1816	16	12	8
500	2268	20	15	10
600	2722	24	18	12



Distributed by:
Huvepharma, Inc.
Peachtree City, GA 30269

CORID 25%
(amprolium)
 Type A medicated article

INDICATIONS (IND USE):
 For use in the prevention and treatment of coccidiosis caused by *Coccidia* spp. in calves.

NET HORSEWEIGHT PERIODS AND RESIDUE WARNINGS:
 Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in growing calves to be processed for meat.

USE SAFETY WARNING:
 Not for human use. Resistant to most antibiotics.
 Contact Huvepharma Inc., 11877 New 40th St, Peachtree City, GA 30269 or (770) 486-4100 for additional information about reporting side effects for animals; contact FDA at 1-888-484-4100 or http://www.fda.gov/oc/ohrt/ohrt.html.
 CAUTION: For medicated feed, it is a complete concentration of the feed should be done to determine if the work is necessary before treatment. When feeding outside, drug should be administered promptly after the work is determined.
 Do not use in feed containing tetracyclines.
 Use in the water source of large animals.
 Medication free (all-chemical) - does not contain tetracyclines.

ADMINISTRATION: CORID 25% (amprolium) Type A medicated article should be thoroughly and evenly mixed in the feed. CORID 25% may be used to manufacture a Type A medicated feed with a concentration of 2.5% amprolium or a Type C medicated feed in the concentration range of 0.0025% to 1.25% amprolium.

- Use 700 pounds of CORID 25% per ton of feed to produce Type B medicated feed containing 2.5% amprolium.
- Use 1 pound of CORID 25% per ton of feed to produce Type C medicated feed containing 0.0025% amprolium.
- Use 100 pounds of CORID 25% per ton of feed to produce Type C medicated feed containing 1.25% amprolium.
- Use 10 pounds of 2.5% Type B per ton of feed to produce Type C medicated feed containing 0.0025% amprolium.
- Use 1000 pounds of 2.5% Type B per ton of feed to produce Type C medicated feed containing 1.25% amprolium.

ADMINISTRATIVE INFORMATION:
Prevention: 222 mg amprolium / 100 lb (5 mg/kg) body weight per day for 21 days during periods of exposure or when exposure indicates that coccidiosis is likely to be a concern.
Treatment: 451 mg amprolium / 100 lb (5 mg/kg) body weight per day for 21 days. Use on a hard feed only when one or more calves show signs of coccidiosis, & it likely that the rest of the group has been exposed and all calves in the group should be treated.

To aid in even distribution of CORID 25% in the finished feed, prepare a mixture of CORID 25% in a portion of complete feed ingredient before adding into the finished feed. Blend the mixture with the remainder of the finished feed and re-challenge. The proper amount of complete feed ingredient to blend with the amount of ration usually consumed in one day. Rechallenge before give suggested examples for using Type C medicated feed.

2.5% Type B medicated feed	1			2					
	Amprolium (lb)	Amprolium (kg)	Amprolium (g)	Amprolium (lb)	Amprolium (kg)	Amprolium (g)			
100	222	4	3	1	100	454	4	3	2
200	454	8	6	4	200	908	8	6	4
300	681	12	9	6	300	1362	12	9	6
400	908	16	12	8	400	1816	16	12	8
500	1135	20	15	10	500	2270	20	15	10
600	1362	24	18	12	600	2724	24	18	12

2.5% Type C medicated feed

2.5% Type C medicated feed	1			2				
	Amprolium (lb)	Amprolium (kg)	Amprolium (g)	Amprolium (lb)	Amprolium (kg)	Amprolium (g)		
100	100	45	34	100	454	4	3	2
200	200	90	68	200	908	8	6	4
300	300	135	102	300	1362	12	9	6
400	400	180	136	400	1816	16	12	8
500	500	225	170	500	2270	20	15	10
600	600	270	204	600	2724	24	18	12

TAKE TIME
OBSERVE LABEL DIRECTIONS

HUVEPHARMA
 KCP-508014
 Version 03
 Rev. 06/2022

CORID 25%
(amprolium)
 Type A medicated article

245532
 11-13
 511403

CORID 25%
(amprolium)
 Type A medicated article
Coccidiostat

To be used in the preparation of medicated feed to aid in the prevention and treatment of coccidiosis in calves.

ACTIVE DRUG INGREDIENT: Amprolium.....25%

INGREDIENTS: Corn Gluten Feed and Soybean Oil

SEE DIRECTIONS ON BACK PANEL

STORAGE: Store at or below 77°F (25°C), excursions permitted to 104°F (40°C). Avoid prolonged exposure to high humidity.

Approved by FDA under NADA # 012-350

©CORID is a trademark of Huvepharma EOOD.

Net Wt.: 50 lb (22.68 kg)

HUVEPHARMA

CORID 25%
(amprolium)
 Type A medicated article

LOT NO. & EXP. DATE

CORID TYPE A MEDICATED ARTICLE			
amprolium granule			
Product Information			
Product Type	OTC TYPE A MEDICATED ARTICLE ANIMAL DRUG	Item Code (Source)	NDC:23243-9704
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
amprolium (UNII: 95CO6N199Q) (AMPROLIUM ION - UNII:H2T307KMZR)	amprolium	250 mg in 0.001 kg	
Inactive Ingredients			

Ingredient Name		Strength		
AMINO ACIDS, CORN GLUTEN (UNII: 0540V8ZD7V)				
SOYBEAN OIL (UNII: 241ATL177A)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:23243-9704-1	22.68 kg in 1 BAG		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
NADA	NADA012350	11/30/2017		

Labeler - Huvepharma, Inc. (619153559)

Registrant - Huvepharma EOOD (507423867)

Establishment

Name	Address	ID/FEI	Business Operations
Huvepharma, Inc.		883128204	medicated animal feed manufacture, analysis, pack, label, manufacture

Revised: 10/2025

Huvepharma, Inc.