

**JANTOVEN- warfarin sodium tablet**  
**Upsher-Smith Laboratories, LLC**

-----

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

**These highlights do not include all the information needed to use JANTOVEN safely and effectively. See full prescribing information for JANTOVEN.**

**Jantoven® Tablets (Warfarin Sodium Tablets, USP), for oral use**  
**Initial U.S. Approval: 1954**

**WARNING: BLEEDING RISK**

***See full prescribing information for complete boxed warning.***

- Warfarin Sodium can cause major or fatal bleeding. ( 5.1)
- Perform regular monitoring of INR in all treated patients. ( 2.1)
- Drugs, dietary changes, and other factors affect INR levels achieved with Warfarin Sodium therapy. ( 7)
- Instruct patients about prevention measures to minimize risk of bleeding and to report signs and symptoms of bleeding. ( 17)

-----  
**RECENT MAJOR CHANGES**  
-----

Dosage and Administration, Renal Impairment ( 2.5)	5/2017
Warnings and Precautions, Calciphylaxis ( 5.3)	9/2016
Warnings and Precautions, Acute kidney injury ( 5.4)	5/2017

-----  
**INDICATIONS AND USAGE**  
-----

JANTOVEN is a vitamin K antagonist indicated for:

- Prophylaxis and treatment of venous thrombosis and its extension, pulmonary embolism ( 1)
- Prophylaxis and treatment of thromboembolic complications associated with atrial fibrillation and/or cardiac valve replacement ( 1)
- Reduction in the risk of death, recurrent myocardial infarction, and thromboembolic events such as stroke or systemic embolization after myocardial infarction ( 1)

**Limitations of Use**

Warfarin Sodium has no direct effect on an established thrombus, nor does it reverse ischemic tissue damage. ( 1)

-----  
**DOSAGE AND ADMINISTRATION**  
-----

- Individualize dosing regimen for each patient, and adjust based on INR response. ( 2.1, 2.2)
- Knowledge of genotype can inform initial dose selection. ( 2.3)
- Monitoring: Obtain daily INR determinations upon initiation until stable in the therapeutic range. Obtain subsequent INR determinations every 1 to 4 weeks. ( 2.4)
- Review conversion instructions from other anticoagulants. ( 2.8)

-----  
**DOSAGE FORMS AND STRENGTHS**  
-----

Scored tablets: 1, 2, 2½, 3, 4, 5, 6, 7½, or 10 mg ( 3)

-----  
**CONTRAINDICATIONS**  
-----

- Pregnancy, except in women with mechanical heart valves ( 4, 5.7, 8.1)
- Hemorrhagic tendencies or blood dyscrasias ( 4)
- Recent or contemplated surgery of the central nervous system (CNS) or eye, or traumatic surgery resulting in large open surfaces ( 4, 5.8)
- Bleeding tendencies associated with certain conditions ( 4)
- Threatened abortion, eclampsia, and preeclampsia ( 4)
- Unsupervised patients with potential high levels of non-compliance ( 4)
- Spinal puncture and other diagnostic or therapeutic procedures with potential for uncontrollable bleeding ( 4)

- Hypersensitivity to warfarin or any component of the product ( 4)
- Major regional or lumbar block anesthesia ( 4)
- Malignant hypertension ( 4)

#### -----**WARNINGS AND PRECAUTIONS**-----

- Tissue necrosis: Necrosis or gangrene of skin or other tissues can occur, with severe cases requiring debridement or amputation. Discontinue JANTOVEN and consider alternative anticoagulants if necessary. ( 5.2)
- Calciphylaxis: Fatal and serious cases have occurred. Discontinue JANTOVEN and consider alternative anticoagulation therapy. ( 5.3)
- Acute kidney injury may occur during episodes of excessive anticoagulation and hematuria. ( 5.4)
- Systemic atheroemboli and cholesterol microemboli: Some cases have progressed to necrosis or death. Discontinue JANTOVEN if such emboli occur. ( 5.5)
- Heparin-induced thrombocytopenia (HIT): Initial therapy with Warfarin Sodium in HIT has resulted in cases of amputation and death. Warfarin Sodium may be considered after platelet count has normalized. ( 5.6)
- Pregnant women with mechanical heart valves: Warfarin Sodium may cause fetal harm; however, the benefits may outweigh the risks. ( 5.7)

#### -----**ADVERSE REACTIONS**-----

Most common adverse reactions to Warfarin Sodium are fatal and nonfatal hemorrhage from any tissue or organ. ( 6)

**To report SUSPECTED ADVERSE REACTIONS, contact Upsher-Smith Laboratories, LLC at 1-855-899-9180 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

#### -----**DRUG INTERACTIONS**-----

- Concomitant use of drugs that increase bleeding risk, antibiotics, antifungals, botanical (herbal) products, and inhibitors and inducers of CYP2C9, 1A2, or 3A4. ( 7)
- Consult labeling of all concurrently used drugs for complete information about interactions with JANTOVEN or increased risks for bleeding. ( 7)

#### -----**USE IN SPECIFIC POPULATIONS**-----

- Pregnant women with mechanical heart valves: JANTOVEN may cause fetal harm; however, the benefits may outweigh the risks. ( 8.1)
- Lactation: Monitor breastfeeding infants for bruising or bleeding. ( 8.2)
- Renal Impairment: Instruct patients with renal impairment to frequently monitor their INR. ( 8.6)

**See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.**

**Revised: 9/2017**

---

## **FULL PRESCRIBING INFORMATION: CONTENTS\***

### **WARNING: BLEEDING RISK**

### **1 INDICATIONS AND USAGE**

### **2 DOSAGE AND ADMINISTRATION**

2.1 Individualized Dosing

2.2 Recommended Target INR Ranges and Durations for Individual Indications

2.3 Initial and Maintenance Dosing

2.4 Monitoring to Achieve Optimal Anticoagulation

2.5 Renal Impairment

2.6 Missed Dose

2.7 Treatment during Dentistry and Surgery

2.8 Conversion from Other Anticoagulants

### **3 DOSAGE FORMS AND STRENGTHS**

## **4 CONTRAINDICATIONS**

## **5 WARNINGS AND PRECAUTIONS**

- 5.1 Hemorrhage
- 5.2 Tissue Necrosis
- 5.3 Calciphylaxis
- 5.4 Acute Kidney Injury
- 5.5 Systemic Atheroemboli and Cholesterol Microemboli
- 5.6 Limb Ischemia, Necrosis, and Gangrene in Patients with HIT and HITTS
- 5.7 Use in Pregnant Women with Mechanical Heart Valves
- 5.8 Other Clinical Settings with Increased Risks
- 5.9 Endogenous Factors Affecting INR

## **6 ADVERSE REACTIONS**

## **7 DRUG INTERACTIONS**

- 7.1 General Information
- 7.2 CYP450 Interactions
- 7.3 Drugs that Increase Bleeding Risk
- 7.4 Antibiotics and Antifungals
- 7.5 Botanical (Herbal) Products and Foods

## **8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females and Males of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 8.7 Hepatic Impairment

## **10 OVERDOSAGE**

- 10.1 Signs and Symptoms
- 10.2 Treatment

## **11 DESCRIPTION**

## **12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics
- 12.5 Pharmacogenomics

## **13 NONCLINICAL TOXICOLOGY**

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

## **14 CLINICAL STUDIES**

- 14.1 Atrial Fibrillation
- 14.2 Mechanical and Bioprosthetic Heart Valves
- 14.3 Myocardial Infarction

## **15 REFERENCES**

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

## **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

### **WARNING: BLEEDING RISK**

- **Warfarin Sodium can cause major or fatal bleeding [see *Warnings and Precautions (5.1)*].**
- **Perform regular monitoring of INR in all treated patients [see *Dosage and Administration (2.1)*].**
- **Drugs, dietary changes, and other factors affect INR levels achieved with Warfarin Sodium therapy [see *Drug Interactions (7)*].**
- **Instruct patients about prevention measures to minimize risk of bleeding and to report signs and symptoms of bleeding [see *Patient Counseling Information (17)*].**

## **1 INDICATIONS AND USAGE**

JANTOVEN<sup>®</sup> is indicated for:

- Prophylaxis and treatment of venous thrombosis and its extension, pulmonary embolism (PE).
- Prophylaxis and treatment of thromboembolic complications associated with atrial fibrillation (AF) and/or cardiac valve replacement.
- Reduction in the risk of death, recurrent myocardial infarction (MI), and thromboembolic events such as stroke or systemic embolization after myocardial infarction.

### **Limitations of Use**

Warfarin Sodium has no direct effect on an established thrombus, nor does it reverse ischemic tissue damage. Once a thrombus has occurred, however, the goals of anticoagulant treatment are to prevent further extension of the formed clot and to prevent secondary thromboembolic complications that may result in serious and possibly fatal sequelae.

## **2 DOSAGE AND ADMINISTRATION**

### **2.1 Individualized Dosing**

The dosage and administration of JANTOVEN must be individualized for each patient according to the patient's International Normalized Ratio (INR) response to the drug. Adjust the dose based on the patient's INR and the condition being treated. Consult the latest evidence-based clinical practice guidelines regarding the duration and intensity of anticoagulation for the indicated conditions.

### **2.2 Recommended Target INR Ranges and Durations for Individual Indications**

**An INR of greater than 4.0 appears to provide no additional therapeutic benefit in most patients and is associated with a higher risk of bleeding.**

**Venous Thromboembolism (including deep venous thrombosis [DVT] and PE)**

Adjust the warfarin dose to maintain a target INR of 2.5 (INR range, 2.0 to 3.0) for all

treatment durations. The duration of treatment is based on the indication as follows:

- For patients with a DVT or PE secondary to a transient (reversible) risk factor, treatment with warfarin for 3 months is recommended.
- For patients with an unprovoked DVT or PE, treatment with warfarin is recommended for at least 3 months. After 3 months of therapy, evaluate the risk-benefit ratio of long-term treatment for the individual patient.
- For patients with two episodes of unprovoked DVT or PE, long-term treatment with warfarin is recommended. For a patient receiving long-term anticoagulant treatment, periodically reassess the risk-benefit ratio of continuing such treatment in the individual patient.

## **Atrial Fibrillation**

In patients with non-valvular AF, anticoagulate with warfarin to target INR of 2.5 (range, 2.0 to 3.0).

- In patients with non-valvular AF that is persistent or paroxysmal and at high risk of stroke (i.e., having any of the following features: prior ischemic stroke, transient ischemic attack, or systemic embolism, or 2 of the following risk factors: age greater than 75 years, moderately or severely impaired left ventricular systolic function and/or heart failure, history of hypertension, or diabetes mellitus), long-term anticoagulation with warfarin is recommended.
- In patients with non-valvular AF that is persistent or paroxysmal and at an intermediate risk of ischemic stroke (i.e., having 1 of the following risk factors: age greater than 75 years, moderately or severely impaired left ventricular systolic function and/or heart failure, history of hypertension, or diabetes mellitus), long-term anticoagulation with warfarin is recommended.
- For patients with AF and mitral stenosis, long-term anticoagulation with warfarin is recommended.
- For patients with AF and prosthetic heart valves, long-term anticoagulation with warfarin is recommended; the target INR may be increased and aspirin added depending on valve type and position, and on patient factors.

## **Mechanical and Bioprosthetic Heart Valves**

- For patients with a bileaflet mechanical valve or a Medtronic Hall (Minneapolis, MN) tilting disk valve in the aortic position who are in sinus rhythm and without left atrial enlargement, therapy with warfarin to a target INR of 2.5 (range, 2.0 to 3.0) is recommended.
- For patients with tilting disk valves and bileaflet mechanical valves in the mitral position, therapy with warfarin to a target INR of 3.0 (range, 2.5 to 3.5) is recommended.
- For patients with caged ball or caged disk valves, therapy with warfarin to a target INR of 3.0 (range, 2.5 to 3.5) is recommended.
- For patients with a bioprosthetic valve in the mitral position, therapy with warfarin to a target INR of 2.5 (range, 2.0 to 3.0) for the first 3 months after valve insertion is recommended. If additional risk factors for thromboembolism are present (AF, previous thromboembolism, left ventricular dysfunction), a target INR of 2.5 (range, 2.0 to 3.0) is recommended.

## **Post-Myocardial Infarction**

- For high-risk patients with MI (e.g., those with a large anterior MI, those with

significant heart failure, those with intracardiac thrombus visible on transthoracic echocardiography, those with AF, and those with a history of a thromboembolic event), therapy with combined moderate-intensity (INR, 2.0 to 3.0) warfarin plus low-dose aspirin ( $\leq 100$  mg/day) for at least 3 months after the MI is recommended.

## Recurrent Systemic Embolism and Other Indications

Oral anticoagulation therapy with warfarin has not been fully evaluated by clinical trials in patients with valvular disease associated with AF, patients with mitral stenosis, and patients with recurrent systemic embolism of unknown etiology. However, a moderate dose regimen (INR 2.0 to 3.0) may be used for these patients.

## 2.3 Initial and Maintenance Dosing

The appropriate initial dosing of JANTOVEN varies widely for different patients. Not all factors responsible for warfarin dose variability are known, and the initial dose is influenced by:

- Clinical factors including age, race, body weight, sex, concomitant medications, and comorbidities
- Genetic factors (CYP2C9 and VKORC1 genotypes) [see *Clinical Pharmacology (12.5)*]

Select the initial dose based on the expected maintenance dose, taking into account the above factors. Modify this dose based on consideration of patient-specific clinical factors. Consider lower initial and maintenance doses for elderly and/or debilitated patients and in Asian patients [see *Use in Specific Populations (8.5)* and *Clinical Pharmacology (12.3)*]. Routine use of loading doses is not recommended as this practice may increase hemorrhagic and other complications and does not offer more rapid protection against clot formation.

Individualize the duration of therapy for each patient. In general, anticoagulant therapy should be continued until the danger of thrombosis and embolism has passed [see *Dosage and Administration (2.2)*].

## Dosing Recommendations without Consideration of Genotype

If the patient's CYP2C9 and VKORC1 genotypes are not known, the initial dose of JANTOVEN is usually 2 to 5 mg once daily. Determine each patient's dosing needs by close monitoring of the INR response and consideration of the indication being treated. Typical maintenance doses are 2 to 10 mg once daily.

## Dosing Recommendations with Consideration of Genotype

Table 1 displays three ranges of expected maintenance JANTOVEN doses observed in subgroups of patients having different combinations of CYP2C9 and VKORC1 gene variants [see *Clinical Pharmacology (12.5)*]. If the patient's CYP2C9 and/or VKORC1 genotype are known, consider these ranges in choosing the initial dose. Patients with CYP2C9 \*1/\*3, \*2/\*2, \*2/\*3, and \*3/\*3 may require more prolonged time (>2 to 4 weeks) to achieve maximum INR effect for a given dosage regimen than patients without these CYP variants.

**Table 1: Three Ranges of Expected Maintenance JANTOVEN Daily Doses Based on CYP2C9 and VKORC1 Genotypes \***

VKORC1	CYP2C9
--------	--------

	<b>*1/*1</b>	<b>*1/*2</b>	<b>*1/*3</b>	<b>*2/*2</b>	<b>*2/*3</b>	<b>*3/*3</b>
GG	5-7 mg	5-7 mg	3-4 mg	3-4 mg	3-4 mg	0.5-2 mg
AG	5-7 mg	3-4 mg	3-4 mg	3-4 mg	0.5-2 mg	0.5-2 mg
AA	3-4 mg	3-4 mg	0.5-2 mg	0.5-2 mg	0.5-2 mg	0.5-2 mg

\* Ranges are derived from multiple published clinical studies. VKORC1-1639G>A (rs9923231) variant is used in this table. Other co-inherited VKORC1 variants may also be important determinants of warfarin dose.

## 2.4 Monitoring to Achieve Optimal Anticoagulation

JANTOVEN has a narrow therapeutic range (index), and its action may be affected by factors such as other drugs and dietary vitamin K. Therefore, anticoagulation must be carefully monitored during JANTOVEN therapy. Determine the INR daily after the administration of the initial dose until INR results stabilize in the therapeutic range. After stabilization, maintain dosing within the therapeutic range by performing periodic INRs. The frequency of performing INR should be based on the clinical situation but generally acceptable intervals for INR determinations are 1 to 4 weeks. Perform additional INR tests when other warfarin products are interchanged with JANTOVEN, as well as whenever other medications are initiated, discontinued, or taken irregularly. Heparin, a common concomitant drug, increases the INR [see *Dosage and Administration (2.8)* and *Drug Interactions (7)*].

Determinations of whole blood clotting and bleeding times are not effective measures for monitoring of JANTOVEN therapy.

## 2.5 Renal Impairment

No dosage adjustment is necessary for patients with renal failure. Monitor INR more frequently in patients with compromised renal function to maintain INR within the therapeutic range [see *Warnings and Precautions (5.4)* and *Use in Specific Populations (8.6)*].

## 2.6 Missed Dose

The anticoagulant effect of JANTOVEN persists beyond 24 hours. If a patient misses a dose of JANTOVEN at the intended time of day, the patient should take the dose as soon as possible on the same day. The patient should not double the dose the next day to make up for a missed dose.

## 2.7 Treatment during Dentistry and Surgery

Some dental or surgical procedures may necessitate the interruption or change in the dose of JANTOVEN therapy. Consider the benefits and risks when discontinuing JANTOVEN even for a short period of time. Determine the INR immediately prior to any dental or surgical procedure. In patients undergoing minimally invasive procedures who must be anticoagulated prior to, during, or immediately following these procedures, adjusting the dosage of JANTOVEN to maintain the INR at the low end of the therapeutic range may safely allow for continued anticoagulation.

## 2.8 Conversion from Other Anticoagulants

### Heparin

Since the full anticoagulant effect of JANTOVEN is not achieved for several days, heparin is preferred for initial rapid anticoagulation. During initial therapy with JANTOVEN, the interference with heparin anticoagulation is of minimal clinical significance. Conversion to JANTOVEN may begin concomitantly with heparin therapy or may be delayed 3 to 6 days. To ensure therapeutic anticoagulation, continue full dose heparin therapy and overlap JANTOVEN therapy with heparin for 4 to 5 days and until JANTOVEN has produced the desired therapeutic response as determined by INR, at which point heparin may be discontinued.

As heparin may affect the INR, patients receiving both heparin and JANTOVEN should have INR monitoring at least:

- 5 hours after the last intravenous bolus dose of heparin, or
- 4 hours after cessation of a continuous intravenous infusion of heparin, or
- 24 hours after the last subcutaneous heparin injection.

JANTOVEN may increase the activated partial thromboplastin time (aPTT) test, even in the absence of heparin. A severe elevation (>50 seconds) in aPTT with an INR in the desired range has been identified as an indication of increased risk of postoperative hemorrhage.

### **Other Anticoagulants**

Consult the labeling of other anticoagulants for instructions on conversion to JANTOVEN.

## **3 DOSAGE FORMS AND STRENGTHS**

JANTOVEN tablets are single scored, compressed tablets with one side scored and debossed with WRF above the score and 1, 2, 2½, 3, 4, 5, 6, 7½, or 10 below the score and with 832 debossed on the opposite side.

JANTOVEN tablets are supplied in the following strengths:

### **JANTOVEN Tablets**

<b>Strength</b>	<b>Color</b>
1 mg	pink
2 mg	lavender
2½ mg	green
3 mg	tan
4 mg	blue
5 mg	peach
6 mg	teal
7½ mg	yellow
10 mg	white (dye free)

## **4 CONTRAINDICATIONS**

JANTOVEN is contraindicated in:

- Pregnancy

JANTOVEN is contraindicated in women who are pregnant except in pregnant women with mechanical heart valves, who are at high risk of thromboembolism [see *Warnings and Precautions (5.7)* and *Use in Specific Populations (8.1)*]. JANTOVEN can cause fetal harm when administered to a pregnant woman. JANTOVEN exposure during pregnancy causes a recognized pattern of major congenital malformations (warfarin embryopathy and fetotoxicity), fatal fetal hemorrhage, and an increased risk of spontaneous abortion and fetal mortality. If JANTOVEN is used during pregnancy or if the patient becomes pregnant while taking this drug, the patient should be apprised of the potential hazard to a fetus [see *Use in Specific Populations (8.1)*].

JANTOVEN is contraindicated in patients with:

- Hemorrhagic tendencies or blood dyscrasias
- Recent or contemplated surgery of the central nervous system or eye, or traumatic surgery resulting in large open surfaces [see *Warnings and Precautions (5.8)*]
- Bleeding tendencies associated with:
  - Active ulceration or overt bleeding of the gastrointestinal, genitourinary, or respiratory tract
  - Central nervous system hemorrhage
  - Cerebral aneurysms, dissecting aorta
  - Pericarditis and pericardial effusions
  - Bacterial endocarditis
- Threatened abortion, eclampsia, and preeclampsia
- Unsupervised patients with conditions associated with potential high level of non-compliance
- Spinal puncture and other diagnostic or therapeutic procedures with potential for uncontrollable bleeding
- Hypersensitivity to warfarin or to any other components of this product (e.g., anaphylaxis) [see *Adverse Reactions (6)*]
- Major regional or lumbar block anesthesia
- Malignant hypertension

## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Hemorrhage**

JANTOVEN can cause major or fatal bleeding. Bleeding is more likely to occur within the first month. Risk factors for bleeding include high intensity of anticoagulation (INR >4.0), age greater than or equal to 65, history of highly variable INRs, history of gastrointestinal bleeding, hypertension, cerebrovascular disease, anemia, malignancy, trauma, renal impairment, certain genetic factors [see *Clinical Pharmacology (12.5)*], certain concomitant drugs [see *Drug Interactions (7)*], and long duration of warfarin therapy.

Perform regular monitoring of INR in all treated patients. Those at high risk of bleeding may benefit from more frequent INR monitoring, careful dose adjustment to desired INR, and a shortest duration of therapy appropriate for the clinical condition. However, maintenance of INR in the therapeutic range does not eliminate the risk of bleeding.

Drugs, dietary changes, and other factors affect INR levels achieved with JANTOVEN therapy. Perform more frequent INR monitoring when starting or stopping other drugs,

including botanicals, or when changing dosages of other drugs [see *Drug Interactions (7)*].

Instruct patients about prevention measures to minimize risk of bleeding and to report signs and symptoms of bleeding [see *Patient Counseling Information (17)*].

## **5.2 Tissue Necrosis**

JANTOVEN can cause necrosis and/or gangrene of skin and other tissues, which is an uncommon but serious risk (<0.1%). Necrosis may be associated with local thrombosis and usually appears within a few days of the start of JANTOVEN therapy. In severe cases of necrosis, treatment through debridement or amputation of the affected tissue, limb, breast, or penis has been reported.

Careful clinical evaluation is required to determine whether necrosis is caused by an underlying disease. Although various treatments have been attempted, no treatment for necrosis has been considered uniformly effective. Discontinue JANTOVEN therapy if necrosis occurs. Consider alternative drugs if continued anticoagulation therapy is necessary.

## **5.3 Calciphylaxis**

JANTOVEN can cause fatal and serious calciphylaxis or calcium uremic arteriopathy, which has been reported in patients with and without end-stage renal disease. When calciphylaxis is diagnosed in these patients, discontinue JANTOVEN and treat calciphylaxis as appropriate. Consider alternative anticoagulation therapy.

## **5.4 Acute Kidney Injury**

In patients with altered glomerular integrity or with a history of kidney disease, acute kidney injury may occur with JANTOVEN, possibly in relation to episodes of excessive anticoagulation and hematuria [ see *Use in Specific Populations (8.6)*]. More frequent monitoring of anticoagulation is advised in patients with compromised renal function.

## **5.5 Systemic Atheroemboli and Cholesterol Microemboli**

Anticoagulation therapy with JANTOVEN may enhance the release of atheromatous plaque emboli. Systemic atheroemboli and cholesterol microemboli can present with a variety of signs and symptoms depending on the site of embolization. The most commonly involved visceral organs are the kidneys followed by the pancreas, spleen, and liver. Some cases have progressed to necrosis or death. A distinct syndrome resulting from microemboli to the feet is known as "purple toes syndrome." Discontinue JANTOVEN therapy if such phenomena are observed. Consider alternative drugs if continued anticoagulation therapy is necessary.

## **5.6 Limb Ischemia, Necrosis, and Gangrene in Patients with HIT and HITTS**

Do not use JANTOVEN as initial therapy in patients with heparin-induced thrombocytopenia (HIT) and with heparin-induced thrombocytopenia with thrombosis syndrome (HITTS). Cases of limb ischemia, necrosis, and gangrene have occurred in patients with HIT and HITTS when heparin treatment was discontinued and warfarin therapy was started or continued. In some patients, sequelae have included amputation of the involved area and/or death. Treatment with JANTOVEN may be considered after

the platelet count has normalized.

## 5.7 Use in Pregnant Women with Mechanical Heart Valves

JANTOVEN can cause fetal harm when administered to a pregnant woman. While JANTOVEN is contraindicated during pregnancy, the potential benefits of using JANTOVEN may outweigh the risks for pregnant women with mechanical heart valves at high risk of thromboembolism. In those individual situations, the decision to initiate or continue JANTOVEN should be reviewed with the patient, taking into consideration the specific risks and benefits pertaining to the individual patient's medical situation, as well as the most current medical guidelines. JANTOVEN exposure during pregnancy causes a recognized pattern of major congenital malformations (warfarin embryopathy and fetotoxicity), fatal fetal hemorrhage, and an increased risk of spontaneous abortion and fetal mortality. If this drug is used during pregnancy, or if the patient becomes pregnant while taking this drug, the patient should be apprised of the potential hazard to a fetus [see *Use in Specific Populations (8.1)*].

## 5.8 Other Clinical Settings with Increased Risks

In the following clinical settings, the risks of JANTOVEN therapy may be increased:

- Moderate to severe hepatic impairment
- Infectious diseases or disturbances of intestinal flora (e.g., sprue, antibiotic therapy)
- Use of an indwelling catheter
- Severe to moderate hypertension
- Deficiency in protein C-mediated anticoagulant response: JANTOVEN reduces the synthesis of the naturally occurring anticoagulants, protein C and protein S. Hereditary or acquired deficiencies of protein C or its cofactor, protein S, have been associated with tissue necrosis following warfarin administration. Concomitant anticoagulation therapy with heparin for 5 to 7 days during initiation of therapy with JANTOVEN may minimize the incidence of tissue necrosis in these patients.
- Eye surgery: In cataract surgery, JANTOVEN use was associated with a significant increase in minor complications of sharp needle and local anesthesia block but not associated with potentially sight-threatening operative hemorrhagic complications. As JANTOVEN cessation or reduction may lead to serious thromboembolic complications, the decision to discontinue JANTOVEN before a relatively less invasive and complex eye surgery, such as lens surgery, should be based upon the risks of anticoagulant therapy weighed against the benefits.
- Polycythemia vera
- Vasculitis
- Diabetes mellitus

## 5.9 Endogenous Factors Affecting INR

The following factors may be responsible for **increased** INR response: diarrhea, hepatic disorders, poor nutritional state, steatorrhea, or vitamin K deficiency.

The following factors may be responsible for **decreased** INR response: increased vitamin K intake or hereditary warfarin resistance.

## 6 ADVERSE REACTIONS

The following serious adverse reactions to JANTOVEN are discussed in greater detail in other sections of the labeling:

- Hemorrhage [see *Boxed Warning, Warnings and Precautions (5.1)*, and *Overdosage (10)*]
- Tissue Necrosis [see *Warnings and Precautions (5.2)*]
- Calciphylaxis [see *Warnings and Precautions (5.3)*]
- Acute Kidney Injury [see *Warnings and Precautions (5.4)*]
- Systemic Atheroemboli and Cholesterol Microemboli [see *Warnings and Precautions (5.5)*]
- Limb Ischemia, Necrosis, and Gangrene in Patients with HIT and HITTS [see *Warnings and Precautions (5.6)*]
- Other Clinical Settings with Increased Risks [see *Warnings and Precautions (5.8)*]

Other adverse reactions to JANTOVEN include:

- Immune system disorders: hypersensitivity/allergic reactions (including urticaria and anaphylactic reactions)
- Vascular disorders: vasculitis
- Hepatobiliary disorders: hepatitis, elevated liver enzymes. Cholestatic hepatitis has been associated with concomitant administration of warfarin sodium and ticlopidine.
- Gastrointestinal disorders: nausea, vomiting, diarrhea, taste perversion, abdominal pain, flatulence, bloating
- Skin disorders: rash, dermatitis (including bullous eruptions), pruritus, alopecia
- Respiratory disorders: tracheal or tracheobronchial calcification
- General disorders: chills

## **7 DRUG INTERACTIONS**

### **7.1 General Information**

Drugs may interact with JANTOVEN through pharmacodynamic or pharmacokinetic mechanisms. Pharmacodynamic mechanisms for drug interactions with JANTOVEN are synergism (impaired hemostasis, reduced clotting factor synthesis), competitive antagonism (vitamin K), and alteration of the physiologic control loop for vitamin K metabolism (hereditary resistance). Pharmacokinetic mechanisms for drug interactions with JANTOVEN are mainly enzyme induction, enzyme inhibition, and reduced plasma protein binding. It is important to note that some drugs may interact by more than one mechanism.

More frequent INR monitoring should be performed when starting or stopping other drugs, including botanicals, or when changing dosages of other drugs, including drugs intended for short-term use (e.g., antibiotics, antifungals, corticosteroids) [see *Boxed Warning*].

Consult the labeling of all concurrently used drugs to obtain further information about interactions with JANTOVEN or adverse reactions pertaining to bleeding.

### **7.2 CYP450 Interactions**

CYP450 isozymes involved in the metabolism of warfarin include CYP2C9, 2C19, 2C8, 2C18, 1A2, and 3A4. The more potent warfarin *S*-enantiomer is metabolized by CYP2C9

while the *R*-enantiomer is metabolized by CYP1A2 and 3A4.

- Inhibitors of CYP2C9, 1A2, and/or 3A4 have the potential to increase the effect (increase INR) of warfarin by increasing the exposure of warfarin.
- Inducers of CYP2C9, 1A2, and/or 3A4 have the potential to decrease the effect (decrease INR) of warfarin by decreasing the exposure of warfarin.

Examples of inhibitors and inducers of CYP2C9, 1A2, and 3A4 are below in Table 2; however, this list should not be considered all-inclusive. Consult the labeling of all concurrently used drugs to obtain further information about CYP450 interaction potential. The CYP450 inhibition and induction potential should be considered when starting, stopping, or changing dose of concomitant medications. Closely monitor INR if a concomitant drug is a CYP2C9, 1A2, and/or 3A4 inhibitor or inducer.

**Table 2: Examples of CYP450 Interactions with Warfarin**

<b>Enzyme</b>	<b>Inhibitors</b>	<b>Inducers</b>
CYP2C9	amiodarone, capecitabine, cotrimoxazole, etravirine, fluconazole, fluvastatin, fluvoxamine, metronidazole, miconazole, oxandrolone, sulfapyridine, tigecycline, voriconazole, zafirlukast	aprepitant, bosentan, carbamazepine, phenobarbital, rifampin
CYP1A2	acyclovir, allopurinol, caffeine, cimetidine, ciprofloxacin, disulfiram, enoxacin, famotidine, fluvoxamine, methoxsalen, mexiletine, norfloxacin, oral contraceptives, phenylpropanolamine, propafenone, propranolol, terbinafine, thiabendazole, ticlopidine, verapamil, zileuton	montelukast, moricizine, omeprazole, phenobarbital, phenytoin, cigarette smoking
CYP3A4	alprazolam, amiodarone, amlodipine, amprenavir, aprepitant, atorvastatin, atazanavir, bicalutamide, cilostazol, cimetidine, ciprofloxacin, clarithromycin, conivaptan, cyclosporine, darunavir/ritonavir, diltiazem, erythromycin, fluconazole, fluoxetine, fluvoxamine, fosamprenavir, imatinib, indinavir, isoniazid, itraconazole, ketoconazole, lopinavir/ritonavir, nefazodone, nelfinavir, nilotinib, oral contraceptives, posaconazole, ranitidine, ranolazine, ritonavir, saquinavir, telithromycin, tipranavir, voriconazole, zileuton	armodafinil, amprenavir, aprepitant, bosentan, carbamazepine, efavirenz, etravirine, modafinil, nafcillin, phenytoin, pioglitazone, prednisone, rifampin, rufinamide

### 7.3 Drugs that Increase Bleeding Risk

Examples of drugs known to increase the risk of bleeding are presented in Table 3. Because bleeding risk is increased when these drugs are used concomitantly with warfarin, closely monitor patients receiving any such drug with warfarin.

**Table 3: Drugs that Can Increase the Risk of Bleeding**

<b>Drug Class</b>	<b>Specific Drugs</b>
Anticoagulants	argatroban, dabigatran, bivalirudin, desirudin, heparin, lepirudin
Antiplatelet Agents	aspirin, cilostazol, clopidogrel, dipyridamole, prasugrel, ticlopidine
Nonsteroidal Anti-Inflammatory Agents	celecoxib, diclofenac, diflunisal, fenoprofen, ibuprofen, indomethacin, ketoprofen, ketorolac, mefenamic acid, naproxen, oxaprozin, piroxicam, sulindac
Serotonin Reuptake Inhibitors	citalopram, desvenlafaxine, duloxetine, escitalopram, fluoxetine, fluvoxamine, milnacipran, paroxetine, sertraline, venlafaxine, vilazodone

#### **7.4 Antibiotics and Antifungals**

There have been reports of changes in INR in patients taking warfarin and antibiotics or antifungals, but clinical pharmacokinetic studies have not shown consistent effects of these agents on plasma concentrations of warfarin.

Closely monitor INR when starting or stopping any antibiotic or antifungal in patients taking warfarin.

#### **7.5 Botanical (Herbal) Products and Foods**

More frequent INR monitoring should be performed when starting or stopping botanicals.

Few adequate, well-controlled studies evaluating the potential for metabolic and/or pharmacologic interactions between botanicals and JANTOVEN exist. Due to a lack of manufacturing standardization with botanical medicinal preparations, the amount of active ingredients may vary. This could further confound the ability to assess potential interactions and effects on anticoagulation.

Some botanicals may cause bleeding events when taken alone (e.g., garlic and Ginkgo biloba) and may have anticoagulant, antiplatelet, and/or fibrinolytic properties. These effects would be expected to be additive to the anticoagulant effects of JANTOVEN. Conversely, some botanicals may decrease the effects of JANTOVEN (e.g., co-enzyme Q<sub>10</sub>, St. John's wort, ginseng). Some botanicals and foods can interact with JANTOVEN through CYP450 interactions (e.g., echinacea, grapefruit juice, ginkgo, goldenseal, St. John's wort).

The amount of vitamin K in food may affect therapy with JANTOVEN. Advise patients taking JANTOVEN to eat a normal, balanced diet maintaining a consistent amount of vitamin K. Patients taking JANTOVEN should avoid drastic changes in dietary habits, such as eating large amounts of green leafy vegetables.

### **8 USE IN SPECIFIC POPULATIONS**

#### **8.1 Pregnancy**

##### **Risk Summary**

JANTOVEN is contraindicated in women who are pregnant except in pregnant women with mechanical heart valves, who are at high risk of thromboembolism, and for whom the benefits of JANTOVEN may outweigh the risks [see *Warnings and Precautions (5.7)*]. JANTOVEN can cause fetal harm. Exposure to warfarin during the first trimester of pregnancy caused a pattern of congenital malformations in about 5% of exposed offspring. Because these data were not collected in adequate and well-controlled studies, this incidence of major birth defects is not an adequate basis for comparison to the estimated incidences in the control group or the U.S. general population and may not reflect the incidences observed in practice. Consider the benefits and risks of JANTOVEN and possible risks to the fetus when prescribing JANTOVEN to a pregnant woman.

Adverse outcomes in pregnancy occur regardless of the health of the mother or the use of medications. The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

## **Clinical Considerations**

### *Fetal/Neonatal Adverse Reactions*

In humans, warfarin crosses the placenta, and concentrations in fetal plasma approach the maternal values. Exposure to warfarin during the first trimester of pregnancy caused a pattern of congenital malformations in about 5% of exposed offspring. Warfarin embryopathy is characterized by nasal hypoplasia with or without stippled epiphyses (chondrodysplasia punctata) and growth retardation (including low birth weight). Central nervous system and eye abnormalities have also been reported, including dorsal midline dysplasia characterized by agenesis of the corpus callosum, Dandy-Walker malformation, midline cerebellar atrophy, and ventral midline dysplasia characterized by optic atrophy. Mental retardation, blindness, schizencephaly, microcephaly, hydrocephalus, and other adverse pregnancy outcomes have been reported following warfarin exposure during the second and third trimesters of pregnancy [see *Contraindications (4)*].

## **8.2 Lactation**

### **Risk Summary**

Warfarin was not present in human milk from mothers treated with warfarin from a limited published study. Because of the potential for serious adverse reactions, including bleeding in a breastfed infant, consider the developmental and health benefits of breastfeeding along with the mother's clinical need for JANTOVEN and any potential adverse effects on the breastfed infant from JANTOVEN or from the underlying maternal condition before prescribing JANTOVEN to a lactating woman.

### **Clinical Considerations**

Monitor breastfeeding infants for bruising or bleeding.

### **Data**

#### *Human Data*

Based on published data in 15 nursing mothers, warfarin was not detected in human

milk. Among the 15 full-term newborns, 6 nursing infants had documented prothrombin times within the expected range. Prothrombin times were not obtained for the other 9 nursing infants. Effects in premature infants have not been evaluated.

### **8.3 Females and Males of Reproductive Potential**

#### **Pregnancy Testing**

JANTOVEN can cause fetal harm [see *Use in Specific Populations (8.1)*].

Verify the pregnancy status of females of reproductive potential prior to initiating JANTOVEN therapy.

#### **Contraception**

##### *Females*

Advise females of reproductive potential to use effective contraception during treatment and for at least 1 month after the final dose of JANTOVEN.

### **8.4 Pediatric Use**

Adequate and well-controlled studies with warfarin sodium have not been conducted in any pediatric population, and the optimum dosing, safety, and efficacy in pediatric patients is unknown. Pediatric use of warfarin sodium is based on adult data and recommendations, and available limited pediatric data from observational studies and patient registries. Pediatric patients administered JANTOVEN should avoid any activity or sport that may result in traumatic injury.

The developing hemostatic system in infants and children results in a changing physiology of thrombosis and response to anticoagulants. Dosing of warfarin in the pediatric population varies by patient age, with infants generally having the highest, and adolescents having the lowest milligram per kilogram dose requirements to maintain target INRs. Because of changing warfarin requirements due to age, concomitant medications, diet, and existing medical condition, target INR ranges may be difficult to achieve and maintain in pediatric patients, and more frequent INR determinations are recommended. Bleeding rates varied by patient population and clinical care center in pediatric observational studies and patient registries.

Infants and children receiving vitamin K-supplemented nutrition, including infant formulas, may be resistant to warfarin therapy, while human milk-fed infants may be sensitive to warfarin therapy.

### **8.5 Geriatric Use**

Of the total number of patients receiving warfarin sodium in controlled clinical trials for which data were available for analysis, 1885 patients (24.4%) were 65 years and older, while 185 patients (2.4%) were 75 years and older. No overall differences in effectiveness or safety were observed between these patients and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

Patients 60 years or older appear to exhibit greater than expected INR response to the anticoagulant effects of warfarin [see *Clinical Pharmacology (12.3)*]. JANTOVEN is contraindicated in any unsupervised patient with senility. Conduct more frequent monitoring for bleeding with administration of JANTOVEN to elderly patients in any

situation or with any physical condition where added risk of hemorrhage is present. Consider lower initiation and maintenance doses of JANTOVEN in elderly patients [see *Dosage and Administration (2.2, 2.3)*].

## **8.6 Renal Impairment**

Renal clearance is considered to be a minor determinant of anticoagulant response to warfarin. No dosage adjustment is necessary for patients with renal impairment. Instruct patients with renal impairment taking warfarin to monitor their INR more frequently [see *Warnings and Precautions (5.4)*]

## **8.7 Hepatic Impairment**

Hepatic impairment can potentiate the response to warfarin through impaired synthesis of clotting factors and decreased metabolism of warfarin. Conduct more frequent monitoring for bleeding when using JANTOVEN in these patients.

## **10 OVERDOSAGE**

### **10.1 Signs and Symptoms**

Bleeding (e.g., appearance of blood in stools or urine, hematuria, excessive menstrual bleeding, melena, petechiae, excessive bruising or persistent oozing from superficial injuries, unexplained fall in hemoglobin) is a manifestation of excessive anticoagulation.

### **10.2 Treatment**

The treatment of excessive anticoagulation is based on the level of the INR, the presence or absence of bleeding, and clinical circumstances. Reversal of JANTOVEN anticoagulation may be obtained by discontinuing JANTOVEN therapy and, if necessary, by administration of oral or parenteral vitamin K<sub>1</sub>.


The use of vitamin K<sub>1</sub> reduces response to subsequent JANTOVEN therapy and patients may return to a pretreatment thrombotic status following the rapid reversal of a prolonged INR. Resumption of JANTOVEN administration reverses the effect of vitamin K, and a therapeutic INR can again be obtained by careful dosage adjustment. If rapid re-anticoagulation is indicated, heparin may be preferable for initial therapy.

Prothrombin complex concentrate (PCC), fresh frozen plasma, or activated Factor VII treatment may be considered if the requirement to reverse the effects of JANTOVEN is urgent. A risk of hepatitis and other viral diseases is associated with the use of blood products; PCC and activated Factor VII are also associated with an increased risk of thrombosis. Therefore, these preparations should be used only in exceptional or life-threatening bleeding episodes secondary to JANTOVEN overdose.

## **11 DESCRIPTION**

JANTOVEN (Warfarin Sodium Tablets, USP) is an anticoagulant that acts by inhibiting vitamin K-dependent coagulation factors. The chemical name of warfarin sodium is 3-( $\alpha$ -acetylbenzyl)-4-hydroxycoumarin sodium salt, which is a racemic mixture of the *R*- and *S*-enantiomers. Crystalline warfarin sodium is an isopropanol clathrate. Its empirical

formula is  $C_{19}H_{15}NaO_4$ , and its structural formula is represented by the following:



Crystalline warfarin sodium occurs as a white, odorless, crystalline powder that is discolored by light. It is very soluble in water, freely soluble in alcohol, and very slightly soluble in chloroform and ether.

JANTOVEN tablets for oral use contain: 1 mg, 2 mg, 2½ mg, 3 mg, 4 mg, 5 mg, 6 mg, 7½ mg or 10 mg of warfarin sodium, USP. They also contain:

---

All strengths:	Lactose monohydrate, magnesium stearate, povidone, and pregelatinized starch (corn).
1 mg:	FD&C Red #40 Aluminum Lake
2 mg:	FD&C Blue #2 Aluminum Lake and FD&C Red #40 Aluminum Lake
2½ mg:	D&C Yellow #10 Aluminum Lake and FD&C Blue #1 Aluminum Lake
3 mg:	Brown #75 Synthetic Brown Iron Oxide
4 mg:	FD&C Blue #1 Aluminum Lake
5 mg:	FD&C Yellow #6 Aluminum Lake
6 mg:	Yellow #10 Synthetic Yellow Iron Oxide, Black #85 Synthetic Black Iron Oxide and FD&C Blue #1 Aluminum Lake
7½ mg:	D&C Yellow #10 Aluminum Lake and FD&C Yellow #6 Aluminum Lake
10 mg:	Dye Free

---

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Warfarin acts by inhibiting the synthesis of vitamin K-dependent clotting factors, which include Factors II, VII, IX, and X, and the anticoagulant proteins C and S. Vitamin K is an essential cofactor for the post ribosomal synthesis of the vitamin K-dependent clotting factors. Vitamin K promotes the biosynthesis of  $\gamma$ -carboxyglutamic acid residues in the proteins that are essential for biological activity. Warfarin is thought to interfere with clotting factor synthesis by inhibition of the C1 subunit of vitamin K epoxide reductase (VKORC1) enzyme complex, thereby reducing the regeneration of vitamin K<sub>1</sub> epoxide

[see *Clinical Pharmacology (12.5)*].

## 12.2 Pharmacodynamics

An anticoagulation effect generally occurs within 24 hours after warfarin administration. However, peak anticoagulant effect may be delayed 72 to 96 hours. The duration of action of a single dose of racemic warfarin is 2 to 5 days. The effects of JANTOVEN may become more pronounced as effects of daily maintenance doses overlap. This is consistent with the half-lives of the affected vitamin K-dependent clotting factors and anticoagulation proteins: Factor II – 60 hours, VII – 4 to 6 hours, IX – 24 hours, X – 48 to 72 hours, and proteins C and S are approximately 8 hours and 30 hours, respectively.

## 12.3 Pharmacokinetics

JANTOVEN is a racemic mixture of the *R*- and *S*-enantiomers of warfarin. The *S*-enantiomer exhibits 2 to 5 times more anticoagulant activity than the *R*-enantiomer in humans, but generally has a more rapid clearance.

### Absorption

Warfarin is essentially completely absorbed after oral administration, with peak concentration generally attained within the first 4 hours.

### Distribution

Warfarin shows a volume of distribution of about 0.14 L/kg. Approximately 99% of the drug is bound to plasma proteins.

### Metabolism

The elimination of warfarin is almost entirely by metabolism. Warfarin is stereoselectively metabolized by hepatic cytochrome P-450 (CYP450) microsomal enzymes to inactive hydroxylated metabolites (predominant route) and by reductases to reduced metabolites (warfarin alcohols) with minimal anticoagulant activity. Identified metabolites of warfarin include dehydrowarfarin, two diastereoisomer alcohols, and 4<sup>l</sup>-, 6-, 7-, 8-, and 10-hydroxywarfarin. The CYP450 isozymes involved in the metabolism of warfarin include CYP2C9, 2C19, 2C8, 2C18, 1A2, and 3A4. CYP2C9, a polymorphic enzyme, is likely to be the principal form of human liver CYP450 that modulates the *in vivo* anticoagulant activity of warfarin. Patients with one or more variant CYP2C9 alleles have decreased *S*-warfarin clearance [see *Clinical Pharmacology (12.5)*].

### Excretion

The terminal half-life of warfarin after a single dose is approximately 1 week; however, the effective half-life ranges from 20 to 60 hours, with a mean of about 40 hours. The clearance of *R*-warfarin is generally half that of *S*-warfarin, thus as the volumes of distribution are similar, the half-life of *R*-warfarin is longer than that of *S*-warfarin. The half-life of *R*-warfarin ranges from 37 to 89 hours, while that of *S*-warfarin ranges from 21 to 43 hours. Studies with radiolabeled drug have demonstrated that up to 92% of the orally administered dose is recovered in urine. Very little warfarin is excreted unchanged in urine. Urinary excretion is in the form of metabolites.

### Geriatric Patients

Patients 60 years or older appear to exhibit greater than expected INR response to the anticoagulant effects of warfarin. The cause of the increased sensitivity to the

anticoagulant effects of warfarin in this age group is unknown but may be due to a combination of pharmacokinetic and pharmacodynamic factors. Limited information suggests there is no difference in the clearance of S-warfarin; however, there may be a slight decrease in the clearance of R-warfarin in the elderly as compared to the young. Therefore, as patient age increases, a lower dose of warfarin is usually required to produce a therapeutic level of anticoagulation [see *Dosage and Administration* (2.3, 2.4)].

## **Asian Patients**

Asian patients may require lower initiation and maintenance doses of warfarin. A non-controlled study of 151 Chinese outpatients stabilized on warfarin for various indications reported a mean daily warfarin requirement of  $3.3 \pm 1.4$  mg to achieve an INR of 2 to 2.5. Patient age was the most important determinant of warfarin requirement in these patients, with a progressively lower warfarin requirement with increasing age.

## **12.5 Pharmacogenomics**

### **CYP2C9 and VKORC1 Polymorphisms**

The S-enantiomer of warfarin is mainly metabolized to 7-hydroxywarfarin by CYP2C9, a polymorphic enzyme. The variant alleles, CYP2C9\*2 and CYP2C9\*3, result in decreased *in vitro* CYP2C9 enzymatic 7-hydroxylation of S-warfarin. The frequencies of these alleles in Caucasians are approximately 11% and 7% for CYP2C9\*2 and CYP2C9\*3, respectively.

Other CYP2C9 alleles associated with reduced enzymatic activity occur at lower frequencies, including \*5, \*6, and \*11 alleles in populations of African ancestry and \*5, \*9, and \*11 alleles in Caucasians.

Warfarin reduces the regeneration of vitamin K from vitamin K epoxide in the vitamin K cycle through inhibition of VKOR, a multiprotein enzyme complex. Certain single nucleotide polymorphisms in the VKORC1 gene (e.g., -1639G>A) have been associated with variable warfarin dose requirements. VKORC1 and CYP2C9 gene variants generally explain the largest proportion of known variability in warfarin dose requirements.

CYP2C9 and VKORC1 genotype information, when available, can assist in selection of the initial dose of warfarin [see *Dosage and Administration* (2.3)].

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

Carcinogenicity, mutagenicity, or fertility studies have not been performed with warfarin.

## **14 CLINICAL STUDIES**

### **14.1 Atrial Fibrillation**

In five prospective, randomized, controlled clinical trials involving 3711 patients with non-rheumatic AF, warfarin significantly reduced the risk of systemic thromboembolism including stroke (see Table 4). The risk reduction ranged from 60% to 86% in all except one trial (CAFA: 45%), which was stopped early due to published positive results from two of these trials. The incidence of major bleeding in these trials ranged from 0.6% to

2.7% (see Table 4).

**Table 4: Clinical Studies of Warfarin in Non-Rheumatic AF Patients \***

Study	N Warfarin- Treated Patients	N Control Patients	PT		% Risk Reduction	p- value	% Major Bleeding	
			Ratio	INR			Warfarin- Treated Patients	Control Patients
AFASAK	335	336	1.5- 2.0	2.8- 4.2	60	0.027	0.6	0.0
SPAF	210	211	1.3- 1.8	2.0- 4.5	67	0.01	1.9	1.9
BAATAF	212	208	1.2- 1.5	1.5- 2.7	86	<0.05	0.9	0.5
CAFA	187	191	1.3- 1.6	2.0- 3.0	45	0.25	2.7	0.5
SPINAF	260	265	1.2- 1.5	1.4- 2.8	79	0.001	2.3	1.5

\* All study results of warfarin vs. control are based on intention-to-treat analysis and include ischemic stroke and systemic thromboembolism, excluding hemorrhagic stroke and transient ischemic attacks.

Trials in patients with both AF and mitral stenosis suggest a benefit from anticoagulation with warfarin sodium [see *Dosage and Administration (2.2)*].

## 14.2 Mechanical and Bioprosthetic Heart Valves

In a prospective, randomized, open-label, positive-controlled study in 254 patients with mechanical prosthetic heart valves, the thromboembolic-free interval was found to be significantly greater in patients treated with warfarin alone compared with dipyridamole/aspirin-treated patients ( $p < 0.005$ ) and pentoxifylline/aspirin-treated patients ( $p < 0.05$ ). The results of this study are presented in Table 5.

**Table 5: Prospective, Randomized, Open-Label, Positive-Controlled Clinical Study of Warfarin in Patients with Mechanical Prosthetic Heart Valves**

Event	Patients Treated With		
	Warfarin	Dipyridamole/Aspirin	Pentoxifylline/Aspirin
Thromboembolism	2.2/100 py	8.6/100 py	7.9/100 py
Major Bleeding	2.5/100 py	0.0/100 py	0.9/100 py

py=patient years

In a prospective, open-label, clinical study comparing moderate (INR 2.65) versus high intensity (INR 9.0) warfarin therapies in 258 patients with mechanical prosthetic heart valves, thromboembolism occurred with similar frequency in the two groups (4.0 and 3.7 events per 100 patient years, respectively). Major bleeding was more common in the high intensity group. The results of this study are presented in Table 6.

**Table 6: Prospective, Open-Label Clinical Study of Warfarin in Patients with Mechanical Prosthetic Heart Valves**

<b>Event</b>	<b>Moderate Warfarin Therapy INR 2.65</b>	<b>High Intensity Warfarin Therapy INR 9.0</b>
Thromboembolism	4.0/100 py	3.7/100 py
Major Bleeding	0.95/100 py	2.1/100 py

py=patient years

In a randomized trial in 210 patients comparing two intensities of warfarin therapy (INR 2.0 to 2.25 vs. INR 2.5 to 4.0) for a three-month period following tissue heart valve replacement, thromboembolism occurred with similar frequency in the two groups (major embolic events 2.0% vs. 1.9%, respectively, and minor embolic events 10.8% vs. 10.2%, respectively). Major hemorrhages occurred in 4.6% of patients in the higher intensity INR group compared to zero in the lower intensity INR group.

### 14.3 Myocardial Infarction

WARIS (The Warfarin Re-Infarction Study) was a double-blind, randomized study of 1214 patients 2 to 4 weeks post-infarction treated with warfarin to a target INR of 2.8 to 4.8. The primary endpoint was a composite of total mortality and recurrent infarction. A secondary endpoint of cerebrovascular events was assessed. Mean follow-up of the patients was 37 months. The results for each endpoint separately, including an analysis of vascular death, are provided in Table 7.

**Table 7: WARIS - Endpoint Analysis of Separate Events**

<b>Event</b>	<b>Warfarin (N=607)</b>	<b>Placebo (N=607)</b>	<b>RR (95% CI)</b>	<b>% Risk Reduction ( p-value)</b>
Total Patient Years of Follow-up	2018	1944		
Total Mortality	94 (4.7/100 py)	123 (6.3/100 py)	0.76 (0.60, 0.97)	24 (p=0.030)
Vascular Death	82 (4.1/100 py)	105 (5.4/100 py)	0.78 (0.60, 1.02)	22 (p=0.068)
Recurrent MI	82 (4.1/100 py)	124 (6.4/100 py)	0.66 (0.51, 0.85)	34 (p=0.001)
Cerebrovascular Event	20 (1.0/100 py)	44 (2.3/100 py)	0.46 (0.28, 0.75)	54 (p=0.002)

RR=Relative risk; Risk reduction=(1 - RR); CI=Confidence interval; MI=Myocardial infarction; py=patient years

WARIS II (The Warfarin, Aspirin, Re-Infarction Study) was an open-label, randomized study of 3630 patients hospitalized for acute myocardial infarction treated with warfarin to a target INR 2.8 to 4.2, aspirin 160 mg per day, or warfarin to a target INR 2.0 to 2.5 plus aspirin 75 mg per day prior to hospital discharge. The primary endpoint was a composite of death, nonfatal reinfarction, or thromboembolic stroke. The mean duration

of observation was approximately 4 years. The results for WARIS II are provided in Table 8.

**Table 8: WARIS II - Distribution of Events According to Treatment Group**

Event	Aspirin plus Warfarin			Rate Ratio (95% CI)	p-value
	Aspirin (N=1206)	Warfarin (N=1216)	Warfarin (N=1208)		
	No. of Events				
Major Bleeding *	8	33	28	3.35 †(ND) 4.00 ‡(ND)	ND ND
Minor Bleeding §	39	103	133	3.21 †(ND) 2.55 ‡(ND)	ND ND
Composite Endpoints ¶	241	203	181	0.81 (0.69-0.95) † 0.71 (0.60-0.83) ‡	0.03 0.001
Reinfarction	117	90	69	0.56 (0.41-0.78) † 0.74 (0.55-0.98) ‡	<0.001 0.03
Thromboembolic Stroke	32	17	17	0.52 (0.28-0.98) † ‡	0.03 0.03
Death	92	96	95		0.82

CI=confidence interval

ND=not determined

\* Major bleeding episodes were defined as nonfatal cerebral hemorrhage or bleeding necessitating surgical intervention or blood transfusion.

† The rate ratio is for aspirin plus warfarin as compared with aspirin.

‡ The rate ratio is for warfarin as compared with aspirin.

§ Minor bleeding episodes were defined as non-cerebral hemorrhage not necessitating surgical intervention or blood transfusion.

¶ Includes death, nonfatal reinfarction, and thromboembolic cerebral stroke.

There were approximately four times as many major bleeding episodes in the two groups receiving warfarin than in the group receiving aspirin alone. Major bleeding episodes were not more frequent among patients receiving aspirin plus warfarin than among those receiving warfarin alone, but the incidence of minor bleeding episodes was higher in the combined therapy group.

## 15 REFERENCES

OSHA Hazardous Drugs. OSHA. <http://www.osha.gov/SLTC/hazardousdrugs/index.html>.

## 16 HOW SUPPLIED/STORAGE AND HANDLING

## Tablets

JANTOVEN tablets are single scored, compressed tablets with one side scored and debossed with WRF above the score and 1, 2, 2½, 3, 4, 5, 6, 7½, or 10 below the score and with 832 debossed on the opposite side. JANTOVEN is available in bottles and unit dose blister packages with potencies and colors as follows:

1 mg – Compressed tablet, pink, round; in bottles of 100 (NDC 0832-1211-00) and 1000 (NDC 0832-1211-10) and in unit dose cartons of 100 tablets (10 cards containing 10 tablets each) (NDC 0832-1211-01).

2 mg – Compressed tablet, lavender, round; in bottles of 100 (NDC 0832-1212-00) and 1000 (NDC 0832-1212-10) and in unit dose cartons of 100 tablets (10 cards containing 10 tablets each) (NDC 0832-1212-01).

2½ mg – Compressed tablet, green, round; in bottles of 100 (NDC 0832-1213-00) and 1000 (NDC 0832-1213-10) and in unit dose cartons of 100 tablets (10 cards containing 10 tablets each) (NDC 0832-1213-01).

3 mg – Compressed tablet, tan, round; in bottles of 100 (NDC 0832-1214-00) and 1000 (NDC 0832-1214-10) and in unit dose cartons of 100 tablets (10 cards containing 10 tablets each) (NDC 0832-1214-01).

4 mg – Compressed tablet, blue, round; in bottles of 100 (NDC 0832-1215-00) and 1000 (NDC 0832-1215-10) and in unit dose cartons of 100 tablets (10 cards containing 10 tablets each) (NDC 0832-1215-01).

5 mg – Compressed tablet, peach, round; in bottles of 100 (NDC 0832-1216-00) and 1000 (NDC 0832-1216-10) and in unit dose cartons of 100 tablets (10 cards containing 10 tablets each) (NDC 0832-1216-01).

6 mg – Compressed tablet, teal, round; in bottles of 100 (NDC 0832-1217-00) and 1000 (NDC 0832-1217-10) and in unit dose cartons of 100 tablets (10 cards containing 10 tablets each) (NDC 0832-1217-01).

7½ mg – Compressed tablet, yellow, round; in bottles of 100 (NDC 0832-1218-00) and 500 (NDC 0832-1218-50) and in unit dose cartons of 100 tablets (10 cards containing 10 tablets each) (NDC 0832-1218-01).

10 mg – Compressed tablet, white (dye-free), round; in bottles of 100 (NDC 0832-1219-00) and 500 (NDC 0832-1219-50) and in unit dose cartons of 100 tablets (10 cards containing 10 tablets each) (NDC 0832-1219-01).

Store at 20° to 25°C (68° to 77°F). Excursions permitted to 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature.] Protect from light and moisture. Dispense in a tight, light-resistant container with a child-resistant closure.

## Special Handling

Procedures for proper handling and disposal of potentially hazardous drugs should be considered. Guidelines on this subject have been published [see *References (15)*].

Pharmacy and clinical personnel who are pregnant should avoid exposure to crushed or broken tablets [see *Use in Specific Populations (8.1)*].

## 17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide).

## **Instructions for Patients**

Advise patients to:

- Strictly adhere to the prescribed dosage schedule [see *Dosage and Administration (2.1)*].
- If the prescribed dose of JANTOVEN is missed, take the dose as soon as possible on the same day but do not take a double dose of JANTOVEN the next day to make up for missed doses [see *Dosage and Administration (2.6)*].
- Obtain prothrombin time tests and make regular visits to their physician or clinic to monitor therapy [see *Dosage and Administration (2.1)*].
- Be aware that if therapy with JANTOVEN is discontinued, the anticoagulant effects of JANTOVEN may persist for about 2 to 5 days [see *Clinical Pharmacology (12.2)*].
- Avoid any activity or sport that may result in traumatic injury [see *Use in Specific Populations (8.4)*]. And to tell their physician if they fall often as this may increase their risk for complications.
- Eat a normal, balanced diet to maintain a consistent intake of vitamin K. Avoid drastic changes in dietary habits, such as eating large amounts of leafy, green vegetables [see *Drug Interactions (7.5)*].
- Contact their physician to report any serious illness, such as severe diarrhea, infection, or fever [see *Warnings and Precautions (5)* and *Adverse Reactions (6)*].
- Immediately contact their physician when experiencing pain and discoloration of the skin (a purple bruise like rash) mostly on areas of the body with a high fat content, such as breasts, thighs, buttocks, hips and abdomen [see *Warnings and Precautions (5.2)*].
- Immediately contact their physician when experiencing any unusual symptom or pain since JANTOVEN may cause small cholesterol or athero emboli. On feet it may appear as a sudden cool, painful, purple discoloration of toe(s) or forefoot [see *Warnings and Precautions (5.5)*].
- Immediately contact their physician when taking JANTOVEN after any heparin formulation therapy and experiencing bloody or black stools or appearance of bruises, or bleeding [see *Warnings and Precautions (5.6)*].
- To tell all of their healthcare professionals and dentists that they are taking JANTOVEN. This should be done before they have any surgery or dental procedure [see *Dosage and Administration (2.7)*].
- Carry identification stating that they are taking JANTOVEN.

## **Bleeding Risks**

Advise patients to:

- Notify their physician immediately if any unusual bleeding or symptoms occur. Signs and symptoms of bleeding include: pain, swelling or discomfort, prolonged bleeding from cuts, increased menstrual flow or vaginal bleeding, nosebleeds, bleeding of gums from brushing, unusual bleeding or bruising, red or dark brown urine, red or tar black stools, headache, dizziness, or weakness [see *Box Warning and Warnings and Precautions (5.1)*].

## **Concomitant Medications and Botanicals (Herbals)**

Advise patients to:

- Not take or discontinue any other drug, including salicylates (e.g., aspirin and topical analgesics), other over-the-counter drugs, and botanical (herbal) products except on advice of your physician [see *Drug Interactions (7)*].

## **Pregnancy and Nursing**

Advise patients to:

- Notify their physician if they are pregnant or planning to become pregnant or considering breast feeding [ see *Use in Specific Populations (8.1, 8.2, 8.3)*].
- Avoid JANTOVEN during pregnancy except in pregnant women with mechanical heart valves, who are at risk of thromboembolism [ see *Contraindications (4)*]. Use effective measures to avoid pregnancy while taking JANTOVEN. This is very important because their unborn baby could be seriously harmed if they take JANTOVEN while they are pregnant [ see *Use in Specific Populations (8.1, 8.3)*].

JANTOVEN is a registered trademark of Upsher-Smith Laboratories, LLC.

Manufactured by

**UPSHER-SMITH LABORATORIES, LLC**

Maple Grove, MN 55369

Revised 0917

<p style="text-align: center;"><b>MEDICATION GUIDE</b> <b>JANTOVEN<sup>®</sup> (JAN-to-ven) Tablets</b> <b>(Warfarin Sodium Tablets, USP)</b></p>
---

**What is the most important information I should know about JANTOVEN?**

JANTOVEN can cause bleeding which can be serious and sometimes lead to death. This is because JANTOVEN is a blood thinner medicine that lowers the chance of blood clots forming in your body.

- You may have a higher risk of bleeding if you take JANTOVEN and:
  - are 65 years of age or older
  - have a history of stomach or intestinal bleeding
  - have high blood pressure (hypertension)
  - have a history of stroke, or "mini-stroke" (transient ischemic attack or TIA)
  - have serious heart disease
  - have a low blood count or cancer
  - have had trauma, such as an accident or surgery
  - have kidney problems
  - take other medicines that increase your risk of bleeding, including:
    - a medicine that contains heparin
    - other medicines to prevent or treat blood clots
    - nonsteroidal anti-inflammatory drugs (NSAIDs)
  - take warfarin sodium for a long time. Warfarin sodium is the active ingredient in JANTOVEN.

**Tell your healthcare provider if you take any of these medicines. Ask your healthcare provider if you are not sure if your medicine is one listed above.**

Many other medicines can interact with JANTOVEN and affect the dose you need or increase JANTOVEN side effects. Do not change or stop any of your medicines or start

any new medicines before you talk to your healthcare provider.

**Do not take other medicines that contain warfarin sodium while taking JANTOVEN.**

- **Get your regular blood test to check for your response to JANTOVEN.** This blood test is called an INR test. The INR test checks to see how fast your blood clots. Your healthcare provider will decide what INR numbers are best for you. Your dose of JANTOVEN will be adjusted to keep your INR in a target range for you.
- **Call your healthcare provider right away if you get any of the following signs or symptoms of bleeding problems:**
  - pain, swelling, or discomfort
  - headaches, dizziness, or weakness
  - unusual bruising (bruises that develop without known cause or grow in size)
  - nosebleeds
  - bleeding gums
  - bleeding from cuts takes a long time to stop
  - menstrual bleeding or vaginal bleeding that is heavier than normal
  - pink or brown urine
  - red or black stools
  - coughing up blood
  - vomiting blood or material that looks like coffee grounds
- **Some foods and beverages can interact with JANTOVEN and affect your treatment and dose.**
  - Eat a normal, balanced diet. Talk to your healthcare provider before you make any diet changes. Do not eat large amounts of leafy, green vegetables. Leafy, green vegetables contain vitamin K. Certain vegetable oils also contain large amounts of vitamin K. Too much vitamin K can lower the effect of JANTOVEN.
- Always tell all of your healthcare providers that you take JANTOVEN.
- Wear or carry information that you take JANTOVEN.

**See " What are the possible side effects of JANTOVEN?" for more information about side effects.**

**What is JANTOVEN?**

JANTOVEN is prescription medicine used to treat blood clots and to lower the chance of blood clots forming in your body. Blood clots can cause a stroke, heart attack, or other serious conditions if they form in the legs or lungs.

**Who should not take JANTOVEN?**

**Do not take JANTOVEN if:**

- **your risk of having bleeding problems is higher than the possible benefit of treatment.** Your healthcare provider will decide if JANTOVEN is right for you.
- **you are pregnant unless you have a mechanical heart valve.** JANTOVEN may cause birth defects, miscarriage, or death of your unborn baby.
- **you are allergic to warfarin or any of the other ingredients in JANTOVEN. See the end of this leaflet for a complete list of ingredients in JANTOVEN.**

**Before taking JANTOVEN, tell your healthcare provider about all of your medical conditions, including if you:**

- have bleeding problems
- fall often

- have liver problems
- have kidney problems or are undergoing dialysis
- have high blood pressure
- have a heart problem called congestive heart failure
- have diabetes
- plan to have any surgery or a dental procedure
- are pregnant or plan to become pregnant. See "**Who should not take JANTOVEN?**"
  - Your healthcare provider will do a pregnancy test before you start treatment with JANTOVEN. Females who can become pregnant should use effective birth control during treatment, and for at least 1 month after the last dose of JANTOVEN.
- are breastfeeding. You and your healthcare provider should decide if you will take JANTOVEN and breastfeed. Check your baby for bruising or bleeding if you take JANTOVEN and breastfeed.

Tell all of your healthcare providers and dentists that you are taking JANTOVEN. They should talk to the healthcare provider who prescribed JANTOVEN for you before you have **any** surgery or dental procedure. Your JANTOVEN may need to be stopped for a short time or you may need your dose adjusted.

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Some of your other medicines may affect the way JANTOVEN works. Certain medicines may increase your risk of bleeding.

See "**What is the most important information I should know about JANTOVEN?**"

#### **How should I take JANTOVEN?**

- **Take JANTOVEN exactly as prescribed.** Your healthcare provider will adjust your dose from time to time depending on your response to JANTOVEN.
- **You must have regular blood tests and visits with your healthcare provider to monitor your condition.**
- **If you miss a dose of JANTOVEN, call your healthcare provider.** Take the dose as soon as possible on the same day. **Do not** take a double dose of JANTOVEN the next day to make up for a missed dose.
- Call your healthcare provider right away if you:
  - take too much JANTOVEN
  - are sick with diarrhea, an infection, or have a fever
  - fall or injure yourself, especially if you hit your head. Your healthcare provider may need to check you.

#### **What should I avoid while taking JANTOVEN?**

- Do not do any activity or sport that may cause a serious injury.

#### **What are the possible side effects of JANTOVEN?**

**JANTOVEN may cause serious side effects, including:**

- See "**What is the most important information I should know about JANTOVEN?**"
  - **Death of skin tissue (skin necrosis or gangrene).** This can happen soon after starting JANTOVEN. It happens because blood clots form and block blood flow to an area of your body. Call your healthcare provider right away if you have

pain, color, or temperature change to any area of your body. You may need medical care right away to prevent death or loss (amputation) of your affected body part.

- **Kidney problems.** Kidney injury may happen in people who take JANTOVEN. Tell your healthcare provider right away if you develop blood in your urine. Your healthcare provider may do tests more often during treatment with JANTOVEN to check for bleeding if you already have kidney problems.
- **"Purple toes syndrome."** Call your healthcare provider right away if you have pain in your toes and they look purple in color or dark in color.

These are not all of the side effects of JANTOVEN. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

### **How should I store JANTOVEN?**

- Store JANTOVEN at 20° to 25°C (68° to 77°F). Excursions permitted to 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature.]
- Keep JANTOVEN in a tightly closed container, and keep JANTOVEN out of the light and moisture.
- Follow your healthcare provider or pharmacist instructions about the right way to throw away outdated or unused JANTOVEN.
- Females who are pregnant should not handle crushed or broken JANTOVEN tablets.

### **Keep JANTOVEN and all medicines out of the reach of children.**

### **General information about the safe and effective use of JANTOVEN.**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use JANTOVEN for a condition for which it was not prescribed. Do not give JANTOVEN to other people, even if they have the same symptoms that you have. It may harm them. You can ask your healthcare provider or pharmacist for information about JANTOVEN that is written for health professionals.

### **What are the ingredients in JANTOVEN Tablets?**

Active ingredient: Warfarin Sodium, USP

Inactive ingredients: Lactose monohydrate, magnesium stearate, povidone, and pregelatinized starch (corn). The following tablets contain:

1 mg: FD&C Red #40 Aluminum Lake

2 mg: FD&C Blue #2 Aluminum Lake and FD&C Red #40 Aluminum Lake

2½ mg: D&C Yellow #10 Aluminum Lake and FD&C Blue #1 Aluminum Lake

3 mg: Brown #75 Synthetic Brown Iron Oxide

4 mg: FD&C Blue #1 Aluminum Lake

5 mg: FD&C Yellow #6 Aluminum Lake

6 mg: Yellow #10 Synthetic Yellow Iron Oxide, Black #85 Synthetic Black Iron Oxide and FD&C Blue #1 Aluminum Lake

7½ mg: D&C Yellow #10 Aluminum Lake and FD&C Yellow #6 Aluminum Lake

10 mg: Dye Free

For Medication Guides, please visit [www.upsher-smith.com](http://www.upsher-smith.com) or call 1-888-650-3789.

The brands listed (other than JANTOVEN<sup>®</sup>) are registered trademarks of their respective owners and are not trademarks of Upsher-Smith Laboratories, LLC.

JANTOVEN is a registered trademark of Upsher-Smith Laboratories, LLC.

Manufactured by  
**UPSHER-SMITH LABORATORIES, LLC**  
Maple Grove, MN 55369

This Medication Guide has been approved by the U.S. Food and Drug Administration.  
Revised 0917

## PRINCIPAL DISPLAY PANEL - 1 mg Tablet Bottle Label

**NDC 0832-1211-00**

**JANTOVEN<sup>®</sup>**

**Warfarin Sodium  
Tablets, USP**

**1 mg**

**PHARMACIST: Dispense the Medication  
Guide provided separately to each patient.**

### **HIGHLY POTENT ANTICOAGULANT WARNING:**

Serious bleeding results from overdosage. Do not use or dispense before reading directions and warnings in accompanying product information.

100 Tablets

**Rx only**

**UPSHER-SMITH**

**NDC 0832-1211-00**  
**JANTOVEN<sup>®</sup>**  
Warfarin Sodium  
Tablets, USP

**1 mg**

**PHARMACIST: Dispense the Medication  
Guide provided separately to each patient.**

**HIGHLY POTENT ANTICOAGULANT WARNING:**  
Serious bleeding results from overdosage. Do not use or dispense before reading directions and warnings in accompanying product information.

100 Tablets      **Rx only**

**UPSHER-SMITH**

Each tablet contains: Warfarin Sodium 1 mg  
Usual Adult Dosage: See package insert for full prescribing information.  
Note to authorized dispenser: Obtain a Medication Guide by visiting [www.upsher-smith.com](http://www.upsher-smith.com) or by calling 1-888-650-3789.

Store at 20° to 25°C (68° to 77°F). Excursions permitted to 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature.] Keep tightly closed. Protect from light and moisture. Dispense in a tight, light-resistant container with a child-resistant closure.

**Keep out of reach of children.**  
SEALED FOR YOUR PROTECTION  
Manufactured by  
**UPSHER-SMITH LABORATORIES, LLC**  
Maple Grove, MN 55369  
© 2017 Upsher-Smith Laboratories, LLC  
112608-01 R0617

WRF

0832-1211-00 7

## PRINCIPAL DISPLAY PANEL - 2 mg Tablet Bottle Label

**NDC 0832-1212-00**

**JANTOVEN<sup>®</sup>**

**Warfarin Sodium  
Tablets, USP**

**2 mg**

**PHARMACIST: Dispense the Medication Guide provided separately to each patient.**

**HIGHLY POTENT ANTICOAGULANT WARNING:**  
Serious bleeding results from overdosage. Do not use or dispense before reading directions and warnings in accompanying product information.

100 Tablets

**Rx only**

**UPSHER-SMITH**

NDC 0832-1212-00  
**JANTOVEN**<sup>®</sup>  
Warfarin Sodium  
Tablets, USP

**2 mg**

PHARMACIST: Dispense the Medication Guide provided separately to each patient.

**HIGHLY POTENT ANTICOAGULANT WARNING:**  
Serious bleeding results from overdosage. Do not use or dispense before reading directions and warnings in accompanying product information.

100 Tablets **Rx only**

**UPSHER-SMITH**

Each tablet contains: Warfarin Sodium 2 mg  
Usual Adult Dosage: See package insert for full prescribing information.  
Note to authorized dispenser: Obtain a Medication Guide by visiting [www.upshe-smith.com](http://www.upshe-smith.com) or by calling 1-888-650-3789.  
Store at 20° to 25°C (68° to 77°F). Excursions permitted to 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature.] Keep tightly closed. Protect from light and moisture. Dispense in a tight, light-resistant container with a child-resistant closure.  
Keep out of reach of children.  
SEALED FOR YOUR PROTECTION  
Manufactured by  
**UPSHER-SMITH LABORATORIES, LLC**  
Maple Grove, MN 55369  
© 2017 Upsher-Smith Laboratories, LLC  
112610-01 R0617

0832-1212-004

**PRINCIPAL DISPLAY PANEL - 2½ mg Tablet Bottle Label**

**NDC 0832-1213-00**

**JANTOVEN**<sup>®</sup>

**Warfarin Sodium  
Tablets, USP**

**2½ mg**

**PHARMACIST: Dispense the Medication Guide provided separately to each patient.**

**HIGHLY POTENT ANTICOAGULANT WARNING:**  
Serious bleeding results from overdosage. Do not use or dispense before reading directions and warnings in accompanying product information.

100 Tablets

**Rx only**

**UPSHER-SMITH**

NDC 0832-1213-00

**JANTOVEN<sup>®</sup>**

Warfarin Sodium  
Tablets, USP

**2½ mg**



**PHARMACIST: Dispense the Medication Guide provided separately to each patient.**

**HIGHLY POTENT ANTICOAGULANT WARNING:**  
Serious bleeding results from overdosage. Do not use or dispense before reading directions and warnings in accompanying product information.

100 Tablets

**Rx only**

**UPSHER-SMITH**

Each tablet contains: Warfarin Sodium 2½ mg

Usual Adult Dosage: See package insert for full prescribing information.

Note to authorized dispenser: Obtain a Medication Guide by visiting [www.upshe-smith.com](http://www.upshe-smith.com) or by calling 1-888-650-3789.

Store at 20° to 25°C (68° to 77°F). Excursions permitted to 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature.] Keep tightly closed. Protect from light and moisture. Dispense in a tight, light-resistant container with a child-resistant closure.

Keep out of reach of children.

SEALED FOR YOUR PROTECTION

Manufactured by  
**UPSHER-SMITH LABORATORIES, LLC**  
Maple Grove, MN 55369

© 2017 Upsher-Smith Laboratories, LLC

112611-01 R0617



## PRINCIPAL DISPLAY PANEL - 3 mg Tablet Bottle Label

NDC 0832-1214-00

**JANTOVEN<sup>®</sup>**

Warfarin Sodium  
Tablets, USP

**3 mg**

**PHARMACIST: Dispense the Medication Guide provided separately to each patient.**

**HIGHLY POTENT ANTICOAGULANT WARNING:**

Serious bleeding results from overdosage. Do not use or dispense before reading directions and warnings in accompanying product information.

100 Tablets

**Rx only**

**UPSHER-SMITH**

NDC 0832-1214-00

**JANTOVEN<sup>®</sup>**

Warfarin Sodium  
Tablets, USP

**3 mg**

**PHARMACIST:** Dispense the Medication Guide provided separately to each patient.

**HIGHLY POTENT ANTICOAGULANT WARNING:**  
Serious bleeding results from overdosage. Do not use or dispense before reading directions and warnings in accompanying product information.

100 Tablets

**Rx only**

**UPSHER-SMITH**



Each tablet contains: Warfarin Sodium 3 mg

**Usual Adult Dosage:** See package insert for full prescribing information.

**Note to authorized dispenser:** Obtain a Medication Guide by visiting [www.upshe-smith.com](http://www.upshe-smith.com) or by calling 1-888-650-3789.

Store at 20° to 25°C (68° to 77°F). Excursions permitted to 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature.] Keep tightly closed. Protect from light and moisture. Dispense in a tight, light-resistant container with a child-resistant closure.

**Keep out of reach of children.**

SEALED FOR YOUR PROTECTION

Manufactured by  
**UPSHER-SMITH LABORATORIES, LLC**  
Maple Grove, MN 55369

© 2017 Upsher-Smith Laboratories, LLC  
112612-01 R0617



## PRINCIPAL DISPLAY PANEL - 4 mg Tablet Bottle Label

NDC 0832-1215-00

**JANTOVEN<sup>®</sup>**

Warfarin Sodium  
Tablets, USP

**4 mg**

**PHARMACIST:** Dispense the Medication Guide provided separately to each patient.

**HIGHLY POTENT ANTICOAGULANT WARNING:**  
Serious bleeding results from overdosage. Do not use or dispense before reading directions and warnings in accompanying product information.

100 Tablets

**Rx only**

**UPSHER-SMITH**

NDC 0832-1215-00

**JANTOVEN**<sup>®</sup>

Warfarin Sodium  
Tablets, USP

**4 mg**

**PHARMACIST: Dispense the Medication  
Guide provided separately to each patient.**

**HIGHLY POTENT ANTICOAGULANT WARNING:**  
Serious bleeding results from overdosage. Do not  
use or dispense before reading directions and  
warnings in accompanying product information.

100 Tablets

**Rx only**

**UPSHER-SMITH**



Each tablet contains: Warfarin Sodium 4 mg

**Usual Adult Dosage:** See package insert for full prescribing information.

**Note to authorized dispenser:** Obtain a Medication Guide by visiting  
[www.upshe-smith.com](http://www.upshe-smith.com) or by calling 1-888-650-3789.

Store at 20° to 25°C (68° to 77°F). Excursions permitted to 15° to 30°C  
(59° to 86°F). [See USP Controlled Room Temperature.] Keep  
tightly closed. Protect from light and moisture. Dispense in a  
tight, light-resistant container with a child-resistant closure.

**Keep out of reach of children.**

SEALED FOR YOUR PROTECTION

Manufactured by  
**UPSHER-SMITH LABORATORIES, LLC**  
Maple Grove, MN 55369

© 2017 Upsher-Smith Laboratories, LLC  
112613-01 R0617



N 0832-1215-00 5

## PRINCIPAL DISPLAY PANEL - 5 mg Tablet Bottle Label

**NDC 0832-1216-00**

**JANTOVEN**<sup>®</sup>

Warfarin Sodium  
Tablets, USP

**5 mg**

**PHARMACIST: Dispense the Medication  
Guide provided separately to each patient.**

**HIGHLY POTENT ANTICOAGULANT WARNING:**

Serious bleeding results from overdosage. Do not  
use or dispense before reading directions and  
warnings in accompanying product information.

100 Tablets

**Rx only**

**UPSHER-SMITH**

NDC 0832-1216-00

**JANTOVEN**<sup>®</sup>

Warfarin Sodium  
Tablets, USP

**5 mg**



**PHARMACIST:** Dispense the Medication Guide provided separately to each patient.

**HIGHLY POTENT ANTICOAGULANT WARNING:** Serious bleeding results from overdosage. Do not use or dispense before reading directions and warnings in accompanying product information.

100 Tablets

**Rx only**

**UPSHER-SMITH**

Each tablet contains: Warfarin Sodium 5 mg

Usual Adult Dosage: See package insert for full prescribing information.

Note to authorized dispenser: Obtain a Medication Guide by visiting [www.upshe-smith.com](http://www.upshe-smith.com) or by calling 1-888-650-3789.

Store at 20° to 25°C (68° to 77°F). Excursions permitted to 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature.] Keep tightly closed. Protect from light and moisture. Dispense in a tight, light-resistant container with a child-resistant closure.

Keep out of reach of children.

SEALED FOR YOUR PROTECTION

Manufactured by  
**UPSHER-SMITH LABORATORIES, LLC**  
Maple Grove, MN 55369

© 2017 Upsher-Smith Laboratories, LLC  
112614-01 R0617



## PRINCIPAL DISPLAY PANEL - 6 mg Tablet Bottle Label

NDC 0832-1217-00

**JANTOVEN**<sup>®</sup>

Warfarin Sodium  
Tablets, USP

**6 mg**

**PHARMACIST:** Dispense the Medication Guide provided separately to each patient.

**HIGHLY POTENT ANTICOAGULANT WARNING:**

Serious bleeding results from overdosage. Do not use or dispense before reading directions and warnings in accompanying product information.

100 Tablets

**Rx only**

**UPSHER-SMITH**

NDC 0832-1217-00

**JANTOVEN<sup>®</sup>**

Warfarin Sodium  
Tablets, USP

**6 mg**



**PHARMACIST:** Dispense the Medication Guide provided separately to each patient.

**HIGHLY POTENT ANTICOAGULANT WARNING:**  
Serious bleeding results from overdosage. Do not use or dispense before reading directions and warnings in accompanying product information.

100 Tablets

**Rx only**

**UPSHER-SMITH**

Each tablet contains: Warfarin Sodium 6 mg

Usual Adult Dosage: See package insert for full prescribing information.

Note to authorized dispenser: Obtain a Medication Guide by visiting [www.upshe-smith.com](http://www.upshe-smith.com) or by calling 1-888-650-3789.

Store at 20° to 25°C (68° to 77°F). Excursions permitted to 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature.] Keep tightly closed. Protect from light and moisture. Dispense in a tight, light-resistant container with a child-resistant closure.

Keep out of reach of children.

SEALED FOR YOUR PROTECTION

Manufactured by  
**UPSHER-SMITH LABORATORIES, LLC**  
Maple Grove, MN 55369

© 2017 Upsher-Smith Laboratories, LLC  
112615-01 R0617



## PRINCIPAL DISPLAY PANEL - 7½ mg Tablet Bottle Label

**NDC 0832-1218-00**

**JANTOVEN<sup>®</sup>**

Warfarin Sodium  
Tablets, USP

**7½ mg**

**PHARMACIST:** Dispense the Medication Guide provided separately to each patient.

**HIGHLY POTENT ANTICOAGULANT WARNING:**  
Serious bleeding results from overdosage. Do not use or dispense before reading directions and warnings in accompanying product information.

100 Tablets

**Rx only**

**UPSHER-SMITH**

NDC 0832-1218-00

**JANTOVEN**<sup>®</sup>

Warfarin Sodium  
Tablets, USP

**7½ mg**



**PHARMACIST:** Dispense the Medication Guide provided separately to each patient.

**HIGHLY POTENT ANTICOAGULANT WARNING:**  
Serious bleeding results from overdosage. Do not use or dispense before reading directions and warnings in accompanying product information.

100 Tablets

**Rx only**

**UPSHER-SMITH**

Each tablet contains: Warfarin Sodium 7½ mg

Usual Adult Dosage: See package insert for full prescribing information.

Note to authorized dispenser: Obtain a Medication Guide by visiting [www.upshe-smith.com](http://www.upshe-smith.com) or by calling 1-888-650-3789.

Store at 20° to 25°C (68° to 77°F). Excursions permitted to 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature.] Keep tightly closed. Protect from light and moisture. Dispense in a tight, light-resistant container with a child-resistant closure.

Keep out of reach of children.

SEALED FOR YOUR PROTECTION

Manufactured by  
**UPSHER-SMITH LABORATORIES, LLC**  
Maple Grove, MN 55369

© 2017 Upsher-Smith Laboratories, LLC  
112616-01 R0617



## PRINCIPAL DISPLAY PANEL - 10 mg Tablet Bottle Label

NDC 0832-1219-50

**JANTOVEN**<sup>®</sup>

Warfarin Sodium  
Tablets, USP

**10 mg**

**PHARMACIST:** Dispense the Medication Guide provided separately to each patient.

**HIGHLY POTENT ANTICOAGULANT WARNING:**

Serious bleeding results from overdosage. Do not use or dispense before reading directions and warnings in accompanying product information.

500 Tablets

**Rx only**

**UPSHER-SMITH**

NDC 0832-1219-50

**JANTOVEN**<sup>®</sup>

Warfarin Sodium  
Tablets, USP



**10 mg**

**PHARMACIST:** Dispense the Medication Guide provided separately to each patient.

**HIGHLY POTENT ANTICOAGULANT WARNING:**  
Serious bleeding results from overdosage. Do not use or dispense before reading directions and warnings in accompanying product information.

500 Tablets

**Rx only**



Each tablet contains: Warfarin Sodium 10 mg

Usual Adult Dosage: See package insert for full prescribing information.

Note to authorized dispenser: Obtain a Medication Guide by visiting [www.upsher-smith.com](http://www.upsher-smith.com) or by calling 1-888-650-3789.

Store at 20° to 25°C (68° to 77°F). Excursions permitted to 15° to 30°C (59° to 86°F). [See USP Controlled Room Temperature.] Keep tightly closed. Protect from light and moisture. Dispense in a tight, light-resistant container with a child-resistant closure.

Keep out of reach of children.

SEALED FOR YOUR PROTECTION

Manufactured by  
**UPsher-SMITH LABORATORIES, LLC**  
Maple Grove, MN 55369

© 2017 Upsher-Smith Laboratories, LLC  
112625-01 R0617



## JANTOVEN

warfarin sodium tablet

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0832-1211
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>WARFARIN SODIUM</b> (UNII: 6153CWM0CL) (WARFARIN - UNII:5Q7ZVV76EI)	WARFARIN SODIUM	1 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>STARCH, CORN</b> (UNII: O8232NY3S)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	

### Product Characteristics

<b>Color</b>	pink	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	WRF;1;832
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC:0832-1211	100 in 1 BOTTLE, Type 0; Not a Combination		

1	NDC:0832-1211-00	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/23/2010	
2	NDC:0832-1211-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	02/23/2010	
3	NDC:0832-1211-01	100 in 1 CARTON	02/23/2010	
3	NDC:0832-1211-89	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040416	02/23/2010	

## JANTOVEN

warfarin sodium tablet

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0832-1212
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
WARFARIN SODIUM (UNII: 6153CWM0CL) (WARFARIN - UNII:5Q7ZVW76EI)	WARFARIN SODIUM	2 mg

### Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	

### Product Characteristics

Color	purple (lavender)	Score	2 pieces
Shape	ROUND	Size	9mm
Flavor		Imprint Code	WRF;2;832
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start	Marketing End
---	-----------	---------------------	-----------------	---------------

#	Item Code	Package Description	Date	Date
1	NDC:0832-1212-00	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/23/2010	
2	NDC:0832-1212-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	02/23/2010	
3	NDC:0832-1212-01	100 in 1 CARTON	02/23/2010	
3	NDC:0832-1212-89	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040416	02/23/2010	

## JANTOVEN

warfarin sodium tablet

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0832-1213
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
WARFARIN SODIUM (UNII: 6153CWM0CL) (WARFARIN - UNII:5Q7ZVV76EI)	WARFARIN SODIUM	2.5 mg

### Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	

### Product Characteristics

Color	green	Score	2 pieces
Shape	ROUND	Size	9mm
Flavor		Imprint Code	WRF;2;1;2;832
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0832-1213-00	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/23/2010	
2	NDC:0832-1213-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	02/23/2010	
3	NDC:0832-1213-01	100 in 1 CARTON	02/23/2010	
3	NDC:0832-1213-89	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040416	02/23/2010	

## JANTOVEN

warfarin sodium tablet

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0832-1214
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
WARFARIN SODIUM (UNII: 6153CWM0CL) (WARFARIN - UNII:5Q7ZVV76EI)	WARFARIN SODIUM	3 mg

### Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
BROWN IRON OXIDE (UNII: 1N032N7MFO)	

### Product Characteristics

Color	brown (tan)	Score	2 pieces
Shape	ROUND	Size	9mm
Flavor		Imprint Code	WRF;3;832
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0832-1214-00	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/23/2010	
2	NDC:0832-1214-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	02/23/2010	
3	NDC:0832-1214-01	100 in 1 CARTON	02/23/2010	
3	NDC:0832-1214-89	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040416	02/23/2010	

## JANTOVEN

warfarin sodium tablet

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0832-1215
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
WARFARIN SODIUM (UNII: 6153CWM0CL) (WARFARIN - UNII:5Q7ZVV76EI)	WARFARIN SODIUM	4 mg

### Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	

### Product Characteristics

Color	blue	Score	2 pieces
Shape	ROUND	Size	9mm
Flavor		Imprint Code	WRF;4;832
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0832-1215-00	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/23/2010	
2	NDC:0832-1215-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	02/23/2010	
3	NDC:0832-1215-01	100 in 1 CARTON	02/23/2010	
3	NDC:0832-1215-89	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040416	02/23/2010	

## JANTOVEN

warfarin sodium tablet

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0832-1216
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
WARFARIN SODIUM (UNII: 6153CWM0CL) (WARFARIN - UNII:5Q7ZVV76EI)	WARFARIN SODIUM	5 mg

### Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

### Product Characteristics

<b>Color</b>	orange (peach)	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	WRF;5;832
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0832-1216-00	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/23/2010	
2	NDC:0832-1216-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	02/23/2010	
3	NDC:0832-1216-01	100 in 1 CARTON	02/23/2010	
3	NDC:0832-1216-89	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040416	02/23/2010	

## JANTOVEN

warfarin sodium tablet

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0832-1217
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
WARFARIN SODIUM (UNII: 6153CWM0CL) (WARFARIN - UNII:5Q7ZVV76EI)	WARFARIN SODIUM	6 mg

### Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	

### Product Characteristics

<b>Color</b>	turquoise (teal)	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	WRF;6;832
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0832-1217-00	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/23/2010	
2	NDC:0832-1217-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	02/23/2010	
3	NDC:0832-1217-01	100 in 1 CARTON	02/23/2010	
3	NDC:0832-1217-89	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040416	02/23/2010	

## JANTOVEN

warfarin sodium tablet

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0832-1218
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
WARFARIN SODIUM (UNII: 6153CWM0CL) (WARFARIN - UNII:5Q7ZVV76EI)	WARFARIN SODIUM	7.5 mg

### Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

### Product Characteristics

Color	yellow	Score	2 pieces
Shape	ROUND	Size	9mm
Flavor		Imprint Code	WRF;7;1;2;832
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0832-1218-00	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/23/2010	
2	NDC:0832-1218-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	02/23/2010	
3	NDC:0832-1218-01	100 in 1 CARTON	02/23/2010	
3	NDC:0832-1218-89	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040416	02/23/2010	

## JANTOVEN

warfarin sodium tablet

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0832-1219
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>WARFARIN SODIUM</b> (UNII: 6153CWM0CL) (WARFARIN - UNII:5Q7ZVW76EI)	WARFARIN SODIUM	10 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	

### Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	9mm
Flavor		Imprint Code	WRF;10;832
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0832-1219-00	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/23/2010	
2	NDC:0832-1219-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	02/23/2010	
3	NDC:0832-1219-01	100 in 1 CARTON	02/23/2010	
3	NDC:0832-1219-89	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040416	02/23/2010	

**Labeler** - Upsher-Smith Laboratories, LLC (047251004)

## Establishment

Name	Address	ID/FEI	Business Operations
Bora Pharmaceuticals, Inc		119296421	analysis(0832-1211, 0832-1212, 0832-1213, 0832-1214, 0832-1215, 0832-1216, 0832-1217, 0832-1218, 0832-1219) , manufacture(0832-1211, 0832-1212, 0832-1213, 0832-1214, 0832-1215, 0832-1216, 0832-1217, 0832-1218, 0832-1219)

Revised: 7/2025

Upsher-Smith Laboratories, LLC