

NICOTINE- nicotine polacrilex lozenge

Walgreens

Drug Facts

Active ingredient (in each lozenge)

Nicotine polacrilex USP, 2 mg

Purpose

Stop smoking aid

Use

- reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding,

only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth problems occur
- persistent indigestion or severe sore throat occurs
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness,

diarrhea, weakness and rapid heartbeat

- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets.

Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **if you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the lozenge on your quit day
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine lozenge
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours

- **nicotine lozenge is a medicine and must be used a certain way to get the best results**
- place the lozenge in your mouth and allow the lozenge to slowly dissolve (about 20 to 30 minutes). Minimize swallowing. **Do not chew or swallow lozenge.**
- you may feel a warm or tingling sensation
- occasionally move the lozenge from one side of your mouth to the other until completely dissolved (about 20 to 30 minutes)
- do not eat or drink 15 minutes before using or while the lozenge is in your mouth
- to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks
- do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects
- **do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day.**
- It is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.

Other information

- **each lozenge contains:** sodium 5 mg
- store at 20° to 25°C (68° to 77°F)
- keep bottle tightly closed and protect from light

Inactive ingredients

acesulfame potassium, colloidal silicon dioxide, cherry flavor, hydroxypropyl cellulose, magnesium stearate, mannitol, potassium bicarbonate, povidone, sodium alginate, sodium carbonate, sucralose, xanthan gum

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredient in Nicorette® Mini Lozenge*

Nicotine Mini Lozenges

NICOTINE POLACRILEX LOZENGE 2 mg

STOP SMOKING AID

Sugar Free

- Includes user's guide

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES OF WAKING UP.

If you smoke your first cigarette **WITHIN 30 MINUTES** of waking up, use Nicotine Polacrilex Lozenge, 4 mg

Cherry Flavor

LOZENGES

TO INCREASE YOUR SUCCESS IN QUITTING:

- 1.** You must be motivated to quit.
- 2.** Use Enough - Use **at least 9** Nicotine Polacrilex lozenges per day during the first six weeks.
- 3.** Use Long Enough - Use Nicotine Polacrilex lozenges for the full 12 weeks.
- 4. Use With a Support Program** as directed in the enclosed User's Guide.
 - **NOT FOR SALE TO THOSE UNDER 18 YEARS OF AGE**
 - **PROOF OF AGE REQUIRED**
 - **NOT FOR SALE IN VENDING MACHINES OR FROM ANY SOURCE WHERE PROOF OF AGE CANNOT BE VERIFIED**

††This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., owners of the registered trademark Nicorette®.

TAMPER EVIDENT FEATURE: DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING.

RETAIN OUTER CARTON FOR FULL PRODUCT USES, DIRECTIONS AND WARNINGS.

DISTRIBUTED BY: WALGREEN CO.

DEERFIELD, IL 60015

Package Label

Drug Facts

Active ingredient (in each lozenge) **Purpose**
Nicotine polacrilex USP, 2 mg. Stop smoking aid

Use ■ reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have ■ heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate. ■ high blood pressure not controlled with medication. Nicotine can increase your blood pressure. ■ stomach ulcer or diabetes ■ history of seizures

Ask a doctor or pharmacist before use if you are ■ using a non-nicotine stop smoking drug ■ taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if ■ mouth problems occur ■ persistent indigestion or severe sore throat occurs ■ irregular heartbeat or palpitations occur ■ you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat ■ you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ if you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you. ■ before using this product, read the enclosed User's Guide for complete directions and other important information ■ begin using the lozenge on your quit day ■ if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine lozenge ■ if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours

■ nicotine lozenge is a medicine and must be used a certain way to get the best results ■ place the lozenge in your mouth and allow the lozenge to slowly dissolve. Minimize swallowing. Do not chew or swallow lozenge. ■ you may feel a warm or tingling sensation ■ occasionally move the lozenge from one side of your mouth to the other until completely dissolved ■ do not eat or drink 15 minutes before using or while the lozenge is in your mouth ■ to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks ■ do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects ■ do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day. ■ it is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.

Other information ■ each lozenge contains: sodium, 5 mg ■ store at 20° to 25°C (68° to 77°F) ■ keep bottle tightly closed and protect from light

Drug Facts (continued)

Inactive ingredients
acesulfame potassium, colloidal silicon dioxide, cherry flavor, hydroxypropyl cellulose, magnesium stearate, mannitol, potassium bicarbonate, povidone, sodium alginate, sodium carbonate, sucralose, xanthan gum

Questions or comments?
Call 1-877-753-3935
Monday-Friday 9AM-5PM EST

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. This product is not affiliated with, manufactured by, or produced by the makers or owners of Nicorette® Lozenge.

For tools, resources and motivation, please visit Walgreens.com/QuitSmoking

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. Use Enough - Use at least 9 Nicotine Polacrilex Lozenges per day during the first six weeks.
3. Use Long Enough - Use Nicotine Polacrilex Lozenges for the full 12 weeks.
4. Use with a support program as directed in the enclosed User's Guide.

Retain outer carton for full product uses, directions and warnings.

NDC 0363-5760-20

Walgreens

Compare to the active ingredient in Nicorette® Mini Lozenge®

Nicotine Mini Lozenge

NICOTINE POLACRILEX LOZENGE 2 mg
STOP SMOKING AID

Sugar Free

Includes user's guide

20
LOZENGES

2
mg

2 mg EACH (1 bottle of 20)

Cherry flavor

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP. If you smoke your first cigarette **WITHIN 30 MINUTES** of waking up, use Nicotine Polacrilex Lozenge, 4 mg.

PAPER BOX PLASTIC BOTTLE

W30RG0122-F
REV-0224

DISTRIBUTED BY: WALGREEN CO., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2021 Walgreen Co.
ITEM 917201 W11042-0224-L
PRODUCT OF SWITZERLAND
PLD-E695A FC009093

Lot No.:
Exp. Date:

3 11917 03253 5

WALGREENS Nicotine Mini Lozenge Cherry Flavor

NICOTINE

nicotine polacrilex lozenge

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-5760
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	2 mg

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MANNITOL (UNII: 3OWL53L36A)	
POTASSIUM BICARBONATE (UNII: HM5Z15LEBN)	
POVIDONE (UNII: FZ989GH94E)	
SODIUM ALGINATE (UNII: C269C4G2ZQ)	
SODIUM CARBONATE (UNII: 45P3261C7T)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor	CHERRY	Imprint Code	A501
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-5760-20	1 in 1 BOX	03/26/2021	
1		20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210711	03/26/2021	

Labeler - Walgreens (008965063)

Revised: 2/2024

Walgreens