

**WINCO FOODS STOMACH RELIEF- bismuth subsalicylate suspension**  
**WINCO FOODS**

-----  
**WinCo<sup>®</sup> Stomach Relief Regular Strength Bismuth Subsalicylate**

**Active ingredient (in each 30 mL dose)**

Bismuth subsalicylate 525 mg

**Purposes**

Upset stomach reliever and antidiarrheal

**Uses**

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn
- indigestion
- nausea
- gas
- belching

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea or vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use if you have**

- an ulcer
- a bleeding problem
- bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool

**Ask a doctor or pharmacist before use if you are**

taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

### **When using this product**

a temporary, but harmless, darkening of the stool and/or tongue may occur

### **Stop use and ask a doctor if**

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- shake well before use
- for accurate dosing, use dose cup
- **adults and children 12 years and over:**(30 mL) every ½ hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
- 30 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (240 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- **children under 12 years:**ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

### **Other information**

- **each 30 mL contains:**sodium 10 mg
- **salicylate 261 mg**
- low sodium
- sugar free
- store at room temperature
- protect from freezing
- avoid excessive heat (more than 104°F 40°C)

### **Inactive ingredients**

carboxymethylcellulose sodium, D&C Red No. 22, D&C Red No. 28, flavor,

microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum

### Questions?

Call **1-800-824-1706 Monday - Friday 9am - 4pm MST**

**Do not use if imprinted shrink band is missing or broken.**

### Principal Display Panel

WinCo® FOODS

NDC# **67091-362-23**

**Compare To the active ingredient in PEPTO-BISMOL® Regular Strength.**

Stomach Relief

**BISMUTH SUBSALICYLATE/**

**UPSET STOMACH RELIEVER & ANTI-DIARRHEAL**

**Soothing Relief from:** Upset Stomach , Indigestion , Nausea , Heartburn & Diarrhea

Regular Strength

8 FL. OZ. (236 mL)

**DISTRIBUTED BY:**

WINCO FOODS, LLC, BOISE, ID 83704

\*This product is not manufactured or distributed by PROCTER & GAMBLE INC., distributor of PEPTO-BISMOL® Regular Strength.

| <p><b>Drug Facts (continued)</b></p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.</p> <p><b>Directions</b> ■ shake well before use ■ only use dose cup provided ■ <b>adults and children 12 years and over:</b> ■ 30 mL (1 dose) every ½ hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea ■ 30mL (1dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) ■ do not exceed 8 doses (240 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ <b>children under 12 years:</b> ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea</p> <p><b>Other information</b> ■ each 30 mL contains: sodium 10 mg ■ salicylate 261 mg ■ low sodium ■ sugar free ■ store at room temperature ■ protect from freezing ■ avoid excessive heat (more than 104°F or 40°C)</p> <p><b>Inactive ingredients</b> carboxymethylcellulose sodium, D&amp;C Red No 22, D&amp;C Red No. 28, flavor, microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum</p> <p><b>Questions?</b> Call 800-824-1706 Monday - Friday 9am - 4pm MST</p> | <p>*Compare to the active ingredient in PEPTO-BISMOL® Regular Strength</p>  <p><b>BISMUTH SUBSALICYLATE/<br/>UPSET STOMACH RELIEVER<br/>&amp; ANTI-DIARRHEAL</b></p> <p>Soothing Relief for: Upset Stomach,<br/>Indigestion, Nausea, Heartburn &amp; Diarrhea</p>  | <p><b>Do not use if imprinted shrink band is missing or broken</b></p> <p><b>Drug Facts</b></p> <table border="1"> <thead> <tr> <th>Active ingredient<br/>(in each 30 mL dose)</th> <th>Purposes</th> </tr> </thead> <tbody> <tr> <td>Bismuth subsalicylate 525 mg</td> <td>Upset stomach reliever and antidiarrheal</td> </tr> </tbody> </table> <p><b>Uses</b> relieves ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including:<br/>■ heartburn ■ indigestion ■ nausea ■ gas ■ belching ■ fullness</p> <p><b>Warnings</b></p> <p><b>Reye's syndrome:</b> Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p><b>Allergy alert: Contains salicylate. Do not take if you are</b><br/>■ allergic to salicylates (including aspirin)<br/>■ taking other salicylate products</p> <p><b>Do not use if you have</b> ■ an ulcer ■ a bleeding problem ■ bloody or black stool</p> <p><b>Ask a doctor before use if you have</b> ■ fever ■ mucus in the stool</p> <p><b>Ask a doctor or pharmacist before use if you are taking any drug for</b> ■ anticoagulation (thinning the blood) ■ diabetes ■ gout ■ arthritis</p> <p><b>When using this product</b> a temporary, but harmless, darkening of the stool and/or tongue may occur.</p> <p><b>Stop use and ask a doctor if</b> ■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days</p> | Active ingredient<br>(in each 30 mL dose) | Purposes | Bismuth subsalicylate 525 mg | Upset stomach reliever and antidiarrheal |
|---|---|---|---|----------|------------------------------|--|
| Active ingredient<br>(in each 30 mL dose)   | Purposes  |   |   |          |                              |  |
| Bismuth subsalicylate 525 mg  | Upset stomach reliever and antidiarrheal  |   |   |          |                              |  |
| <p>LOT: UNVARNISHED AREA EXP:</p> <p>DISTRIBUTED BY:<br/>WINCO FOODS, LLC, BOISE, ID 83704</p>  <p>0 70552 24150 3</p>   | <p><small>*This product is not manufactured or distributed by PROCTER &amp; GAMBLE INC., distributor of PEPTO-BISMOL® Regular Strength.</small></p>   |   |   |          |                              |  |

# WINCO FOODS STOMACH RELIEF

bismuth subsalicylate suspension

## Product Information

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:67091-362 |
| <b>Route of Administration</b> | ORAL           |                           |               |

## Active Ingredient/Active Moiety

| Ingredient Name  | Basis of Strength     | Strength        |
|--|-----------------------|-----------------|
| <b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ) | BISMUTH SUBSALICYLATE | 525 mg in 30 mL |

## Inactive Ingredients

| Ingredient Name  | Strength |
|--|----------|
| <b>CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED</b> (UNII: K679OBS311) |          |
| <b>D&amp;C RED NO. 22</b> (UNII: 1678RKX8RT)                         |          |
| <b>D&amp;C RED NO. 28</b> (UNII: 767IPOY5NH)                         |          |
| <b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)                 |          |
| <b>POTASSIUM HYDROXIDE</b> (UNII: WZH3C48M4T)                        |          |
| <b>POTASSIUM SORBATE</b> (UNII: 1VPU26JZZ4)                          |          |
| <b>WATER</b> (UNII: 059QF0K00R)                                      |          |
| <b>SALICYLIC ACID</b> (UNII: O414PZ4LPZ)                             |          |
| <b>DIMETHICONE, UNSPECIFIED</b> (UNII: 92RU3N3Y1O)                   |          |
| <b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)                            |          |
| <b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)                                  |          |
| <b>XANTHAN GUM</b> (UNII: TTV12P4NEE)                                |          |

## Packaging

| # | Item Code        | Package Description                                   | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:67091-362-23 | 236 mL in 1 BOTTLE; Type 0: Not a Combination Product | 02/24/2020           |                    |

## Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M008                                     | 02/24/2020           |                    |

**Labeler** - WINCO FOODS (056098817)

