

**WINCO FOODS STOMACH RELIEF- bismuth subsalicylate suspension**  
**WINCO FOODS**

-----  
**WinCo® Stomach Relief Regular Strength Bismuth Subsalicylate**

**Active ingredient (in each 30 mL dose)**

Bismuth subsalicylate 525 mg

**Purposes**

Upset stomach reliever and antidiarrheal

**Uses**

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn
- indigestion
- nausea
- gas
- belching

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea or vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use if you have**

- an ulcer
- a bleeding problem
- bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool

## **Ask a doctor or pharmacist before use if you are**

taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

## **When using this product**

a temporary, but harmless, darkening of the stool and/or tongue may occur

## **Stop use and ask a doctor if**

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

## **If pregnant or breast-feeding,**

ask a health professional before use.

## **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

## **Directions**

- shake well before use
- for accurate dosing, use dose cup
- **adults and children 12 years and over:** (30 mL) every ½ hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
- 30 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (240 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- **children under 12 years:** ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

## **Other information**

- **each 30 mL contains:** sodium 10 mg
- **salicylate 261 mg**
- low sodium
- sugar free
- store at room temperature
- protect from freezing
- avoid excessive heat (more than 104°F 40°C)

## **Inactive ingredients**

carboxymethylcellulose sodium, D&C Red No. 22, D&C Red No. 28, flavor, microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum

## **Questions?**

Call **1-800-824-1706 Monday - Friday 9am - 4pm MST**

**Do not use if imprinted shrink band is missing or broken.**

## **Principal Display Panel**

WinCo® FOODS

NDC# **67091-362-23**

**Compare To the active ingredient in PEPTO-BISMOL® Regular Strength.**

Stomach Relief

**BISMUTH SUBSALICYLATE/**

**UPSET STOMACH RELIEVER & ANTI-DIARRHEAL**

**Soothing Relief from:** Upset Stomach, Indigestion, Nausea, Heartburn & Diarrhea

Regular Strength

8 FL. OZ. (236 mL)

**DISTRIBUTED BY:**

WINCO FOODS, LLC, BOISE, ID 83704

\*This product is not manufactured or distributed by PROCTER & GAMBLE INC., distributor of PEPTO-BISMOL® Regular Strength.

**Drug Facts (continued)**

If pregnant or breast-feeding, ask a health professional before use.  
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.

**Directions** ■ shake well before use ■ only use dose cup provided ■ **adults and children 12 years and over:** ■ 30 mL (1 dose) every ½ hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea ■ 30 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) ■ do not exceed 8 doses (240 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ **children under 12 years:** ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information** ■ each 30 mL contains: sodium 10 mg ■ salicylate 261 mg ■ low sodium ■ sugar free ■ store at room temperature ■ protect from freezing ■ avoid excessive heat (more than 104°F or 40°C)

**Inactive ingredients** carboxymethylcellulose sodium, D&C Red No 22, D&C Red No. 28, flavor, microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum

**Questions?** Call 800-824-1706 Monday - Friday 9am - 4pm MST

\*Compare to the active ingredient in PEPTO-BISMOL® Regular Strength



**BISMUTH SUBSALICYLATE/  
UPSET STOMACH RELIEVER  
& ANTI-DIARRHEAL**

Soothing Relief for: Upset Stomach,  
Indigestion, Nausea, Heartburn & Diarrhea

8 FL OZ (236 mL)

Do not use if imprinted shrink band is missing or broken

**Drug Facts**

**Active ingredient (in each 30 mL dose)** Bismuth subsalicylate 525 mg .....Upset stomach reliever and antidiarrheal

**Uses** relieves ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including: ■ heartburn ■ indigestion ■ nausea ■ gas ■ belching ■ fullness

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are** ■ allergic to salicylates (including aspirin) ■ taking other salicylate products

**Do not use if you have** ■ an ulcer ■ a bleeding problem ■ bloody or black stool

**Ask a doctor before use if you have** ■ fever ■ mucus in the stool

**Ask a doctor or pharmacist before use if you are taking any drug for** ■ anticoagulation (thinning the blood) ■ diabetes ■ gout ■ arthritis

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur.

**Stop use and ask a doctor if** ■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days

\*This product is not manufactured or distributed by PROCTER & GAMBLE INC., distributor of PEPTO-BISMOL® Regular Strength.

LOT:  
UNVARNISHED  
AREA  
EXP:

DISTRIBUTED BY:  
WINCO FOODS, LLC, BOISE, ID 83704

**WINCO FOODS STOMACH RELIEF**

bismuth subsalicylate suspension

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:67091-362
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:Z59CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 30 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED</b> (UNII: K679OBS311)	
<b>D&amp;C RED NO. 22</b> (UNII: 1678RKX8RT)	
<b>D&amp;C RED NO. 28</b> (UNII: 7671P0Y5NH)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POTASSIUM HYDROXIDE</b> (UNII: WZ H3C48M4T)	
<b>POTASSIUM SORBATE</b> (UNII: 1VPU26JZ4)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SALICYLIC ACID</b> (UNII: O414PZ4LPZ)	
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67091-362-23	236 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/24/2020	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	02/24/2020	

**Labeler - WINCO FOODS (056098817)**

Revised: 12/2023

WINCO FOODS