

**VICKS NYQUIL KIDS COLD AND COUGH PLUS FEVER- acetaminophen,
dextromethorphan hydrobromide, doxylamine succinate liquid
The Procter & Gamble Manufacturing Company**

VICKS® NyQuil™ Kids COLD & COUGH + FEVER

Drug Facts

Active ingredients (in each 15 mL)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Doxylamine succinate 6.25 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat • headache • minor aches & pains • fever
- runny nose & sneezing

Warnings

Liver warning:

This product contains acetaminophen.

Severe liver damage may occur if

- adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product
- child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy Alert:

Acetaminophen may cause severe skin reactions.

Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 15 mL contains:**sodium 13 mg
- store at no greater than 25°C

Inactive ingredients

citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

Questions?

1-800-362-1683

TAMPER EVIDENT: DO NOT USE IF PRINTED SHRINKBAND ON BOTTLE IS BROKEN OR MISSING.

DIST. BY:

PROCTER & GAMBLE

CINCINNATI, OH 45202

PRINCIPAL DISPLAY PANEL - 236 ml Bottle Label

VICKS[®]

NyQuil[™]

Kids

COLD & COUGH

+ FEVER

Acetaminophen, Doxylamine Succinate, Dextromethorphan HBr

Fever, Sore Throat, Minor Aches & Pains

Sneezing, Runny Nose

Cough

AGES 6+

Nighttime Relief

8 FL OZ (236 mL)



Drug Facts (continued)

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur • avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present • new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 15 mL contains: sodium 13 mg
- store at no greater than 25°C

Inactive ingredients citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

Questions? 1-800-362-1683

Drug Facts (continued)

liver damage may occur if

- adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product
- child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product
- adult has 3 or more alcoholic drinks every day while using this product
- taken with other drugs containing acetaminophen

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy

Ask a doctor before use if you have

- liver disease • glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland



VICKS NYQUIL KIDS COLD AND COUGH PLUS FEVER

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:84126-305
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH)	DEXTROMETHORPHAN	10 mg

(DEXTROMETHORPHAN - UNII:7355X3ROTS)	HYDROBROMIDE	in 15 mL
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
SORBITOL (UNII: 506T60A25R)	
XANTHAN GUM (UNII: TTV12P4NEE)	
GLYCERIN (UNII: PDC6A3C0OX)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	

Product Characteristics

Color	purple	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84126-305-08	236 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/06/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/06/2024	

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Revised: 12/2025

The Procter & Gamble Manufacturing Company