

**EXTRA FAST COLTALIN- acetaminophen, chlorpheniramine maleate,
phenylephrine hydrochloride tablet
FORTUNE PHARMACAL COMPANY LIMITED**

Drug Facts

Active ingredients(in each tablet)

Acetaminophen 325 mg

Chlorpheniramine maleate 2 mg

Phenylephrine hydrochloride 5 mg

Purpose

Pain reliever-fever reducer

Antihistamine

Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold, the flu, or hay fever:
 - minor aches and pains
 - sore throat
 - headache
 - runny nose
 - sinus congestion and pressure
 - nasal congestion due to a cold or hay fever
 - runny nose
 - sneezing, itching of the nose or throat, and itchy, watery eyes due to hay fever
- temporarily reduces fever
- promotes nasal and/or sinus drainage

Warnings

Liver Warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 12 tablets in 24 hours, which is the maximum daily amount for this product
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks everyday while using this product

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin rash occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know whether a prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if the user has

- heart disease
- high blood pressure
- diabetes
- glaucoma
- liver disease
- thyroid disease
- a high fever
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if the user is

- taking tranquilizers or sedatives
- taking the blood thinning drug warfarin

When using this product

- **do not exceed recommended dosage**
- may cause excitability especially in children
- use caution when driving or operating machinery
- avoid alcoholic beverages
- may cause drowsiness; alcohol, sedatives, and tranquilizers may increase the drowsiness effect

Stop use and ask a doctor if

- pain or nasal congestion gets worse or lasts more than 5 days (children) or 7 days (adults)
- sore throat persists for more than 2 days
- fever gets worse or lasts more than 3 days
- nasal congestion is accompanied by fever
- nervousness, dizziness, or sleeplessness occur
- redness or swelling is present
- new symptoms occur
- any of the following occurs (these could be signs of a serious condition):
 - severe sore throat
 - sore throat is accompanied or followed by fever, headache, rash, nausea or vomiting

Do not give to children under 6 years of age unless directed by doctor.

If pregnant or breast-feeding,ask a health professional before use.

Keep out of reach of children.In case of overdose, get medical help or contact a Poison Control Center right away.

Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

■ adults and children 12 years of age and over: 2 tablets every 4 hours, while symptoms persist, not to exceed 6 doses (12 tablets) in 24 hours, or as directed by a doctor

■ children 6 to under 12 years of age: 1 tablet every 4 hours, while symptoms persist, not to exceed 5 doses (5 tablets) in 24 hours, or as directed by a doctor

■ children under 6 of age : consult a doctor

Other information

■ keep tightly sealed

■ store between below 25 C (77 F) in a dry place, away from direct sunlight

Inactive ingredients

Microcrystalline Cellulose, corn starch, FD&C yellow no. 6 aluminum lake, sodium starch glycolate, Citric Acid Monohydrate, Hydroxypropyl Methylcellulose, Colloidal silicon dioxide, magnesium stearate

Questions or comments? (888) 221-3496 M-F 9 am to 5 pm

■ you may also report serious side effects to this phone number

EXTRA FAST COLTALIN

Safely Sealed

NDC 51467-014-01

Acetaminophen - Pain Reliever-Fever Reducer

Chlorpheniramine maleate - Antihistamine

Phenylephrine hydrochloride - Nasal Decongestant

Cold, Flu & Hay Fever Multi-Symptom Relief

■ Minor Aches & Pain, Sore Throat

■ Headache, Fever

■ Nasal Congestion, Sinus Pressure/Drainage

■ Sneezing, Runny Nose

■ Itchy, Watery Eyes

24 Tablets

In Individually Sealed Strip

Aspirin Free

Product code

LOT
EXP

EXTRA-FAST

SAFELY SEALED • NDC 51467-014-01

COLTALIN[®]

GOLD & FLU

速效幸福傷風感冒素

Acetaminophen - Pain Reliever - Fever Reducer
Chlorpheniramine Maleate - Antihistamine
Phenylephrine Hydrochloride - Nasal Decongestant

Cold, Flu & Hay Fever Multi-Symptom Relief

傷風、流感、花粉熱 舒緩各種症狀

- Minor Aches & Pain, Sore Throat 輕微疼痛、喉嚨痛
- Headache, Fever 發熱、頭痛
- Nasal Congestion, Sinus Pressure/Drainage 鼻塞、鼻竇充血
- Sneezing, Runny Nose 打噴嚏、流鼻涕
- Itchy, Watery Eyes 眼睛發癢、淚眼



24
Tablets

In Individually Sealed Strip
獨立鋁箔密封包裝

ASPIRIN FREE



PRINTED WITH SOY INK

24 Tablets/H

速效幸福傷風感冒素

Please refer to enclosed package insert for Chinese translation of drug facts.
請參閱內附之說明書以查看中文藥品資料。



Distributed by:
SOLSTICE MEDICINE COMPANY
美國蘇氏藥業有限公司
215 W. Ave Street
Los Angeles, CA 90012 U.S.A.
Tel: (888) 221-3496
Fax: (323) 221-2448
www.smrmed.com

PAPER

Pharma code

Pharma code

Drug Facts

Active Ingredients (in each tablet)
Acetaminophen 325 mg, Pain Reliever/Fever Reducer
Chlorpheniramine maleate 2 mg, Antihistamine
Phenylephrine hydrochloride 5 mg, Nasal decongestant

Uses
Temporarily relieves these signs and symptoms due to a cold, the flu, or hay fever:
• minor aches and pains • sore throat • headache
• sinus congestion and pressure • nasal congestion • runny nose
• sneezing, itching of the nose or throat, and itchy, watery eyes
Temporarily reduces fever.
Promotes nasal and/or sinus drainage

Warnings
Liver Warning: This product contains acetaminophen. Severe liver damage may occur if adult takes more than 12 tablets in 24 hours, which is the maximum daily amount for this product.
Child takes more than 5 doses in 24 hours.
Taken with other drugs containing acetaminophen.
Adult has 3 or more alcoholic drinks everyday while using this product.
Always alert: Acetaminophen may cause severe skin reactions.
If a skin reaction occurs, stop use and seek medical help right away.
Symptoms may include: skin redness, blisters, rash.

Do not use
• with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, Parkinson's disease, or migraine) or Parkinson's disease, or for 2 weeks after stopping the MAOI drug.
If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if the user has
heart disease • high blood pressure • diabetes • glaucoma
liver disease • thyroid disease • high fever
difficulty in urination due to enlargement of the prostate gland
• a breathing problem such as emphysema or chronic bronchitis

Ask a doctor or pharmacist before use if the user is
taking tranquilizers or sedatives • taking the blood thinning drug warfarin
• using caution when driving or operating machinery
• may cause excitability especially in children
• do not exceed recommended dosage

When using this product
• avoid alcoholic beverages

Other information
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Forming medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Forming medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
adult and childen 12 years of age and over
take 2 tablets every 4 hours, while symptoms persist, not to exceed 6 doses (12 tablets) in 24 hours, or as directed by a doctor.
children 6 to under 12 years of age
take 1 tablet every 4 hours, while symptoms persist, not to exceed 5 doses (5 tablets) in 24 hours, or as directed by a doctor.
children under 6 years of age
consult a doctor

Inactive ingredient
FD&C Yellow No. 6 Aluminum Lake, Sodium Starch Glycolate, Citric Acid Monohydrate, Hydroxypropyl Methylcellulose, Colloidal silicon dioxide, Magnesium stearate.

Questions or comments? (888) 221-3496 M-F 9 am to 5 pm
• you may also report serious side effects to this phone number

COLTALIN[®]
EXTRA-FAST
GOLD & FLU

Do not use if individually sealed foil unit dose pouches are torn, broken, or show evidence of being opened. Keep outer carton for complete warnings and product information.

Manufactured by:
Fortune Pharmaceutical Co., Ltd.
香港 幸福藥業有限公司
Made in China



EXTRA FAST COLTALIN			
acetaminophen, chlorpheniramine maleate, phenylephrine hydrochloride tablet			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51467-014
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	325 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)		CHLORPHENIRAMINE MALEATE	2 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)		PHENYLEPHRINE	5 mg
Inactive Ingredients			
Ingredient Name			Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
STARCH, CORN (UNII: O8232NY3S)			
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)			
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)			
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)			

HYDROXYPROPYL METHYLCELLULOSE (UNII: 3NXW29V3WO)

Product Characteristics

Color	orange	Score	no score
Shape	ROUND	Size	12mm
Flavor		Imprint Code	SHIELDTM
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51467-014-01	2 in 1 BOX	01/30/2026	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:51467-014-02	3 in 1 BOX	01/30/2026	
2		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/30/2026	

Labeler - FORTUNE PHARMACAL COMPANY LIMITED (686280561)

Registrant - FORTUNE PHARMACAL COMPANY LIMITED (686280561)

Establishment

Name	Address	ID/FEI	Business Operations
FORTUNE PHARMACAL COMPANY LIMITED		686280561	manufacture(51467-014)

Revised: 2/2026

FORTUNE PHARMACAL COMPANY LIMITED