GILTUSS DM- dextromethorphan hbr syrup Gil Pharmaceutical Corp

Giltuss ® DM High Blood Pressure Dextromethorphan HBr, 30 mg Maximum Strength

Drug Facts

Active ingredient (in each 10 mL) Purpose

Dextromethorphan HBr. 30 mg Cough Suppresant

Uses

- temporarily relieves:
- cough due to minor throat and bronchial irritation associated with a common cold or inhaled irritants
- the intensity of coughing
- the impulse to cough to help you get to sleep

Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- When using this product do not use more than directed.

Ask a doctor before use if you have

- a cough with too much phlegm (mucus)
- a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor if

cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache as these could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

- Keep out of reach of children.
- In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not exceed recommended dosage
- do not take more than 4 doses in any 24-hour period
- use enclosed dosing cup
- ml= milliliter

Adults and children over 12	10 mL every 8	
years	hours	
Children under 6 years to	5 mL every 8	
under 12 years	hours	
Children 4 years to under 6	2.5 mL* every	
years	8 hours	
Children under 4 years	Do not use	

^{*}use appropriate measuring syringe or measuring cup

Other information

- store at room temperature 15°-30°c (59°-86°F)
- close cap tightly
- SAVE CARTON FOR COMPLETE DRUG FACTS

Inactive ingredients

Acesulfame potassium, citric acid, flavor, glycerin, menthol, polyethylene glycol, povidone, propylene glycol, purified water, sodium benzoate, sodium saccharin.

Questions or comments?

Call **1-787-848-9114**, Monday-Friday 9AM-5PM EST. Call your doctor for medical advice in the event of side effects.

Label revised: 02/23

MANUFACTURED FOR GIL PHARMACEUTICAL CORP. PONCE, PUERTO RICO 00717-1565

PRINCIPAL DISPLAY PANEL

Giltuss DM High Blood Pressure - NDC 58552-150-04 - Bottle Label

Uses

temporarily relieves:

- · cough due to minor throat and bronchial irritation associated with a common cold or inhaled irritants
- · the intensity of coughing
- · the impulse to cough to help you get to sleep

Do not use

- No not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- · When using this product do not use more than directed.

Ask a doctor before use if you have

- · a cough with too much phlegm (mucus)
- a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache as these could be signs of a serious condition.



Drug Facts (continued).

If pregnant or breast-feeding, ask a health professional before use.

· Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- · do not exceed recommended dosage
- . do not take more than 4 doses in any 24-hour period
- · use enclosed dosing cup
- mL = milliliter

Adults and children over 12 years	10 mL every 8 hours
Children under 6 years to under 12 years	5 mL every 8 hours
Children 4 years to under 6 years	2.5 mL* every 8 hours
Children under 4 years	Do not use

* use appropriate measuring syringe or measuring cup

- Other information
 store at room temperature 15°-30°C (59°-86°F)
- SAVE CARTON FOR COMPLETE DRUG FACTS

Acesulfame potassium, citric acid, flavor, glycerin, menthol, polyethylene glycol, povidone, propylene glycol, purified water, sodium benzoate, sodium saccharin.

Questions or comments?

Call 1-787-848-9114, Monday—Friday 9AM-5PM EST, Call your doctor for medical advice in the event of side effects.

revised: 02/23 abe

MANUFACTURED FOR GIL PHARMACEUTICAL CORP. PONCE, PUERTO RICO 00717-1565

Giltuss DM High Blood Pressure - NDC 58552-150-04 - Carton Label



MAXIMUM STRENGTH



HIGH BLOOD PRESSURE

Dextromethorphan HBr, 30 mg

ALCOHOL FREE. SUGAR FREE. DYE FREE.

RELIEVES ALIVIA

PERSISTENT ✓ TOS COUGH PERSISTENTE UP TO 8 HOURS HASTA POR 8 HORAS




4FL OZ (118 mL)

HIGH BLOOD PRESSURE

ANTITUSSIVE STRENGTH TO CONTROL COUGH **SYMPTOMS UP TO 8 HOURS**

ALCOHOL FREE. SUGAR FREE. DYE FREE.



Drug Facts

Active ingredient (in each 10 mL) Purpose Dextromethorphan HBr. 30 mg_Cough Suppresant Drug Facts (contin

 use enclosed dosing cup · mL = milliter

Adults and children over 12 years

Children 4 years to under 6 years

use appropriate measuring syringe or measuring cup

Children under 4 years

do not exceed recommended dosage do not take more than 4 doses in any 24-hour period

Children under 6 years to under 12 years 5 ml. every 8 hours

store at room temperature 15°-30°C (59°-86°F)

close cap tightly
 SAVE CARTON FOR COMPLETE DRUG FACTS

Inactive ingredients Acesulfame potassium, citric acid, flavor, glycerin,

menthol, polyethylene glycol, povidone, propylene glycol, purified water, sodium benzoate, sodium

Questions or comments? Call 1-787-848-9114, Monday—Friday 9AM-5PM EST.

ALCOHOL FREE, SUGAR FREE, DYE FREE,

dent: do not use if safety seal under or aroun

Call your doctor for medical advice in the event of

2,5 mL* every 8 hours

Label revised: 02/23

Directions

- temporarily relieves:
- + cough due to minor throat and bronchial irritation associated with a common cold or inhaled irritants
- · the intensity of coughing
- . the impulse to cough to help you get to sleep

Warnings

Do not use

- . if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- . When using this product do not use more than directed.

Ask a doctor before use if you have

- a cough with too much philegm (mucus)
- . a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor if

cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache as these could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

- Keep out of reach of children.
- · In case of overdose, get medical help or contact a Poison Control Center right away.



MANUFACTURED FOR GIL PHARMACEUTICAL CORP. PONCE, PUERTO RICO 00717-1565



giltuss.com

GILTUSS DM

dextromethorphan hbr syrup

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:58552-150

Route of Administration ORAL

Active Ingredient/Active Moiety

Basis of Strength Ingredient Name Strength **DEXTROMETHORPHAN HYDROBROMIDE** (UNII: 9D2RTI9KYH) **DEXTROMETHORPHAN** 30 mg (DEXTROMETHORPHAN - UNII:7355X3ROTS) **HYDROBROMIDE** in 10 mL

Inactive Ingredients

Ingredient Name

Strength

ACESULFAME POTASSIUM (UNII: 230V73Q5G9)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
RASPBERRY (UNII: 4N14V5R27W)	
GLYCERIN (UNII: PDC6A3C0OX)	
MENTHOL (UNII: L7T10EIP3A)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ 989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	

Product Characteristics		
Color	white (clear color)	Score
Shape		Size
Flavor	RASPBERRY	Imprint Code
Contains		

ı	Packaging				
4	# Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:58552-150- 04	1 in 1 CARTON	08/09/2023		
1	L	118 mL in 1 BOTTLE; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/27/2023	

Labeler - Gil Pharmaceutical Corp (176826592)

Revised: 12/2023 Gil Pharmaceutical Corp