

**FOSTER AND THRIVE MEDICATED CORN REMOVERS- 40% w/w salicylic acid
corn remover plaster
Strategic Sourcing Services LLC**

Foster and Thrive Medicated Corn Remover

Active Ingredient

Salicylic Acid 40%

Purpose

Corn Remover

Uses

- for removal of corns
- relieves pain by removing corns

Warning

For external use only

Do Not Use

- if you are a diabetic
- if you have poor blood circulation
- on irritated skin or area that is infected or reddened

Stop use and ask a doctor

if discomfort lasts

Keep out of reach of children

If swallowed get medical help or contact a Poison Control Center right away. 800-222-1222

Directions

- wash affected area, dry thoroughly
- If necessary, cut medicated disk to fit corn
- apply medicated disk with sticky side adhering to skin
- cover medicated disk with enclosed cushion
- after 48 hours, remove medicated disk
- repeat procedure every 48 hours as needed for up to 14 days (until corn is removed)

- may soak corn in warm water for 5 minutes to assist in removal

Other Information

Store between 20 °C to 25 °C (68 °F to 77 °F)

Inactive Ingredient

vinyl acetate

Clear of Copy or Logo

*This product is not manufactured or distributed by Yellow Wood Partners LLC, distributor of Dr. Scholl's®.

©2023 McKesson Corporation
Distributed by: McKesson Corp.,
via Strategic Sourcing Services LLC
Memphis, TN 38141
Money Back Guarantee
www.fosterandthrive.com
Product of USA

R004528-A



0 10939 96029 0

NDC 79677-1237-01

COMPARE TO DR. SCHOLL'S®

Foster & Thrive™

Medicated Corn Removers

Salicylic Acid

CLEAR WINDOW
SHOWING
PRODUCT



- Medicated discs safely remove corns
- Soft cushions provide pressure relief

9 CUSHIONS
9 MEDICATED DISCS

Drug Facts	
Active Ingredient	Purpose
Salicylic acid 40%.....	Corn Remover
Uses	
<ul style="list-style-type: none"> for removal of corns relieves pain by removing corns 	
Warning	
For external use only.	
Do not use	
<ul style="list-style-type: none"> if you are a diabetic if you have poor blood circulation on irritated skin or any area that is infected or reddened 	
Stop use and ask a doctor if	
discomfort lasts	
Keep out of reach of children.	
If swallowed, get medical help or contact a Poison Control Center right away. 800-222-1222	
Directions	
<ul style="list-style-type: none"> wash affected area and dry thoroughly if necessary, cut medicated disk to fit corn apply medicated disc with sticky side adhering to skin cover medicated disc with enclosed cushion after 48 hours, remove medicated disc repeat procedure every 48 hours as needed for up to 14 days (until corn is removed) may soak corn in warm water for 5 minutes to assist in removal 	
Other information	
Store between 20°C to 25°C (68°F to 77°F)	
Inactive ingredient	
vinyl acetate	
Questions or Comments? Call	
833-358-6431 Monday to Friday 9:00am to 7:00pm EST	
KEEP OUTER CARTON FOR COMPLETE WARNING AND PRODUCT INFORMATION.	
TAMPER EVIDENT: DO NOT USE IF CARTON IS OPEN/SEALS BROKEN.	
Rev. 02/24	

No Copy or Logo (Except R#'s)

Clear of Ink Except Eyespots

Clear of Ink Except Eyespots

FOSTER AND THRIVE MEDICATED CORN REMOVERS

40% w/w salicylic acid corn remover plaster

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70677-1237
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SALICYLIC ACID (UNII: O414PZ4LPZ) (SALICYLIC ACID - UNII:O414PZ4LPZ)	SALICYLIC ACID	40 mg in 100 mg

Inactive Ingredients

Ingredient Name	Strength
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
ACRYLIC ACID/ETHYLENE COPOLYMER (600 MPA.S) (UNII: 1PEZ3NLY6I)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70677-1237-1	100 mg in 1 BOX; Type 0: Not a Combination Product	05/07/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M030	05/07/2024	

Labeler - Strategic Sourcing Services LLC (116956644)