

**COLD AND FLU DAYTIME RELIEF- acetaminophen, dextromethorphan hbr,
phenylephrine hcl solution
L.N.K. International, Inc.**

Convenience Valet 44-012

Active ingredients (in each 15 mL)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Nasal decongestant

Uses

- temporarily relieves common cold and flu symptoms:
 - sore throat
 - nasal congestion
 - headache
 - minor aches and pains
 - cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if the user has

- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- heart disease
- diabetes
- thyroid disease
- liver disease
- high blood pressure

Ask a doctor or pharmacist before use if the user is

taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**

- do not exceed 4 doses per 24 hours
- mL = milliliter
- only use the dose cup provided

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children under 6 years	do not use

Other information

- **each 15 mL contains:** sodium 13 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

anhydrous citric acid, FD&C yellow #6, flavors, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sodium saccharin, sorbitol, sucralose

Questions or comments?

1-877-507-6516

Principal display panel

24/7 life

by **7-ELEVEN™**

Cold&Flu

DayTime Relief

For Ages 6 Years & Over

Acetaminophen / Pain Reliever/Fever Reducer

Dextromethorphan HBr / Cough Suppressant

Phenylephrine HCl / Nasal Decongestant

Relieves: Minor Aches & Pains, Sore Throat, Fever, Headache, Nasal Congestion, Cough

compare to Vicks®

DayQuil® Cold & Flu

active ingredients*

Menthol

Flavor

**6 FL OZ
(177 mL)**

**TAMPER EVIDENT: DO NOT USE IF PRINTED
NECK WRAP IS BROKEN OR MISSING**

*This product is not
manufactured or
distributed by The Procter
& Gamble Company,
owner of the registered
trademark Vicks®
DayQuil® Cold & Flu.

50844 REV0323A01245

65154L-SE-02-25-1

DISTRIBUTED BY
7-ELEVEN, INC.
IRVING, TX 75063
WWW.7-ELEVEN.COM

**Satisfaction Guaranteed
1-800-255-0711**

PARENTS:

Learn about teen medicine abuse
www.StopMedicineAbuse.org

Drug Facts (continued)

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if the user has

■ difficulty in urination due to enlargement of the prostate gland

■ cough that occurs with too much phlegm (mucus)

■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema

■ heart disease ■ diabetes ■ thyroid disease

■ liver disease ■ high blood pressure

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

■ nervousness, dizziness, or sleeplessness occur

■ fever gets worse or lasts more than 3 days

■ redness or swelling is present ■ new symptoms occur

■ pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)

■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed

■ do not exceed 4 doses per 24 hours ■ mL = milliliter

■ only use the dose cup provided

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children under 6 years	do not use

Other information

■ each 15 mL contains: sodium 13 mg

■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package

Inactive ingredients anhydrous citric acid, FD&C yellow #6, flavors, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sodium saccharin, sorbitol, sucralose

Questions or comments? 1-877-607-6516

STOP PEELING

Drug Facts (continued)

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if ■ adult takes more than 4,000 mg of acetaminophen in 24 hours

■ child takes more than 5 doses in 24 hours

■ taken with other drugs containing acetaminophen

■ adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

DISTRIBUTED BY
7-ELEVEN, INC.
IRVING, TX 75063
www.7-ELEVEN.COM

50844 REV0323A01245
65154L-SE-02-25-11

Satisfaction Guaranteed
1-800-255-0711

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

PEEL HERE FOR MORE DRUG FACTS

24/7 life by 7-ELEVEN

Cold & Flu For Ages 6 Years & Over

DayTime Relief

Acetaminophen / Pain Reliever/Fever Reducer
Dextromethorphan HBr / Cough Suppressant
Phenylephrine HCl / Nasal Decongestant

Relieves: Minor Aches & Pains, Sore Throat, Fever, Headache, Nasal Congestion, Cough

6 FL OZ (177 mL)

Menthol Flavor

compare to Vicks® DayQuil® Cold & Flu active ingredients*

No Print/No Varnish Lot & EXP

Drug Facts TAMPER EVIDENT: DO NOT USE IF PRINTED NECK WRAP IS BROKEN OR MISSING

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves common cold and flu symptoms:
- sore throat ■ nasal congestion
- headache ■ minor aches and pains
- cough due to minor throat and bronchial irritation
- temporarily reduces fever

*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademark Vicks® DayQuil® Cold & Flu.

0 52548 59553 1

Convenience Valet 44-012

COLD AND FLU DAYTIME RELIEF			
acetaminophen, dextromethorphan hbr, phenylephrine hcl solution			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50844-251
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL	
Inactive Ingredients			
Ingredient Name	Strength		
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			

FD&C YELLOW NO. 6 (UNII: H77VEI93A8)
GLYCERIN (UNII: PDC6A3C0OX)
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)
WATER (UNII: 059QF0KO0R)
SODIUM BENZOATE (UNII: OJ245FE5EU)
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)
SODIUM METABISULFITE (UNII: 4VON5FNS3C)
SACCHARIN SODIUM (UNII: SB8ZUX40TY)
SORBITOL (UNII: 506T60A25R)
SUCRALOSE (UNII: 96K6UQ3ZD4)

Product Characteristics

Color	orange	Score	
Shape		Size	
Flavor	MENTHOL	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50844-251-45	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/20/2019	
2	NDC:50844-251-02	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/20/2019	04/30/2026

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/20/2019	

Labeler - L.N.K. International, Inc. (038154464)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	manufacture(50844-251) , pack(50844-251)

Revised: 9/2025

L.N.K. International, Inc.