

ASPIRIN 325MG- aspirin tablet
Innovida Pharmaceutique Corporation

Active Ingredient

Aspirin 325 mg(NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- temporarily relieves minor aches and pains due to:
- headache
- menstrual pain
- a cold
- toothache
- muscle pain
- minor pain of arthritis
- or as recommended by your doctor

Warnings

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product.

When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product

- take more or for a longer time than directed

Do not use if

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma you
- are taking a prescription drug for diabetes, arthritis, or gout

Stop use and ask a doctor if

- an allergic reaction occurs.
- Seek medical help right away
- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - feel faint
 - have bloody or black stools
 - have stomach pain that does not get better
 - ringing in the ears or a loss of hearing occurs
 - new symptoms occur · pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 to 2 tablets every 4 hours not to exceed 12 tablets in 24 hours unless directed by a doctor ·
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive Ingredients

black iron oxide, cellulose, colloidal anhydrous silica, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, polydextrose, polyethylene glycol, polyvinyl acetate phthalate, propylene glycol, shellac wax, simethicone, sodium alginate, sodium bicarbonate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate

Questions or Comments? 1-888-462-4166

Product label

NDC: 71800-041-01

INNVIDA
PHARMACEUTIQUE CORPORATION

Aspirin

Pain Reliever (NSAID),
325 mg

Enteric Coated Tablet

100 Tablets

Tamper Evident: Do not use if imprinted seal under cap is missing or broken.

Drug Facts

Active ingredient (in each tablet) **Purpose**
Aspirin 325 mg(NSAID)*Pain reliever
*nonsteroidal anti-inflammatory drug

Uses • temporarily relieves minor aches and pains due to:
• headache • menstrual pain • a cold • toothache • muscle pain
• minor pain of arthritis • or as recommended by your doctor

Warnings

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
• are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed

Do not use if • if you are allergic to aspirin or any other pain reliever/fever reducer • if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you are taking a diuretic • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you have asthma you • are taking a prescription drug for diabetes, arthritis, or gout

Mfd. for:
Innovida Pharmaceutique
Corporation, Phoenix, AZ 85040

Tel.: 888-462-4166



PEEL HERE FOR MORE DRUG FACTS

Questions or Comments? 1-888-462-4166

Inactive ingredients: Black iron oxide, cellulose, colloidal amorphous silica, corn starch, D&C yellow #10 aluminum lake, F&D&C yellow #6 aluminum lake, hypromellose, polyethylene glycol, polyethylene glycol, polyvinyl acetate phthalate, polyethylene glycol, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate, shellac wax, simethicone, sodium alginate, sodium bicarbonate,

Other information • store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) • see end flap for expiration date and lot number

Directions • do not take more than directed • drink a full glass of water with each dose • adults and children 12 years and over take 1 to 2 tablets every 4 hours not to exceed 12 tablets in 24 hours unless directed by a doctor • children under 12 years: do not use unless directed by a doctor

In case of overdose, get medical help or contact a Poison Control Center right away.

Keep out of reach of children.

because it may cause problems in the unborn child or complications during delivery.
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless directed to do so by a doctor.

Stop use and ask a doctor if • an allergic reaction occurs. Seek medical help right away • you experience any of the following signs of stomach bleeding: • vomit blood • feel faint • have bloody or black stools • have stomach pain that does not get better • ringing in the ears or a loss of hearing occurs • new symptoms occur • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • redness or swelling is present. These could be signs of a serious condition.

Drug Facts (continued)

aspirin tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:71800-041
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ACETYLSALICYLIC ACID - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
POWDERED CELLULOSE (UNII: SMD1X3X09M)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
FD&C YELLOW NO. 6 ALUMINUM LAKE (UNII: GYP6Z2JR6Q)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
POLYVINYL ACETATE PHTHALATE (UNII: 58QVG85GW3)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B710)	
DIMETHICONE (UNII: 92RU3N3Y10)	
WATER (UNII: 059QF0KO0R)	
SODIUM ALGINATE (UNII: C269C4G2ZQ)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71800-041-	100 in 1 BOTTLE; Type 0: Not a Combination	02/10/2025	

01	Product	02/10/2025	
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/10/2025	

Labeler - Innovida Pharmaceutique Corporation (080892908)

Revised: 2/2025

Innovida Pharmaceutique Corporation