

**FLU HBP MAXIMUM STRENGTH- acetaminophen, chlorpheniramine maleate,
dextromethorphan hbr tablet, film coated
Meijer Distribution Inc**

Meijer 44-738

Active ingredients (in each caplet)

Acetaminophen 325 mg
Chlorpheniramine maleate 2 mg
Dextromethorphan HBr 10 mg

Purpose

Pain reliever/fever reducer
Antihistamine
Cough suppressant

Uses

- temporarily relieves these common cold and flu symptoms:
 - minor aches and pains
 - sore throat
 - sneezing
 - headache
 - runny nose
 - cough
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a

doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- avoid alcoholic beverages
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- excitability may occur, especially in children
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- adults and children 12 years and over: take 2 caplets every 4 hours while symptoms persist. Do not take more than 10 caplets in 24 hours.
- children under 12 years: ask a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- protect from excessive moisture
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, crospovidone, FD&C red #40 aluminum lake, lecithin, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments?

1-800-426-9391

Principal display panel

meijer®

NDC 41250-638-09

Compare to Coricidin® HBP

Maximum Strength

Multi-Symptom Flu

active ingredients*

MAXIMUM STRENGTH

flu HBP

Acetaminophen 325 mg

Chlorpheniramine maleate 2 mg

Dextromethorphan HBr 10 mg

Pain Reliever/Fever Reducer

Antihistamine

Cough Suppressant

For People with High Blood Pressure

Multi-Symptom

Relieves: Body Aches, Pains & Headache,

Fever, Cough, Runny Nose, Sore Throat, Sneezing

20 Caplets

Actual
Size

**TAMPER EVIDENT: DO NOT USE IF PACKAGE
IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS OF TAMPERING**

**DIST. BY MEIJER
DISTRIBUTION, INC.
GRAND RAPIDS, MI 49544
www.meijer.com**

*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Coricidin® HBP Maximum Strength Multi-Symptom Flu. 50844 REV1019A73809

OUR QUALITY
GUARANTEE
*The Meijer
Family*
WWW.MEIJER.COM/SATISFACTION

*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Coricidin® HBP Maximum Strength Multi-Symptom Flu. 50844 REV1019A73809

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

DIST. BY MEIJER
DISTRIBUTION, INC.
GRAND RAPIDS, MI 49544
www.meijer.com

OUR QUALITY
GUARANTEE
*The Meijer
Family*
WWW.MEIJER.COM/SATISFACTION

NDC 41250-638-09

meijer

Compare to Coricidin® HBP
Maximum Strength
Multi-Symptom Flu
active ingredients*

**MAXIMUM STRENGTH
flu HBP**

**Acetaminophen 325 mg
Chlorpheniramine maleate 2 mg
Dextromethorphan HBr 10 mg**

Pain Reliever/Fever Reducer
Antihistamine
Cough Suppressant

For People with High Blood Pressure
Multi-Symptom

Relieves: Body Aches, Pains & Headache,
Fever, Cough, Runny Nose, Sore Throat, Sneezing

20 Caplets  Actual Size

Drug Facts (continued)
microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Drug Facts (continued)
Store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F).
Protect from excessive moisture.
See end flap for expiration date and lot number.

Inactive ingredients corn starch, croscopollose, FD&C red #40 aluminum lake, lecithin, magnesium stearate.

Drug Facts (continued)
Temporarily relieves these common cold and flu symptoms:
■ sore throat ■ sore throat ■ sore throat
■ minor aches and pains ■ headache ■ headache
■ sneezing ■ sneezing ■ sneezing
■ temporarily reduces fever ■ temporarily reduces fever ■ temporarily reduces fever

Drug Facts (continued)
Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ blisters ■ rash ■ skin reddening
If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
■ difficulty in urination due to enlargement of the prostate gland
■ liver disease ■ glaucoma
■ cough that occurs with too much phlegm (mucus)
■ a breathing problem or persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

Drug Facts (continued)
Ask a doctor or pharmacist before use if you are
■ taking sedatives or tranquilizers
■ taking the blood thinning drug warfarin

When using this product
■ avoid alcoholic beverages
■ marked drowsiness may occur
■ alcohol, sedatives, and tranquilizers may increase drowsiness
■ excitability may occur, especially in children
■ use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if
■ pain or cough gets worse or lasts more than 7 days
■ new symptoms occur
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

if pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
■ do not take more than directed
■ adults and children 12 years and over: take 2 caplets every 4 hours while symptoms persist. Do not take more than 10 caplets in 24 hours.
■ children under 12 years: ask a doctor

Other information
■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

Drug Facts (continued)
Uses
Temporarily relieves these common cold and flu symptoms:
■ sore throat ■ sore throat ■ sore throat
■ minor aches and pains ■ headache ■ headache
■ sneezing ■ sneezing ■ sneezing
■ temporarily reduces fever ■ temporarily reduces fever ■ temporarily reduces fever

Drug Facts (continued)
Active ingredients (in each caplet)
Acetaminophen 325 mg Pain reliever/fever reducer
Chlorpheniramine maleate 2 mg Antihistamine
Dextromethorphan HBr 10 mg Cough suppressant

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

PID 4338255

7 19283 64071 0

B-1214-738-09-R
REV1019A73809

No Print / No Varnish Area
Lot # and Exp. Info

Meijer 44-738

FLU HBP MAXIMUM STRENGTH

acetaminophen, chlorpheniramine maleate, dextromethorphan hbr tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-638
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
FD&C RED NO. 40 ALUMINUM LAKE (UNII: 6T47AS764T)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	red	Score	no score
Shape	OVAL	Size	16mm
Flavor		Imprint Code	44;738
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-638-09	2 in 1 CARTON	02/28/2020	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination		

Product			
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	02/28/2020	

Labeler - Meijer Distribution Inc (006959555)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(41250-638) , pack(41250-638)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(41250-638)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(41250-638)

Revised: 8/2025

Meijer Distribution Inc