

BACLOFEN- baclofen tablet
Method Pharmaceuticals LLC


Baclofen Tablets, USP

Rx only

DESCRIPTION

Baclofen, USP is a muscle relaxant and antispastic.

Its chemical name is 4-amino-3-(4-chlorophenyl)-butanoic acid, and its structural formula is



$\text{C}_{10}\text{H}_{12}\text{ClNO}_2$ M.W. 213.66

Baclofen, USP is a white to off-white odorless or practically odorless crystalline powder. It is slightly soluble in water, very slightly soluble in methanol and insoluble in chloroform.

Each tablet, for oral administration, contains 5 mg or 10 mg or 20 mg baclofen, USP. In addition, each tablet contains the following inactive ingredients: croscarmellose sodium, lactose monohydrate, microcrystalline cellulose, zinc stearate.

CLINICAL PHARMACOLOGY

The precise mechanism of action of baclofen is not fully known. Baclofen is capable of inhibiting both monosynaptic and polysynaptic reflexes at the spinal level, possibly by hyperpolarization of afferent terminals, although actions at supraspinal sites may also occur and contribute to its clinical effect. Although baclofen is an analog of the putative inhibitory neurotransmitter gamma-aminobutyric acid (GABA), there is no conclusive evidence that actions on GABA systems are involved in the production of its clinical effects. In studies with animals baclofen has been shown to have general CNS depressant properties as indicated by the production of sedation with tolerance, somnolence, ataxia, and respiratory and cardiovascular depression. Baclofen is rapidly and extensively absorbed and eliminated. Absorption may be dose-dependent, being reduced with increasing doses. Baclofen is excreted primarily by the kidney in unchanged form and there is relatively large intersubject variation in absorption and/or

elimination.

INDICATIONS AND USAGE

Baclofen tablets USP are useful for the alleviation of signs and symptoms of spasticity resulting from multiple sclerosis, particularly for the relief of flexor spasms and concomitant pain, clonus, and muscular rigidity.

Patients should have reversible spasticity so that baclofen treatment will aid in restoring residual function. Baclofen tablets USP may also be of some value in patients with spinal cord injuries and other spinal cord diseases.

Baclofen tablets USP are not indicated in the treatment of skeletal muscle spasm resulting from rheumatic disorders.

The efficacy of baclofen in stroke, cerebral palsy, and Parkinson's disease has not been established and, therefore, it is not recommended for these conditions.

CONTRAINDICATIONS

Hypersensitivity to baclofen.

WARNINGS

a. Neonatal Withdrawal Symptoms:

Withdrawal symptoms have been reported starting hours to days after delivery in neonates whose mothers were treated with oral baclofen throughout pregnancy. The symptoms of withdrawal in these infants have included increased muscle tone, tremor, jitteriness, and seizure. If the potential benefit justifies the potential risk to the fetus and oral baclofen is continued during pregnancy, gradually reduce the dose and discontinue baclofen before delivery. If slow withdrawal is not feasible, advise the parents or caregivers of the potential for neonatal withdrawal.

b. Abrupt Drug Withdrawal:

Hallucinations and seizures have occurred on abrupt withdrawal of baclofen. Therefore, except for serious adverse reactions, the dose should be reduced slowly when the drug is discontinued.

c. Impaired Renal Function:

Because baclofen is primarily excreted unchanged through the kidneys, it should be given with caution, and it may be necessary to reduce the dosage.

d. Stroke:

Baclofen has not significantly benefited patients with stroke. These patients have also shown poor tolerability to the drug.

e. Pregnancy:

Baclofen has been shown to increase the incidence of omphaloceles (ventral hernias) in fetuses of rats given approximately 13 times the maximum dose recommended for

human use, at a dose which caused significant reductions in food intake and weight gain in dams. This abnormality was not seen in mice or rabbits.

There was also an increased incidence of incomplete sternebral ossification in fetuses of rats given approximately 13 times the maximum recommended human dose, and an increased incidence of unossified phalangeal nuclei of forelimbs and hindlimbs in fetuses of rabbits given approximately 7 times the maximum recommended human dose. In mice, no teratogenic effects were observed, although reductions in mean fetal weight with consequent delays in skeletal ossification were present when dams were given 17 and 34 times the human daily dose. There are no studies in pregnant women. Baclofen should be used during pregnancy only if the benefit clearly justifies the potential risk to the fetus.

PRECAUTIONS

Because of the possibility of sedation, patients should be cautioned regarding the operation of automobiles or other dangerous machinery, and activities made hazardous by decreased alertness. Patients should also be cautioned that the central nervous system effects of baclofen may be additive to those of alcohol and other CNS depressants.

Baclofen should be used with caution where spasticity is utilized to sustain upright posture and balance in locomotion or whenever spasticity is utilized to obtain increased function.

In patients with epilepsy, the clinical state and electroencephalogram should be monitored at regular intervals, since deterioration in seizure control and EEG have been reported occasionally in patients taking baclofen.

It is not known whether this drug is excreted in human milk. As a general rule, nursing should not be undertaken while a patient is on a drug since many drugs are excreted in human milk.

A dose-related increase in incidence of ovarian cysts and a less marked increase in enlarged and/or hemorrhagic adrenal glands was observed in female rats treated chronically with baclofen.

Ovarian cysts have been found by palpation in about 4% of the multiple sclerosis patients that were treated with baclofen for up to one year. In most cases these cysts disappeared spontaneously while patients continued to receive the drug. Ovarian cysts are estimated to occur spontaneously in approximately 1% to 5% of the normal female population.

Pediatric Use

Safety and effectiveness in pediatric patients below the age of 12 years have not been established.

ADVERSE REACTIONS

The most common is transient drowsiness (10 to 63%). In one controlled study of 175 patients, transient drowsiness was observed in 63% of those receiving baclofen compared to 36% of those in the placebo group. Other common adverse reactions are

dizziness (5 to 15%), weakness (5 to 15%) and fatigue (2 to 4%).

Others reported:

Neuropsychiatric:

Confusion (1 to 11%), headache (4 to 8%), insomnia (2 to 7%); and, rarely, euphoria, excitement, depression, hallucinations, paresthesia, muscle pain, tinnitus, slurred speech, coordination disorder, tremor, rigidity, dystonia, ataxia, blurred vision, nystagmus, strabismus, miosis, mydriasis, diplopia, dysarthria, epileptic seizure.

Cardiovascular:

Hypotension (0 to 9%). Rare instances of dyspnea, palpitation, chest pain, syncope.

Gastrointestinal:

Nausea (4 to 12%), constipation (2 to 6%); and, rarely, dry mouth, anorexia, taste disorder, abdominal pain, vomiting, diarrhea, and positive test for occult blood in stool.

Genitourinary:

Urinary frequency (2 to 6%); and, rarely, enuresis, urinary retention, dysuria, impotence, inability to ejaculate, nocturia, hematuria.

Other:

Instances of rash, pruritus, ankle edema, excessive perspiration, weight gain, nasal congestion.

Some of the CNS and genitourinary symptoms may be related to the underlying disease rather than to drug therapy. The following laboratory tests have been found to be abnormal in a few patients receiving baclofen: increased SGOT, elevated alkaline phosphatase, and elevation of blood sugar.

OVERDOSAGE

Signs and Symptoms:

Vomiting, muscular hypotonia, drowsiness, accommodation disorders, coma, respiratory depression and seizures.

Treatment:

In the alert patient, empty the stomach promptly by induced emesis followed by lavage. In the obtunded patient, secure the airway with a cuffed endotracheal tube before beginning lavage (do not induce emesis). Maintain adequate respiratory exchange, do not use respiratory stimulants.

DOSAGE AND ADMINISTRATION

The determination of optimal dosage requires individual titration. Start therapy at a low dosage and increase gradually until optimum effect is achieved (usually between 40 to 80 mg daily).

The following dosage titration schedule is suggested:

5 mg t.i.d. for 3 days
10 mg t.i.d. for 3 days
15 mg t.i.d. for 3 days
20 mg t.i.d. for 3 days

Thereafter additional increases may be necessary but the total daily dose should not exceed a maximum of 80 mg daily (20 mg q.i.d.).

The lowest dose compatible with an optimal response is recommended. If benefits are not evident after a reasonable trial period, patients should be slowly withdrawn from the drug (see **WARNINGS**, Abrupt Drug Withdrawal).

HOW SUPPLIED

Baclofen Tablets USP, 5 mg – white to off-white, round, flat tablet with bevel edge, debossed “167”

on one side and “l” on the other side.

Bottles of 100..... NDC 58657-730-01

Bottles of 1000..... NDC 58657-730-10

Carton of 50 tablets (10 tablets per blister pack x 5) NDC 58657-930-05

Carton of 100 tablets (10 tablets per blister pack x 10)..... NDC 58657-930-01

Baclofen Tablets USP, 10 mg – white to off-white, round, flat tablet with bevel edge, debossed “l

114” on one side and scored on the other side.

Bottles of 100..... NDC 58657-731-01

Bottles of 500..... NDC 58657-731-50

Bottles of 1000..... NDC 58657-731-10

Carton of 50 tablets (10 tablets per blister pack x 5) NDC 58657-931-05

Carton of 100 tablets (10 tablets per blister pack x 10)..... NDC 58657-931-01

Baclofen Tablets USP, 20 mg – white to off-white, round, flat tablet with bevel edge, debossed “l

115” on one side and scored on the other side.

Bottles of 100..... NDC 58657-732-01

Bottles of 500..... NDC 58657-732-50

Bottles of 1000..... NDC 58657-732-10

Carton of 50 tablets (10 tablets per blister pack x 5) NDC 58657-

932-05

Carton of 100 tablets (10 tablets per blister pack x 10)..... NDC 58657-932-01

PHARMACIST: Dispense in a well-closed container as defined in the USP, with a child-resistant closure (as required).

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Distributed by:

Method Pharmaceuticals, LLC

Southlake, TX 76092

Rev. 06/2025

Baclofen Tablets USP, 5 mg- 100 Tablets

Rx Only

NDC 58657- 0730 -01	
Baclofen Tablets, USP	
5 mg	
Rx Only 100 Tablets	
Each tablet contains 5 mg baclofen, USP. Usual Dosage: See package insert for full prescribing information. Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. Dispense in a well-closed container as defined in the USP, with a child-resistant closure (as required). KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.	
Rev.03/2024 Dist. by: Method Pharmaceuticals, LLC Southlake, TX 76092	
 N 3 58657 73001 7	
1.0"L x 1.5"W	
NO VARNISH	

Baclofen Tablets USP, 5 mg- 1000 Tablets

Rx Only

NDC 58657-0730-10

Baclofen Tablets, USP

5 mg

Rx Only

1000 Tablets



Each tablet contains 5 mg baclofen, USP.

Usual Dosage: See package insert for full prescribing information.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Dispense in a well-closed container as defined in the USP, with a child-resistant closure (as required).

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Rev.03/2024

Dist. by: **Method Pharmaceuticals, LLC**
Southlake, TX 76092



1.0"L x 2.0"W

NO VARNISH

Baclofen Tablets USP, 10 mg- 100 Tablets

Rx Only

NDC 58657-0731-01

Baclofen Tablets, USP

10 mg

Rx Only

100 Tablets



Each tablet contains 10 mg baclofen, USP.

Usual Dosage: See package insert for full prescribing information.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Dispense in a well-closed container as defined in the USP, with a child-resistant closure (as required).

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Rev.03/2024

Dist. by: **Method Pharmaceuticals, LLC**
Southlake, TX 76092



1.0"L x 1.5"W

NO VARNISH

Baclofen Tablets USP, 10 mg- 500 Tablets

Rx Only

NDC 58657-0731-50

Baclofen Tablets, USP

10 mg

Rx Only
500 Tablets



Each tablet contains 10 mg baclofen, USP.

Usual Dosage: See package insert for full prescribing information.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Dispense in a well-closed container as defined in the USP, with a child-resistant closure (as required).

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Rev.03/2024

Dist. by: **Method Pharmaceuticals, LLC**
Southlake, TX 76092



1.0"L x 2.0"W

NO VARNISH

Baclofen Tablets USP, 10 mg- 1000 Tablets

Rx Only

NDC 58657-0731-10

Baclofen Tablets, USP

10 mg

Rx Only
1000 Tablets



Each tablet contains 10 mg baclofen, USP.

Usual Dosage: See package insert for full prescribing information.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Dispense in a well-closed container as defined in the USP, with a child-resistant closure (as required).

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Rev.03/2024

Dist. by: **Method Pharmaceuticals, LLC**
Southlake, TX 76092



1.2"L x 2.5"W

NO VARNISH

Baclofen Tablets USP, 20 mg- 100 Tablets

Rx only

NDC 58657-0732-01

Baclofen Tablets, USP

20 mg

Rx Only
100 Tablets



Each tablet contains 20 mg baclofen, USP.

Usual Dosage: See package insert for full prescribing information.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Dispense in a well-closed container as defined in the USP, with a child-resistant closure (as required).

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Rev.03/2024

Dist. by: **Method Pharmaceuticals, LLC**
Southlake, TX 76092



1.0"L x 2.0"W

NO VARNISH

Baclofen Tablets USP, 20 mg- 500 Tablets

Rx only

NDC 58657-0732-50

Baclofen Tablets, USP

20 mg

Rx Only
500 Tablets



Each tablet contains 20 mg baclofen, USP.

Usual Dosage: See package insert for full prescribing information.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Dispense in a well-closed container as defined in the USP, with a child-resistant closure (as required).

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Rev.03/2024

Dist. by: **Method Pharmaceuticals, LLC**
Southlake, TX 76092



1.2"L x 2.5"W

NO VARNISH

Baclofen Tablets USP, 20 mg- 1000 Tablets

Rx only

NDC 58657-0732-10

Baclofen Tablets, USP

20 mg

Rx Only

1000 Tablets



Each tablet contains 20 mg baclofen, USP.

Usual Dosage: See package insert for full prescribing information.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Dispense in a well-closed container as defined in the USP, with a child-resistant closure (as required).

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Rev.03/2024

Dist. by: **Method Pharmaceuticals, LLC**
Southlake, TX 76092



1.5"L x 3.0"W

NO VARNISH

Baclofen Tablets USP, 5 mg- 50 Tablets

Rx Only



NDC 58657-930-05
Unit Dose
Baclofen Tablets, USP

5 mg per tablet

Rx Only
50 Tablets (5 x 10 Unit Dose)



NDC 58657-930-05
Unit Dose
Baclofen Tablets, USP

5 mg per tablet

Rx Only
50 Tablets (5 x 10 Unit Dose)

Each tablet contains:
Baclofen, USP..... 5 mg

Usual Dosage: See package insert for full
prescribing information.

Store at 20° to 25°C (68° to 77°F).
[See USP Controlled Room Temperature].

Dispense in a well-closed container as defined in the USP,
with a child-resistant closure (as required).

Keep this and all medications out of
the reach of children. This unit
dose package is not child resistant and is intended for
Institutional Use Only.

Rev: 06/2025

Mfg by: Innogenix, LLC.
Amityville, NY 11701

Dist. by: Method Pharmaceuticals LLC
Southlake, TX 76092



Baclofen Tablets USP, 5 mg- 100 Tablets
Rx Only



NDC 58657-930-01

Unit Dose

Baclofen Tablets, USP

5 mg per tablet

Rx Only

100 Tablets (10 x 10 Unit Dose)



NDC 58657-930-01

Unit Dose

Baclofen Tablets, USP

5 mg per tablet

Rx Only

100 Tablets (10 x 10 Unit Dose)

Each tablet contains:

Baclofen, USP..... 5 mg

Usual Dosage: See package insert for full
prescribing information.

Store at 20° to 25°C (68° to 77°F).

[See USP Controlled Room Temperature].

Dispense in a well-closed container as defined in the USP,
with a child-resistant closure (as required).

Keep this and all medications out of
the reach of children. This unit
dose package is not child resistant and is intended for
Institutional Use Only.

Rev: 04/2025

Mfg by: Innogenix, LLC.

Amityville, NY 11701

Dist. by: Method Pharmaceuticals LLC

Southlake, TX 76092



Baclofen Tablets USP, 10 mg- 50 Tablets
Rx Only



NDC 58657-931-05

Unit Dose

Baclofen Tablets, USP

10 mg per tablet

Rx Only

50 Tablets (5 x 10 Unit Dose)



NDC 58657-931-05

Unit Dose

Baclofen Tablets, USP

10 mg per tablet

Rx Only

50 Tablets (5 x 10 Unit Dose)

Each tablet contains:

Baclofen, USP..... 10 mg

Usual Dosage: See package insert for full
prescribing information.

Store at 20° to 25°C (68° to 77°F).

[See USP Controlled Room Temperature].

Dispense in a well-closed container as defined in the USP,
with a child-resistant closure (as required).

Keep this and all medications out of
the reach of children. This unit
dose package is not child resistant and is intended for
Institutional Use Only.

Rev: 06/2025

Mfg by: Innogenix, LLC.
Amityville, NY 11701

Dist. by: Method Pharmaceuticals LLC
Southlake, TX 76092



Baclofen Tablets USP, 10 mg- 100 Tablets

Rx Only



NDC 58657-931-01

Unit Dose

Baclofen Tablets, USP

10 mg per tablet

Rx Only

100 Tablets (10 x 10 Unit Dose)



NDC 58657-931-01

Unit Dose

Baclofen Tablets, USP

10 mg per tablet

Rx Only

100 Tablets (10 x 10 Unit Dose)

Each tablet contains:

Baclofen, USP..... 10 mg

Usual Dosage: See package insert for full
prescribing information.

Store at 20° to 25°C (68° to 77°F).
[See USP Controlled Room Temperature].

Dispense in a well-closed container as defined in the USP,
with a child-resistant closure (as required).

Keep this and all medications out of
the reach of children. This unit
dose package is not child resistant and is Intended for
Institutional Use Only.

Rev: 06/2025

Mfg by: Innogenix, LLC.
Amityville, NY 11701

Dist. by: Method Pharmaceuticals LLC
Southlake, TX 76092



Baclofen Tablets USP, 20 mg- 50 Tablets
Rx Only



NDC 58657-932-05

Unit Dose

Baclofen Tablets, USP

20 mg per tablet

Rx Only

50 Tablets (5 x 10 Unit Dose)



NDC 58657-932-05

Unit Dose

Baclofen Tablets, USP

20 mg per tablet

Rx Only

50 Tablets (5 x 10 Unit Dose)

Each tablet contains:

Baclofen, USP..... 20 mg

Usual Dosage: See package insert for full
prescribing information.

Store at 20° to 25°C (68° to 77°F).

[See USP Controlled Room Temperature].

Dispense in a well-closed container as defined in the USP,
with a child-resistant closure (as required).

Keep this and all medications out of
the reach of children. This unit
dose package is not child resistant and is intended for
Institutional Use Only.

Rev: 06/2025

Mfg by: Innogenix, LLC.
Amityville, NY 11701

Dist. by: Method Pharmaceuticals LLC
Southlake, TX 76092



Baclofen Tablets USP, 20 mg- 100 Tablets

Rx Only



NDC 58657-932-01

Unit Dose

Baclofen Tablets, USP

20 mg per tablet

Rx Only

100 Tablets (10 x 10 Unit Dose)



NDC 58657-932-01

Unit Dose

Baclofen Tablets, USP

20 mg per tablet

Rx Only

100 Tablets (10 x 10 Unit Dose)

Each tablet contains:

Baclofen, USP..... 20 mg

Usual Dosage: See package insert for full
prescribing information.

Store at 20° to 25°C (68° to 77°F).
[See USP Controlled Room Temperature].

Dispense in a well-closed container as defined in the USP,
with a child-resistant closure (as required).

Keep this and all medications out of
the reach of children. This unit
dose package is not child resistant and is intended for
Institutional Use Only.

Rev: 04/2025

Mfg by: Innogenix, LLC.
Amityville, NY 11701

Dist. by: Method Pharmaceuticals LLC
Southlake, TX 76092



BACLOFEN

baclofen tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:58657-932
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
BACLOFEN (UNII: H789N3FKE8) (BACLOFEN - UNII:H789N3FKE8)		BACLOFEN	20 mg	
Inactive Ingredients				
Ingredient Name			Strength	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
ZINC STEARATE (UNII: H92E6QA4FV)				
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)				
Product Characteristics				
Color	white (off-white)	Score	2 pieces	
Shape	ROUND	Size	11mm	
Flavor		Imprint Code	l115	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58657-932-05	5 in 1 CARTON	06/01/2025	
1	NDC:58657-932-10	10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:58657-932-01	10 in 1 CARTON	06/01/2025	
2	NDC:58657-932-10	10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA212378	06/15/2024	

BACLOFEN

baclofen tablet

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:58657-730
Route of Administration	ORAL		
Active Ingredient/Active Moiety			

Ingredient Name		Basis of Strength		Strength
BACLOFEN (UNII: H789N3FKE8) (BACLOFEN - UNII:H789N3FKE8)		BACLOFEN		5 mg
Inactive Ingredients				
Ingredient Name			Strength	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)				
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
ZINC STEARATE (UNII: H92E6QA4FV)				
Product Characteristics				
Color	white (off-white)		Score	no score
Shape	ROUND		Size	6mm
Flavor			Imprint Code	167;l
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58657-730-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	06/15/2024	
2	NDC:58657-730-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	06/15/2024	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA212378	06/15/2024	

BACLOFEN			
baclofen tablet			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:58657-731
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
BACLOFEN (UNII: H789N3FKE8) (BACLOFEN - UNII:H789N3FKE8)		BACLOFEN	10 mg

Inactive Ingredients				
Ingredient Name			Strength	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)				
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
ZINC STEARATE (UNII: H92E6QA4FV)				
Product Characteristics				
Color	white (off-white)	Score	2 pieces	
Shape	ROUND	Size	9mm	
Flavor		Imprint Code	I114	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58657-731-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	06/15/2024	
2	NDC:58657-731-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	06/15/2024	
3	NDC:58657-731-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	06/15/2024	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA212378	06/15/2024	

BACLOFEN			
baclofen tablet			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:58657-732
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
BACLOFEN (UNII: H789N3FKE8) (BACLOFEN - UNII:H789N3FKE8)		BACLOFEN	20 mg
Inactive Ingredients			
Ingredient Name			Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)			

LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
ZINC STEARATE (UNII: H92E6QA4FV)				
Product Characteristics				
Color	white (off-white)		Score	2 pieces
Shape	ROUND		Size	11mm
Flavor			Imprint Code	l115
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58657-732-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	06/15/2024	
2	NDC:58657-732-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	06/15/2024	
3	NDC:58657-732-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	06/15/2024	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA212378	06/15/2024	

BACLOFEN			
baclofen tablet			
Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:58657-931
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
BACLOFEN (UNII: H789N3FKE8) (BACLOFEN - UNII:H789N3FKE8)		BACLOFEN	10 mg
Inactive Ingredients			
Ingredient Name			Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)			
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)			
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)			
ZINC STEARATE (UNII: H92E6QA4FV)			

Product Characteristics

Color	white (off-white)	Score	2 pieces
Shape	ROUND	Size	9mm
Flavor		Imprint Code	l114
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58657-931-05	5 in 1 CARTON	06/01/2025	
1	NDC:58657-931-10	10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:58657-931-01	10 in 1 CARTON	06/01/2025	
2	NDC:58657-931-10	10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA212378	06/01/2025	

BACLOFEN

baclofen tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:58657-930
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BACLOFEN (UNII: H789N3FKE8) (BACLOFEN - UNII:H789N3FKE8)	BACLOFEN	5 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
ZINC STEARATE (UNII: H92E6QA4FV)	

Product Characteristics

Color	white (off-white)	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	167;l
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58657-930-05	5 in 1 CARTON	06/01/2025	
1	NDC:58657-930-10	10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:58657-930-01	10 in 1 CARTON	06/01/2025	
2	NDC:58657-930-10	10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA212378	06/15/2024	

Labeler - Method Pharmaceuticals LLC (060216698)

Revised: 7/2025

Method Pharmaceuticals LLC