

STOMACH RELIEF REGULAR STRENGTH- bismuth subsalicylate liquid

CVS Pharmacy

Drug Facts

Active ingredient (in each 30 mL dose)

Bismuth subsalicylate 525 mg

Purpose

Upset stomach reliever/Antidiarrheal

Uses

relieves

- diarrhea
- travelers' diarrhea
- upset stomach due to overindulgence in food and drink, including:
 - heartburn
 - gas
 - indigestion
 - nausea
 - fullness
 - belching

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are:

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- diabetes
- gout
- arthritis
- anticoagulation (thinning the blood)

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- mL = milliliter
- shake well before using
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- adults and children 12 years and over
 - 30 mL (1 dose) every 1/2 hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
 - 30 mL (1dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
 - do not exceed 8 doses (240 mL) in 24 hours
 - use until diarrhea stops but not more than 2 days
- children under 12 years of age: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- **each 30 mL contains:** potassium 8 mg, sodium 9 mg
- **each 30 mL contains:** salicylate 206 mg
- low sodium
- keep tightly closed
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

Inactive ingredients

benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to Pepto-Bismol® active ingredient*

Original

STOMACH RELIEF

Bismuth Subsalicylate 525 mg per 30 mL

Upset stomach reliever/

Antidiarrheal

Relieves 5 symptoms: Heartburn, Indigestion, Upset stomach, Nausea, Diarrhea

Original Flavor | **SHAKE WELL**

FL OZ (mL)

*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

Distributed by: CVS Pharmacy, Inc.

One CVS Drive, Woonsocket, RI 02895

Package Label

*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

TWIN PACK
NDC 69842-227-32

CVS

Original
STOMACH RELIEF
Bismuth Subsalicylate, 525 mg per 30 mL
Upset Stomach Reliever / Antidiarrheal

Compare to Pepto-Bismol® active ingredient*

Relieves 5 Symptoms: Heartburn, Indigestion, Upset stomach, Nausea, Diarrhea

Original Flavor **SHAKE WELL**

2 - 16 FL OZ (473 mL) BOTTLES
TOTAL 32 FL OZ (946 mL)

Distributed by:
 CVS Pharmacy, Inc.
 One CVS Drive
 Woodbridge, NJ 07095
 © 2026 CVS/Pharmacy
 CVS Store
 1-800-SHOP CVS
 V-10579
 100% money back
 guaranteed.
 cvs.com/returnpolicy

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL
 (MAYBE) IS NOT ON THIS CAP OR IS CRACKED OR MISSING.

#9900864
 0-50428-14032-6

PLASTIC BOTTLE
 PLASTIC CAP

Active Ingredient (in each 30 mL dose)	Purpose
Bismuth subsalicylate 525 mg	Upset stomach relief/antidiarrheal

Uses relieves ■ diarrhea ■ traveler's diarrhea
 ■ upset stomach due to overindulgence in food and drink, including
 ■ heartburn ■ gas ■ indigestion ■ nausea ■ bloating ■ belching

Warnings
 ■ **Key symptoms:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Shiga's syndrome, a rare but serious illness.
 ■ **Allergy alert:** Contains salicylate. Do not take if you are:
 ■ allergic to salicylates (including aspirin) ■ taking other salicylate products
 Do not use if you have: ■ an ulcer ■ a bleeding problem ■ bloody or black stool
 Ask a doctor before use if you have: ■ fever ■ muscle in the stool
 Ask a doctor or pharmacist before use if you are taking any drug for: ■ diabetes ■ gout ■ arthritis ■ anticoagulation (thinning the blood)
 When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.
 Stop use and ask a doctor if:
 ■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days
 If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children.
 In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions
 ■ **Use:** - milliliters ■ shake well before using
 ■ measure only with dosing cap provided. Do not use any other dosing device. ■ keep dosing cap with product
 ■ adults and children 12 years and over
 ■ 30 mL (1 dose) every 1/2 hour or 30 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
 ■ 30 mL (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
 ■ do not exceed 8 doses (240 mL) in 24 hours
 ■ use until diarrhea stops but not more than 2 days
 ■ children under 12 years of age, ask a doctor
 ■ store opens of clear fluids to help prevent dehydration caused by diarrhea

Other information
 ■ each 30 mL contains: potassium 8 mg, sodium 9 mg
 ■ each 30 mL contains: salicylate 205 mg
 ■ low sodium ■ keep tightly closed ■ protect from freezing ■ avoid excessive heat (over 104°F or 40°C)

Inactive ingredients benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucrose, xanthan gum

Questions or comments? call 1-877-733-3888 Monday-Friday 9AM-5PM EST

CVS Original Stomach Relief

STOMACH RELIEF REGULAR STRENGTH

bismuth subsalicylate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69842-227
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
BENZOIC ACID (UNII: 8SKNOB0MIM)	
D&C RED NO. 22 (UNII: 1678RKX8RT)	
D&C RED NO. 28 (UNII: 7671POY5NH)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842-227-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/31/2019	
2	NDC:69842-227-04	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/31/2019	

3	NDC:69842-227-16	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/31/2019	
4	NDC:69842-227-32	2 in 1 PACKAGE	01/31/2019	
4		473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	01/31/2019	

Labeler - CVS Pharmacy (062312574)

Revised: 11/2025

CVS Pharmacy