

**NIGHTTIME COLD AND FLU RELIEF MULTI SYMPTOM- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled**  
**United Natural Foods, Inc. dba UNFI**

-----  
**Drug Facts**

**Active ingredients (in each softgel)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

**Purposes**

**Pain reliever/fever reducer**

Cough suppressant

Antihistamine

**Uses**

- temporarily relieves common cold and flu symptoms
  - sore throat
  - headache
  - minor aches and pains
  - fever
  - runny nose and sneezing
  - cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult

a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### **When using this product**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- **when using other Daytime or Nighttime products, carefully read each label to insure correct dosing**

## Other information

- store between 15-30°C (59-86°F)
- avoid excessive heat

## Inactive ingredients

D&C yellow #10, FD&C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

## Questions or comments?

Call toll free **1-877-932-7948**

## Principal Display Panel

compare to Vicks® NyQuil® Cold & Flu LiquiCaps® active ingredients†

multi-symptom

Nighttime

### Cold & Flu Relief

acetaminophen (aches/fever reducer)

dextromethorphan HBr (cough suppressant)

doxylamine succinate (antihistamine)

relieves:

- aches, fever & sore throat
- cough, sneezing & runny nose

Softgels\*\* (\*\*liquid-filled capsules)

†This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, NyQuil®, and LiquiCaps® are registered trademarks of the Procter & Gamble Company.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS**

**TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

DISTRIBUTED BY SUPERVALU INC.,

EDEN PRAIRIE, MN 55344 USA

[supervaluprivatebrands.com](http://supervaluprivatebrands.com)

**Product Label**

**EQUALINE**<sup>®</sup>

NDC 41163-849-24

compare to Vicks<sup>®</sup>  
NyQuil<sup>®</sup> Cold & Flu  
LiquiCaps<sup>®</sup> active  
ingredients!

# multi-symptom nighttime cold & flu relief

acetaminophen (aches/fever reducer)  
dextromethorphan HBr (cough suppressant)  
doxylamine succinate (antihistamine)

relieves:

- aches, fever & sore throat
- cough, sneezing & runny nose



24 capsules

No. 3. Date:

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING. KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

<p><b>Drug Facts (continued)</b></p> <p><b>Warnings</b> ■ Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product ■ Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash ■ If a skin reaction occurs, stop use and seek medical help right away. ■ Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p><b>Do not use</b> ■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product ■ liver disease ■ glaucoma ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema ■ cough that occurs with too much phlegm (mucus)</p> <p><b>Ask a doctor or pharmacist before use if you are</b> ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers ■ avoid alcoholic drinks ■ marked drowsiness may occur ■ excitability may occur, especially in children ■ do not exceed recommended dosage</p> <p><b>When using this product</b></p>	<p><b>Drug Facts (continued)</b></p> <p><b>Directions</b> ■ do not take more than directed (see Overdose warning) ■ adults as well as for children even if you do not notice any signs or symptoms. ■ do not take more than 4 doses in 24 hours ■ adults and children 12 years and over: take 2 softgels with water every 6 hours ■ swallow whole; do not crush, chew, or dissolve ■ children under 12 years: do not use ■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing</p> <p><b>Other information</b> ■ store between 15-30°C (59-86°F) ■ avoid excessive heat</p> <p><b>Inactive ingredients</b> D&amp;C yellow #10, FD&amp;C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink</p> <p><b>Questions or comments?</b> Call toll free 1-877-932-7948</p>
<p><b>Drug Facts (continued)</b></p> <p><b>Stop use and ask a doctor if</b> ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ be careful when driving a motor vehicle or operating machinery</p> <p><b>Keep out of reach of children.</b> Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p><b>If pregnant or breast-feeding,</b> ask a health professional before use. These could be signs of a serious condition.</p> <p>■ cough comes back or occurs with rash or headache that lasts. ■ redness or swelling is present ■ new symptoms occur ■ fever gets worse or lasts more than 3 days ■ pain or cough gets worse or lasts more than 7 days ■ Stop use and ask a doctor if</p>	<p><b>Drug Facts</b></p> <p><b>Active ingredients (in each softgel)</b> Acetaminophen 325 mg.....Pain reliever/fever reducer Dextromethorphan HBr 15 mg.....Cough suppressant Doxylamine succinate 6.25 mg.....Antihistamine</p> <p><b>Purposes</b></p> <p><b>Uses</b> ■ temporarily relieves common cold and flu symptoms: ■ sore throat ■ headache ■ minor aches and pains ■ fever ■ runny nose and sneezing ■ cough due to minor throat and bronchial irritation</p> <p><b>Drug Facts (continued)</b></p>

**24 SOTTgeis** (\*\*liquid-filled capsules)

actual size

Lot Exp

DISTRIBUTED BY SUPERVALU INC., EDEN PRAIRIE, MN 55344 USA

\*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, NyQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.



Like it or let us make it right. That's our quality promise. 877-932-7948 [supervaluprivatebrands.com](http://supervaluprivatebrands.com)



**EQUALINE multi-Symptom Nighttime Cold & Flu Relief**

**NIGHTTIME COLD AND FLU RELIEF MULTI SYMPTOM**

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41163-849
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	

**Product Characteristics**

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	P30
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41163-849-24	24 in 1 CARTON	09/30/2019	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/30/2019	

**Labeler** - United Natural Foods, Inc. dba UNFI (943556183)

Revised: 5/2024

United Natural Foods, Inc. dba UNFI