NIGHTTIME COLD AND FLU RELIEF MULTI SYMPTOM- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled EQUALINE (SuperValu)

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

- temporarily relieves common cold and flu symptoms
 - sore throat
 - headache
 - minor aches and pains
 - fever
 - runny nose and sneezing
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not exceed recommended dosage
- excitability may occur, especially in children
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause liver damage.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- when using other Daytime or Nighttime products, carefully read each label to insure correct dosing

Other information

- store between 15-30°C (59-86°F)
- avoid excessive heat

Inactive ingredients

D&C yellow #10, FD&C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

Questions or comments?

Call toll free**1-877-932-7948**

Principal Display Panel

compare to Vicks® NyQuil® Cold & Flu LiquiCaps® active ingredients†

multi-symptom

Nighttime

Cold & Flu Relief

acetaminophen (aches/fever reducer)

dextromethorphan HBr (cough suppressant)

doxylamine succinate (antihistamine)

relieves:

- aches, fever & sore throat
- cough, sneezing & runny nose

Softgels** (**liquid-filled capsules)

†This product is not manufacture or distributed by The Procter & Gamble Company.

 $\mathsf{Vicks}\, \mathbbm{B}\,,\,\mathsf{NyQuil}\, \mathbbm{B}\,,\,\,\mathsf{and}\,\,\mathsf{LiquiCaps}\, \mathbbm{B}\,\,\mathsf{are}\,\,\mathsf{registered}\,\,\mathsf{trademarks}\,\,\mathsf{of}\,\,\mathsf{the}\,\,\mathsf{Procter}\,\,\&\,\,\mathsf{Gamble}\,\,\mathsf{Company}.$

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DISTRIBUTED BY SUPERVALU INC.,

EDEN PRAIRIE, MN 55344 USA

supervaluprivatebrands.com

Product Label

cough, sneezing & runny nose

- aches, fever & sore throat

- relieves:
- doxylamine succinate (antihistamine)

dextromethorphan HBr (cough suppressant)

nighttime cold & flu relief acetaminophen (aches/fever reducer)

compare to Vicks® NyQuil[®] Cold & Flu LiquiCaps[®] active ingredients⁺

EJUALINE[®]

 fevet fund the forming model and sneezing cough due to minor throat and bronchial irritation 	Actionation of the second seco	
Uses temporarily relieves common cold and flu symptoms: some sond pains some sond pains	Active ingredients (in each softgel) Active ingredients (in each softgel) Active ingredient and the soft soft soft soft soft soft soft soft	
(panuțuoo) รเวย₋₁ ธิกม ู		
		1

NDC 41163-849-24

Call toll tree 1-817-932-7948 Constitutions or comments?	When using this product a do not exceed recommended dosage excitability may occur, especially in children a svoid alcoholic drinks a marked drowsiness may occur to a state of the state
Inactive ingredients D&C yellow #10, FD&C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sophise schiftel white lok	Ask a doctor or pharmacist before use if you are taking the blood thinning drug wartarin taking sedatives or tranquilizers
Other information a store between 15-30°C (59-86°F) a svoid excessive heat	 cough that occurs with too much phiegm (mucus)
children under 12 years: do not use when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing	Ask a doctor before use if you have m liver disease m glaucoma trouble urinating due to an enlarged prostate gland a breathing problem or chronic cough that lasts such as occurs with
Directions a do not take more than directed (see Overdose warning) a do not take more than 4 doses in 24 hours a adults and children 12 years and over: take 2 softgels with water every 6 hours b swallow whole; do not crush, chew, or dissolve	If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Overdose warning: Taking more than the recommended dose can cause liver damage, in case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away, Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.	Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Stop use and ask a doctor if a pain or cough gets worse or lasts more than 7 days tever gets worse or lasts more than 3 days cough comes back or occurs with resh or headache that lasts. These could be signs of a serious condition.	may occur if you take: more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen a 3 or more alcoholic drinks every day while using this product Allergy alert. Acetaminophen may cause severe skin reactions. Symptoms may include: m skin redening mblisters mash if a skin reaction occurs, stop use and seek medical help right away. If a skin reaction occurs, stop use and seek medical help right away.
alcohol, sedatives, and tranquilizers may increase drowsiness be careful when driving a motor vehicle or operating machinery	Warnings Liver warning: This product contains acetaminophen. Severe liver damage
Drug Facts (continued)	Drug Facts (continued)

ER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN. BROKEN OR SHOWS ANY SIGNS OF TAMPERING.



EQUALINE multi-Symptom Nighttime Cold & Flu Relief

NIGHTTIME COLD AND FLU RELIEF MULTI SYMPTOM

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled

Product Type	HUMAN OTC DR	RUG I	tem Code (Source)	NDC:41	163-849
Route of Administration	n ORAL					
Active Ingredient/Act	tive Moiety					
Ir	ngredient Name			Basis of Str	rength	Strength
ACETAMINOPHEN (UNII: 362	O9ITL9D) (ACETAMINO	PHEN - UNII:3	362O9ITL9D)	ACETAMINOPHEN		325 mg
DEXTROMETHORPHAN HYD (DEXTROMETHORPHAN - UNII:		D2RTI9KYH)		DEXTROMETHORP HYDROBROMIDE	HAN	15 mg
DOXYLAMINE SUCCINATE (UNII:95QB77JKPL)	UNII: V9BI9B5YI2) (DO)	XYLAMINE -		DOXYLAMINE SUC	CINATE	6.25 mg
Inactive Ingredients						
	Ingredien	t Name				Strength
D&C YELLOW NO. 10 (UNII:	35SW5USQ3G)					
FD&C BLUE NO. 1 (UNII: H3	R47K3TBD)					
GELATIN (UNII: 2G86QN327L))					
GLYCERIN (UNII: PDC6A3C00)X)					
POLYETHYLENE GLYCOL, U	JNSPECIFIED (UNII: 3)	WJQ0SDW1A)				
POVIDONE (UNII: FZ989GH94	4E)					
PROPYLENE GLYCOL (UNII:	6DC9Q167V3)					
WATER (UNII: 059QF0KO0R)						
SORBITAN (UNII: 6092ICV9RI						
SORBITOL (UNII: 506T60A25	•					
MANNITOL (UNII: 30WL53L36	5A)					
Draduct Characterist						
Product Characterist		-				
Color	green	Score			no score	
Product Characterist Color Shape Flavor		Score Size Imprint			no score 20mm P30	

Pa	ckaging			
#	ltem Code	Package Description	Marketing Start Date	Marketing End Date
	NDC:41163-849- 24	24 in 1 CARTON	09/30/2019	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination		
		Product		
-		Product		
	arketing	Information		
	arketing Marketing Category		Marketing Start Date	Marketing End Date

Labeler - EQUALINE (SuperValu) (006961411)

Revised: 12/2022

EQUALINE (SuperValu)