

**IBUPROFEN- ibuprofen capsule, liquid filled  
AMERISOURCE BERGEN**

-----  
***Drug Facts***

***Active ingredient (in each capsule)***

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\*  
(present as the free acid and potassium salt)

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/Fever reducer

***Uses***

- temporarily relieves minor aches and pains due to:
- headache
- toothache
- backache
- menstrual cramps
- the common cold
- muscular aches
- minor pain of arthritis
- temporarily reduces fever

***Warnings***

**Allergy alert:**Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:**This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin,

ibuprofen, naproxen, or others]

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

- redness or swelling is present in the painful area
- any new symptoms appear

### **If pregnant or breast- feeding,**

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause delivery problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

### ***Directions***

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

### ***Other information***

- each capsule contains: **potassium 20 mg**
- read all warnings and directions before use
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F). Protect from light.

### ***Inactive ingredients***

FD&C Blue #1, gelatin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan and sorbitol

For India Manufacturing Site:

FD&C Blue #1, gelatin, polyethylene glycol, potassium hydroxide, purified water, sorbitan and sorbitol

### ***Questions or comments?***

call toll free **1-888-235-2466**

DO NOT USE IF TAMPER-EVIDENT SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED for YOUR PROTECTION" IS BROKEN OR MISSING.

†All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Advil® LIQUI-GELS®.

Distributed By

AmerisourceBergen

1 West First Ave

Conshohocken, PA 19428

Questions or Concerns?

R0422

L0000600

**For India Manufacturing Site:**

**MADE IN INDIA**

Code: TN/DRUGS/TN/25/00074

P3450/00/25

R0625

L0001063

**120's Bottle Label**

**Compare to Advil<sup>®</sup> Liqui-Gels<sup>®</sup>**

**active ingredient<sup>†</sup>**

**GOOD**

**NEIGHBOUR**

**PHARMACY<sup>®</sup>**

**NDC 46122-580-55**

**Ibuprofen**

**Capsules, 200 mg**

***Pain Reliever/Fever Reducer (NSAID)***

**120**

**softgels\*\***

**(\*\*Liquid filled capsules)**

**GOOD NEIGHBOR PHARMACY®**  
NDC 46122-580-55

**Ibuprofen Capsules, 200 mg**  
Pain Reliever/Fever Reducer (NSAID)

**120 softgels\*\*** (\*\*Liquid filled capsules)

Compare to Advil® Liqui-Gel® active ingredient

DO NOT USE IF TAMPER-EVIDENT SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED BY YOUR PROTECTION" IS BROKEN OR MISSING.

**Drug Facts**

**Active ingredient (in each capsule) Purpose**  
Sulbutozole (Ibuprofen) equivalent to 200 mg Ibuprofen (NSAID)\* Pain reliever/fever reducer (present as the free acid and potassium salt)  
\*non-steroidal anti-inflammatory drug

**Uses** ■ temporarily relieves minor aches and pains due to:  
■ headache ■ toothache ■ backache ■ menstrual cramps  
■ the common cold ■ muscular aches ■ minor pain of arthritis  
■ temporarily reduces fever

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
■ hives ■ facial swelling (wheezing) ■ shock  
■ skin redness/itching ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems. ■ take a blood thinner (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed  
**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

**Ask a doctor before use if** ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach

Distributed by  
Amgen Consumer  
1 World Center Ave.  
Conshohocken, PA 19428  
Questions or Comments?

ABC#: 10211368  
0 87701 43191 3  
Lot No: 14141  
10022  
L01000010

PEEL HERE

All trademarks are property of their respective owners. This product is not affiliated with the trademarks of Advil® LIQUI-GELS®.

Call toll-free 1-888-255-2466

**Questions or comments?**

**Drug Facts (continued)**

problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic.

**Ask a doctor or pharmacist before use if you are**  
■ under a doctor's care for any serious condition  
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin  
■ taking any other drug

**When using this product:**  
■ take with food or milk if stomach upset occurs

**Stop use and ask a doctor if** ■ you experience any of the following signs of stomach bleeding:  
■ feel tired ■ vomit blood ■ have bloody or black stools  
■ have stomach pain that does not get better  
■ you have symptoms of heart problems or stroke:  
■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling  
■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present in the painful area  
■ any new symptoms develop

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** ■ do not take more than directed  
■ the smallest effective dose should be used  
■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist. ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used  
■ do not exceed 6 capsules in 24 hours, unless directed by a doctor ■ children under 12 years: ask a doctor

**Other information** ■ each capsule contains:  
■ potassium 20 mg ■ read all warnings and directions before use ■ store at 20° to 25°C (68° to 77°F) ■ avoid excessive heat above 40°C (104°F). Protect from light.

**Inactive ingredients** FD &C Blue #1, gelatin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan and sorbitol

## 120's label-Made In India

Compare to Advil® Liqui-Gels®

active ingredient †

GOOD

NEIGHBOUR

PHARMACY®

NDC 46122-580-55

Ibuprofen

Capsules, 200 mg

Pain Reliever/Fever Reducer (NSAID)

120

softgels\*\*

(\*\*Liquid filled capsules)



Compare to Advil® Liqui-Gels® active ingredient!

NDC 46122-580-55

# Ibuprofen Capsules, 200 mg

Pain Reliever/Fever Reducer (NSAID)

120 softgels\*\* (\*\*Liquid filled capsules)



DO NOT USE IF TAMPER-EVIDENT SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED BY YOUR PROTECTION" IS BROKEN OR UNUSUAL.

## Drug Facts

**Active ingredient (in each capsule)** Purpose  
Sulfonized ibuprofen equal to 200 mg of ibuprofen (NSAID)\*. Pain reliever/fever reducer (present as the free acid and potassium salt) nonsteroidal anti-inflammatory drug

**Uses**  
temporarily relieves minor aches and pains due to:  
headache toothache backache menstrual cramps  
the common cold muscular aches minor pain of arthritis  
temporarily reduces fever

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:  
hives facial swelling asthma (wheezing) shock  
skin redness rash ulcers. If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
are age 60 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer right before or after heart surgery  
**Ask a doctor before use if** stomach bleeding warning applies to you you have problems or serious side effects from taking pain relievers or fever reducers you have a history of stomach pain relievers or fever reducers

Rev 07/2025 ABCR: 10300111  
Distributed By: AmriseSourceBergien West First PA 15428  
www.mygnph.com Questions or Concerns?  
Code: TN / DRUGS / TN25/00074 3 46122-5805517  
Lot No: Exp. Date:  
MADE IN INDIA

Non Varnish Area



T00073R0125

This product is not manufactured or distributed by the owners of Advil® (LDA) 12/13/2017.

## Questions or comments?

**Inactive ingredients:** FD&C Blue #1, gelatin, polyethylene glycol, potassium hydroxide, purified water, sorbitan and sorbitol

**Other information** each capsule contains:  
potassium 20 mg read all warnings and directions before use store at 20° to 25°C (68° to 77°F) avoid excessive heat above 40°C (104°F). Protect from light.

**Directions** do not take more than directed  
the smallest effective dose should be used  
adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist if pain or fever does not respond to 1 capsule, 2 capsules may be used do not exceed 6 capsules in 24 hours, unless directed by a doctor children under 12 years: ask a doctor

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**When using this product**  
take with food or milk if stomach upset occurs

**Stop use and ask a doctor if** you experience any of the following signs of stomach bleeding:  
feel faint vomit blood have bloody or black stools have stomach pain that does not get better

you have symptoms of heart problems or stroke:  
chest pain trouble breathing weakness in one part or side of body slurred speech leg swelling pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days redness or swelling is present in the painful area any new symptoms appear

**Ask a doctor or pharmacist before use if you are** under a doctor's care for any serious condition taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin taking any other drug

**Drug Facts (continued)**  
problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke you are taking a diuretic

**When using this product**  
take with food or milk if stomach upset occurs

**Stop use and ask a doctor if** you experience any of the following signs of stomach bleeding:  
feel faint vomit blood have bloody or black stools have stomach pain that does not get better

you have symptoms of heart problems or stroke:  
chest pain trouble breathing weakness in one part or side of body slurred speech leg swelling pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days redness or swelling is present in the painful area any new symptoms appear

## IBUPROFEN

ibuprofen capsule, liquid filled

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:46122-580
Route of Administration	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

## Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDWIA)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
WATER (UNII: 059QF0KO0R)	

**SORBITAN** (UNII: 6O92ICV9RU)

**SORBITOL** (UNII: 506T60A25R)

### Product Characteristics

<b>Color</b>	blue	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	IB1
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46122-580-55	120 in 1 BOTTLE; Type 0: Not a Combination Product	03/01/2019	
2	NDC:46122-580-58	40 in 1 CARTON	09/04/2019	
2		40 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:46122-580-41	80 in 1 CARTON	09/04/2019	
3		80 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078682	03/01/2019	

**Labeler** - AMERISOURCE BERGEN (007914906)

**Registrant** - Bionpharma Inc. (079637826)

### Establishment

Name	Address	ID/FEI	Business Operations
Patheon Softgels Inc.		002193829	manufacture(46122-580)

### Establishment

Name	Address	ID/FEI	Business Operations
ESJAY PHARMA PRIVATE LIMITED		957309497	manufacture(46122-580)