RVLAIM SILICONE SCAR CREAM- allantoin cream Shenzhen Jiucai Ecommerce Co., Ltd

RVLAIM SILICONE SCAR CREAM

RVLAIM SILICONE SCAR CREAM

Allantoin 0.5%

Emollient Scar Removal

This product can be used day and night, and thescar should be kept clean and dry before use.

Squeeze out the appropriate amount according to the size of the scar and massage gently until thescar cream is absorbed and does not feel shiny, wet, or sticky. Use 2-3 times a day, increasing or decreasing the frequency depending on the condition of the scar. Individual results will be different, and the results will take time to verify.

For external use only, avoid contact with the eyesif symptoms of redness or allergy occur, discontinue use and consuit a physician.

if you are using other medications or treatmentsespecially those for skin problems. If you have a history of allergies, especially toingredients in medications, cosmetics or any otherproducts.

If you are pregnant, breastfeeding, a child or anelderly person, or if you have any chronic health problems.

After consulting a doctor, they can provide you withprofessional advice on the use of this product basedon your specific situation and needs.

Preparation:

Make sure the scar area is clean and dry. Wash your hands before use to ensure that your hands are clean.

How to use:

Gently squeeze out an appropriate amount of scarcream to avoid overdose. Applythe scar cream evenlyon the scar area. Massage for a while to help the product absorb better.

Frequency ofuse:

Use [2-3 times a day] according to the productinstructions or doctor's advice. It is recommended to apply the scar cream after it is completely absorbed before the next application.

Precautions:

For external use only, do not take internally. Avoid contact with eyes and mucous membranes. If allergic symptoms such as skin irritation, redness, swelling, etc. occur, please stop using it immediately and consult a doctor. Pregnant women, breastfeeding women and children please consult a doctor before use.

Use in special circumstances:

Do not use this product on open wounds or broken skin. Hard scars are difficult to treat, so it is recommended to consult a doctor before use.

Storage:

Store in a cool and dry place, avoid direct sunlight. Keep this product out of the reach ofchildren.

Important tips:

Before using this product, please read the productins tructions carefully and follow the advice of your doctor or professional.

if symptoms of redness or allergy occur, discontinueuse and consult a physician.

KEEP OUT OF REACH OF CHILDREN SECTION

Use 2-3 times a day, increasing or decreasing the frequency depending on the condition of the scar. Individual results will be different, and the results will take time to verify.

Store in a dark and cool place

WATER

Glycerin

Allantoin

Retinol

Panthenol

Vitamin C

Vitamin E

DIMETHICONE

Carbomer

Laurocapram

Propylene Glycol

Caprylic/capric triglyceride

Stearic Acid

Isopropyl myristate

SODIUM CARBOXYMETHYL .BETA.-GLUCAN

Centella Asiatica Extract

包装刀模: 38*38*128mm



RVLAIM SILICONE SCAR CREAM

allantoin cream

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:84433-001	
Route of Administration	TOPICAL			

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
ALLANTOIN (UNII: 344S277G0Z) (ALLANTOIN - UNII:344S277G0Z)	ALLANTOIN	5 g in 100 mL		

Inactive Ingredients	
Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
ALPHA-TOCOPHEROL (UNII: H4N855PNZ1)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
ASCORBIC ACID (UNII: PQ6CK8PD0R)	
CARBOMER HOMOPOLYMER, UNSPECIFIED TYPE (UNII: 0A5MM307FC)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
LAUROCAPRAM (UNII: 1F3X9DRV9X)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
GLYCERIN (UNII: PDC6A3C0OX)	
PANTHENOL (UNII: WW9CM0O67Z)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM CARBOXYMETHYL .BETAGLUCAN (DS 0.65-0.85) (UNII: 2YGO1190AP)	
CENTELLA ASIATICA TRITERPENOIDS (UNII: 4YS74Q4G4J)	
RETINOL (UNII: G2SH0XKK91)	
WATER (UNII: 059QF0KO0R)	

ı	Packaging				
	#	# Item Code Package Description		Marketing Start Date	Marketing End Date
		NDC:84433-001- 01	50 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/19/2024	

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M016	06/19/2024		

Labeler - Shenzhen Jiucai Ecommerce Co., Ltd (621699187)

Establishment				
Name	Address	ID/FEI	Business Operations	
Shenzhen Jiucai Ecommerce Co., Ltd		621699187	manufacture(84433-001)	

Revised: 6/2024 Shenzhen Jiucai Ecommerce Co., Ltd