VICKS DAYQUIL AND VICKS NYQUIL ELDERBERRY COLD AND FLUacetaminophen, dextromethorphan hbr, doxylamine succinate The Procter & Gamble Manufacturing Company

VICKS [®] DayQuil™ NyQuil™ ELDERBERRY COLD & FLU Liquid Convenience Pack

VICKS ® DayQuil™ ELDERBERRY COLD & FLU

Drug Facts

Active ingredients (in each 15 mL)

Acetaminophen 325 mg Dextromethorphan HBr 10 mg

Purpose

Pain reliever/ fever reducer Cough suppressant

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product
- child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 15 mL contains: sodium 13 mg
- store at no greater than 25°C and do not refrigerate

Inactive ingredients

citric acid, FD&C Blue No. 1, FD&C Red No. 40, flavor (with elderberry), glycerin, propylene glycol, saccharin sodium, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate, water, xanthan gum

Questions?

1-800-362-1683

TAMPER EVIDENT: DO NOT USE IF PRINTED SHRINKBAND ON BOTTLE IS BROKEN OR MISSING.

DIST. BY: PROCTER & GAMBLE,

CINCINNATI, OH 45202

VICKS® NyQuil™ ELDERBERRY COLD & FLU

Drug Facts

Active ingredients (in each 15 mL)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Doxylamine succinate 6.25 mg

Purpose

Pain reliever/Fever reducer

Cough suppressant

Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever
- runny nose & sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product
- child takes more than 4 doses (15 mL eah) in 24 hrs, which is the maximum daily amount for this product
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy Alert:Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning:If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked dowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough get worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 15 mL contains: sodium 13 mg
- store at no greater than 25°C and do not refrigerate

Inactive ingredients

citric acid, FD&C Blue No. 1, FD&C Red No. 40, flavor (with elderberry), glycerin, propylene glycol, saccharin sodium, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate, water, xanthan gum

Questions?

1-800-362-1683

TAMPER EVIDENT: DO NOT USE IF PRINTED SHRINKBAND ON BOTTLE IS BROKEN OR MISSING.

DIST. BY: PROCTER & GAMBLE,

CINCINNATI, OH 45202

PRINCIPAL DISPLAY PANEL - Kit Carton

POWERFUL RELIEF

BLACK ELDERBERRY FLAVORED

Vicks DayQuil and NyQuil ELDERBERRY

COLD & FLU

Vicks DayQuil ELDERBERRY

COLD & FLU

THE DAYTIME, COUGHING, ACHING, SORE THROAT, FEVER, POWER THROUGH YOUR DAY, MEDICINE.

Acetaminophen, Dextromethorphan HBr

Headache, Fever, Sore Throat,

Minor Aches & Pains

Cough

Vicks NyQuil ELDERBERRY

COLD & FLU

THE NIGHTTIME, SNIFFLING, SNEEZING, COUGHING, ACHING, FEVER, BEST SLEEP WITH A COLD, MEDICINE.

Acetaminophen, Dextromethorphan HBr, Doxylamine succinate

Headache, Fever, Sore Throat, minor Aches & Pains

Sneezing, Runny Nose

Cough

2 BOTTLES - 12 FL OZ (354 mL) EACH; TOTAL 24 FL OZ (708 mL)



VICKS DAYQUIL AND VICKS NYQUIL ELDERBERRY COLD AND FLU

acetaminophen, dextromethorphan hbr, doxylamine succinate kit

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:84126-309

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84126-309- 24	1 in 1 PACKAGE; Type 0: Not a Combination Product	07/09/2024	

_			P
_	uantity	ΔT	Darte
•	ualitity	UI.	raits

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE, PLASTIC	354 mL
Part 2	1 BOTTLE, PLASTIC	354 mL

Part 1 of 2

VICKS DAYQUIL ELDERBERRY COLD AND FLU

acetaminophen, dextromethorphan hbr liquid

Product Information

Item Code (Source) NDC:84126-307

Route of Administration ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg in 15 mL	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL	

Inactive Ingredients			
Ingredient Name	Strength		
XANTHAN GUM (UNII: TTV12P4NEE)			
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)			
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			
GLYCERIN (UNII: PDC6A3C0OX)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
WATER (UNII: 059QF0KO0R)			
SACCHARIN SODIUM (UNII: SB8ZUX40TY)			
SODIUM BENZOATE (UNII: OJ245FE5EU)			
SODIUM CITRATE (UNII: 1Q73Q2JULR)			
SORBITOL (UNII: 506T60A25R)			
SUCRALOSE (UNII: 96K6UQ3ZD4)			

Product Characteristics			
Color	purple	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

nDate Date Date Date Date	Packaging			
NDC:84126- 354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a	# Item Code	Package Description	_	Marketing End Date
507 IZ	1 NDC:84126- 307-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing InformationMarketing CategoryApplication Number or Monograph CitationMarketing Start DateMarketing End DateOTC Monograph DrugM01207/09/2024

Part 2 of 2

VICKS NYQUIL ELDERBERRY COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate liquid

Product Information	
Item Code (Source)	NDC:84126-308
Route of Administration	ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg in 15 mL	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL	
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg in 15 mL	

Inactive Ingredients			
Ingredient Name	Strength		
SUCRALOSE (UNII: 96K6UQ3ZD4)			
SORBITOL (UNII: 506T60A25R)			
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)			
FD&C RED NO. 40 (UNII: WZB9127XOA)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
WATER (UNII: 059QF0KO0R)			
SACCHARIN SODIUM (UNII: SB8ZUX40TY)			
SODIUM BENZOATE (UNII: OJ245FE5EU)			
SODIUM CITRATE (UNII: 1Q73Q2JULR)			
XANTHAN GUM (UNII: TTV12P4NEE)			
GLYCERIN (UNII: PDC6A3C0OX)			

Product Characteristics			
Color	purple	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84126- 308-12	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information				
Marketing Category	• • • • • • • • • • • • • • • • • • • •		Marketing End Date	
OTC Monograph Drug	M012	07/09/2024		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/09/2024	

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Establishment				
Name	Address	ID/FEI	Business Operations	
Novacyl Wuxi Pharmaceutical Co., Ltd.		263624659	api manufacture(84126-308)	

Revised: 6/2024 The Procter & Gamble Manufacturing Company