CHILDRENS IBUPROFEN- ibuprofen tablet, chewable Kroger Company

Kroger Co. Children's Ibuprofen Drug Facts

Active ingredient (in each chewable tablet)

Ibuprofen 100 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- · has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more

than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- · child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- · child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- mouth or throat burning may occur; give with food or water
- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
- feels faint
- vomits blood
- has bloody or black stools
- has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days

- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- chew or crush tablets completely before swallowing
- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day

Dosing Chart			
Weight (lb)	Age (yr)	Tablets	
under 24	under 2	ask a doctor	
24-35	2-3	1	
36-47	4-5	1 ½	
48-59	6-8	2	
60-71	9-10	2 ½	
72-95	11	3	

Other information

- phenylketonurics: contains phenylalanine 6 mg per tablet
- store between 20-25°C (68-77°F)
- do not use if printed seal under cap is broken or missing

Inactive ingredients

acesulfame potassium, ammonium glycyrrhizin, aspartame, carnauba wax, croscarmellose sodium, D&C red no. 27 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, mannitol, natural and artificial flavors, silicon dioxide, sodium lauryl sulfate, soybean oil, succinic acid

Questions or comments?

1-800-632-6900

Package/Label Principal Display Panel

COMPARE TO the active ingredient of CHILDREN'S MOTRIN® See side panel OUR PHARMACIST RECOMMENDED

Ages 2-11 Years

children's Ibuprofen

Chewable Tablets, 100 mg

Pain Reliever/Fever Reducer (NSAID)

actual size

Grape Flavor

Chew or crush tablets completely before swallowing

CHEWABLES

Lasts up to 8 Hours

24 TABLETS





KEEP CARTON FOR REFERENCE

is not affiliated with I he Kroger Co. or this product. "Motrin" is a registered trademark of Johnson & Johnson Johnson & Johnson

Drug Facts

Active ingredient (in each chewable tablet) Purposes buprofen 100 m.g (NSAID)*..... *nonsteroidal anti-inflammatory drug

Uses temporarily: ■ reduces fever
■ relieves minor aches and pains due to the common cold, flu, sore
• fireat, headaches and toofbaches

Warni ngs

Marrii ngs
Allergy a lert lbupofen may cause a severe allergic reaction,
especially inpeople allergic to apprin. Symptoms may include:

It hives If facial swelling If as thima (wheezing)
If shock If skin reddening If rash If blates
If an allergic reaction occurs, dop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSMD, which
may cause severe a somach bleeding. The chance is higher if your child:
If has had a tomach ubers or bleeding problems
If takes althood thinning (annicosqualming or sheroid drug
If takes other druss containing usescription or nonnesscription

Takes other drugs containing prescription or nonprescription
NS AIDs (aspirin, ibuprofen, naproxen, or others)

Takes more or for a longer time than directed
Heart a ttack and stroke warning: NSA Ds, except aspirin, increase
the risk of heart attack, heart failure, and stroke. These can be fatal. The
risk is higher if you use more than directed or for longer than directed. Sore throat warning: Severe or persistent one throat or so se throat accompanied by high even headache, nausea, and wontiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Drug Facts (continued)

Do notuse

■ if the child has ever had an allergic reaction to ibuprofenor any other pain reliever/fever reducer

■ right before or after heart surgery

Ask a doctor before use if

Ask a doctor be fore use if

stomach bleeding warning applies to your child

child has a history of stom ash problems, such as hearthum

child has problems or serious side effects from taking pain relievers
or feer reducers

child has notbeen drinking fluids

child has notbeen drinking fluids

child has best a bt of fluid due to vomiting or diarrhea.

child has high bood pressure, heart disease, liver cirrhosis, kidney

disease schedus at the decided.

disease, or had a stroke ■child has asthma ■child is taking a diuret ic

Ask a doctor or pharmacist before use if the child is

■ under a doctor's care for any serious condition ■taking any other drug

When using this product ■ mouth or throat buming may occur; give with food or water ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if

■child experiences any of the following signs of stormach bleeding: ■feels faint ■vormits blood ■ has bloody or black stools

■ has stomach pain that does not getbetter
■ child has symptoms of heart problems or stroke
■ chest pain ■ trouble breathing

■ weakness in one part or side of body ■ slurred speech ■ leg swelling

Drug Facts (continued)

■ the child does not get any relief within first day (24 hours) of treatment ■ fever or pain gets worse or lasts more than 3 days

redness or swelling is present in the painful area.

■any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contacta Poison Control Center right away. (1-800-222-1222)

Directions

٠

Chewor crush tablets completely before swallowing
 this product does not contain directions or complete warnings
 for adult use

■ donot give more than directed

■find right dose on chart below. If possible, use weight to dose;

otherwise use age. ■ if needed, repeat dose every 6-8 hours ■ do not use more than 4 times a day

Dosing Chart

Weight (lb)	Age (yr)	Tablets
under 24	under 2	aska doctor
24-35	2-3	1
36-47	4-5	1 1/2
48 - 59	6-8	2
60-71	9-10	21/2
72-95	11	3

hacthe ingredients aceateme passium, annoi um giopritich, aspatame, carautha wa, o ocamelbesectium, D& et oc. 27 duminum bke, FD& the na 1 duminum bke, hpromstoe, magresium starate, mannto, natra alandar fitidal flavos, stomotoris, sodium tarris utate, sopeanol, socializaci Other information

phenylectonuries:combins phenyletanine 6 mg pertablet

steine between 20-25 °C(68-77°F)

do not use if printed seal under cap is broken o rmissi gund Facts (continued OHO www.kroger.com 21 40046 32499 00 GLUTEN FREE



CHILDRENS IBUPROFEN

ibuprofen tablet, chewable

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:30142-402

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	100 mg

Inactive Ingredients	
Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 230V73Q5G9)	
AMMONIUM GLYCYRRHIZATE (UNII: 3VRD35U26C)	
ASPARTAME (UNII: Z0H242BBR1)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
CROSCARMELLOSE SODIUM (UNII: M280L1HH48)	
D&C RED NO. 27 (UNII: 2LRS185U6K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MANNITOL (UNII: 30WL53L36A)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SOYBEAN OIL (UNII: 241ATL177A)	
SUCCINIC ACID (UNII: AB6MNQ6J6L)	

Product Characteristics				
Color	PURPLE (Lavender)	Score	2 pieces	
Shape	ROUND	Size	12mm	
Flavor	GRAPE	Imprint Code	L521	
Contains				

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:30142-402- 62	1 in 1 CARTON	10/09/2019		
1		24 in 1 BOTTLE; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076359	10/09/2019	

Labeler - Kroger Company (006999528)

Revised: 10/2022 Kroger Company